

thm\_2Ereal\_\_topology\_2EBILINEAR\_\_RADD  
(TMKDRVYZmhFzaza-  
MjD8h2LvEXmZtUv2vjaM)

October 26, 2020

**Definition 1** We define  $c\_2Emin\_2E\_3D$  to be  $\lambda A.\lambda x \in A.\lambda y \in A.inj\_o (x = y)$  of type  $\iota \Rightarrow \iota$ .

**Definition 2** We define  $c\_2Ebool\_2E\_21$  to be  $(ap (ap (c\_2Emin\_2E\_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

**Definition 3** We define  $c\_2Ebool\_2E\_21$  to be  $\lambda A.27a : \iota.(\lambda V0P \in (2^{A-27a}).(ap (ap (c\_2Emin\_2E\_3D (2^{A-27a}))$

**Definition 4** We define  $c\_2Ebool\_2E\_21$  to be  $(ap (c\_2Ebool\_2E\_21 2) (\lambda V0t \in 2.V0t))$ .

**Definition 5** We define  $c\_2Emin\_2E\_3D\_3D\_3E$  to be  $\lambda P \in 2.\lambda Q \in 2.inj\_o (p P \Rightarrow p Q)$  of type  $\iota$ .

**Definition 6** We define  $c\_2Ebool\_2E\_7E$  to be  $(\lambda V0t \in 2.(ap (ap c\_2Emin\_2E\_3D\_3D\_3E V0t) c\_2Ebool\_2E\_21))$

Let  $ty\_2Ehreal\_2Ehreal : \iota$  be given. Assume the following.

$$nonempty\ ty\_2Ehreal\_2Ehreal \tag{1}$$

Let  $ty\_2Epair\_2Eprod : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A0.nonempty\ A0 \Rightarrow \forall A1.nonempty\ A1 \Rightarrow nonempty\ (ty\_2Epair\_2Eprod\ A0\ A1) \tag{2}$$

Let  $ty\_2Erealax\_2Ereal : \iota$  be given. Assume the following.

$$nonempty\ ty\_2Erealax\_2Ereal \tag{3}$$

Let  $c\_2Erealax\_2Ereal\_REP\_CLASS : \iota$  be given. Assume the following.

$$c\_2Erealax\_2Ereal\_REP\_CLASS \in ((2^{(ty\_2Epair\_2Eprod\ ty\_2Ehreal\_2Ehreal\ ty\_2Ehreal\_2Ehreal)})\ ty\_2Erealax) \tag{4}$$

**Definition 7** We define  $c\_2Emin\_2E\_40$  to be  $\lambda A.\lambda P \in 2^A.if (\exists x \in A.p (ap P x))$  **then** (the  $(\lambda x.x \in A \wedge p x)$ ) of type  $\iota \Rightarrow \iota$ .



Assume the following.

$$\forall A\_27a.nonempty\ A\_27a \Rightarrow (\forall V0x \in A\_27a. (\forall V1y \in A\_27a. ((V0x = V1y) \Leftrightarrow (V1y = V0x)))) \quad (13)$$

Assume the following.

$$(\forall V0t1 \in 2. (\forall V1t2 \in 2. (\forall V2t3 \in 2. (((p\ V0t1) \Rightarrow ((p\ V1t2) \Rightarrow (p\ V2t3))) \Leftrightarrow (((p\ V0t1) \wedge (p\ V1t2)) \Rightarrow (p\ V2t3)))))) \quad (14)$$

Assume the following.

$$\begin{aligned} & (\forall V0x \in 2. (\forall V1x\_27 \in 2. (\forall V2y \in 2. (\forall V3y\_27 \in \\ & 2. (((((p\ V0x) \Leftrightarrow (p\ V1x\_27)) \wedge ((p\ V1x\_27) \Rightarrow ((p\ V2y) \Leftrightarrow (p\ V3y\_27)))) \Rightarrow \\ & (((p\ V0x) \Rightarrow (p\ V2y)) \Leftrightarrow ((p\ V1x\_27) \Rightarrow (p\ V3y\_27)))))))) \end{aligned} \quad (15)$$

**Theorem 1**

$$\begin{aligned} & (\forall V0h \in ((ty\_2Erealax\_2Ereal^{ty\_2Erealax\_2Ereal})^{ty\_2Erealax\_2Ereal}). \\ & (\forall V1x \in ty\_2Erealax\_2Ereal. (\forall V2y \in ty\_2Erealax\_2Ereal. \\ & (\forall V3z \in ty\_2Erealax\_2Ereal. ((p\ (ap\ c\_2Ereal\_topology\_2Ebilinear \\ & V0h)) \Rightarrow ((ap\ (ap\ V0h\ V1x)\ (ap\ (ap\ c\_2Erealax\_2Ereal\_add\ V2y)\ V3z)) = \\ & (ap\ (ap\ c\_2Erealax\_2Ereal\_add\ (ap\ (ap\ V0h\ V1x)\ V2y))\ (ap\ (ap\ V0h \\ & V1x)\ V3z)))))))) \end{aligned}$$