

thm_2Ereal__topology_2ECARD__EQ__REAL__IMP__UNCOUNTAI (TMMR8ALtfKqNjLxpuCcvGe1XuLAYdnbdifR)

October 26, 2020

Definition 1 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$
of type $\iota \Rightarrow \iota$.

Definition 2 We define c_2Ebool_2ET to be $(ap (ap (c_2Emin_2E_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

Definition 3 We define c_2Ebool_2EIN to be $\lambda A_27a : \iota.(\lambda V0x \in A_27a.(\lambda V1f \in (2^{A_27a}).(ap V1f V0x)))$

Definition 4 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p P \Rightarrow p Q)$
of type ι .

Definition 5 We define $c_2Ebool_2E_21$ to be $\lambda A_27a : \iota.(\lambda V0P \in (2^{A_27a}).(ap (ap (c_2Emin_2E_3D (2^{A_27a})))$

Definition 6 We define $c_2Ebool_2E_2F_5C$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21 2)) (\lambda V2t \in 2.V2t)))$

Definition 7 We define $c_2Emin_2E_40$ to be $\lambda A.\lambda P \in 2^A.if (\exists x \in A.p (ap P x)) \mathbf{then} (the (\lambda x.x \in A \wedge p x))$
of type $\iota \Rightarrow \iota$.

Definition 8 We define $c_2Ebool_2E_3F$ to be $\lambda A_27a : \iota.(\lambda V0P \in (2^{A_27a}).(ap V0P (ap (c_2Emin_2E_40 A_27a P))))$

Definition 9 We define $c_2Epred_set_2ESURJ$ to be $\lambda A_27a : \iota.\lambda A_27b : \iota.\lambda V0f \in (A_27b^{A_27a}).\lambda V1s \in (2^{A_27a})$

Definition 10 We define $c_2Epred_set_2EINJ$ to be $\lambda A_27a : \iota.\lambda A_27b : \iota.\lambda V0f \in (A_27b^{A_27a}).\lambda V1s \in (2^{A_27a})$

Definition 11 We define $c_2Epred_set_2EBIJ$ to be $\lambda A_27a : \iota.\lambda A_27b : \iota.\lambda V0f \in (A_27b^{A_27a}).\lambda V1s \in (2^{A_27a})$

Definition 12 We define $c_2Ecardinal_2Ecardeq$ to be $\lambda A_27a : \iota.\lambda A_27b : \iota.\lambda V0s1 \in (2^{A_27a}).\lambda V1s2 \in (2^{A_27a})$

Definition 13 We define $c_2Epred_set_2EUNIV$ to be $\lambda A_27a : \iota.(\lambda V0x \in A_27a.c_2Ebool_2ET)$.

Let $ty_2Erealax_2Ereal : \iota$ be given. Assume the following.

$$nonempty\ ty_2Erealax_2Ereal \tag{1}$$

Let $ty_2Enum_2Enum : \iota$ be given. Assume the following.

$$nonempty\ ty_2Enum_2Enum \tag{2}$$

Definition 14 We define $c_2Epred_set_2Ecountable$ to be $\lambda A.27a : \iota.\lambda V0s \in (2^{A.27a}).(ap (c_2Ebool_2E_3F$

Definition 15 We define c_2Ebool_2EF to be $(ap (c_2Ebool_2E.21 2) (\lambda V0t \in 2.V0t))$.

Definition 16 We define $c_2Ebool_2E.5C.2F$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E.21 2) (\lambda V2t \in$

Definition 17 We define $c_2Ebool_2E.7E$ to be $(\lambda V0t \in 2.(ap (ap c_2Emin_2E.3D.3D.3E V0t) c_2Ebool_2E$

Assume the following.

$$True \tag{3}$$

Assume the following.

$$(\forall V0t1 \in 2.(\forall V1t2 \in 2.(((p V0t1) \Rightarrow (p V1t2)) \Rightarrow (((p V1t2) \Rightarrow (p V0t1)) \Rightarrow ((p V0t1) \Leftrightarrow (p V1t2)))))) \tag{4}$$

Assume the following.

$$(\forall V0t \in 2.(False \Rightarrow (p V0t))) \tag{5}$$

Assume the following.

$$(\forall V0t \in 2.(((p V0t) \Rightarrow False) \Rightarrow (\neg(p V0t)))) \tag{6}$$

Assume the following.

$$(\forall V0t \in 2.((\neg(p V0t)) \Rightarrow ((p V0t) \Rightarrow False))) \tag{7}$$

Assume the following.

$$(\forall V0t \in 2.(((True \Rightarrow (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \Rightarrow True) \Leftrightarrow True) \wedge (((False \Rightarrow (p V0t)) \Leftrightarrow True) \wedge (((p V0t) \Rightarrow (p V0t)) \Leftrightarrow True) \wedge ((p V0t) \Rightarrow False) \Leftrightarrow (\neg(p V0t)))))) \tag{8}$$

Assume the following.

$$((\forall V0t \in 2.((\neg(\neg(p V0t))) \Leftrightarrow (p V0t))) \wedge ((\neg True) \Leftrightarrow False) \wedge ((\neg False) \Leftrightarrow True)) \tag{9}$$

Assume the following.

$$(\forall V0t \in 2.(((True \Leftrightarrow (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \Leftrightarrow True) \Leftrightarrow (p V0t)) \wedge (((False \Leftrightarrow (p V0t)) \Leftrightarrow (\neg(p V0t))) \wedge (((p V0t) \Leftrightarrow False) \Leftrightarrow (\neg(p V0t)))))) \tag{10}$$

Assume the following.

$$(\forall V0A \in 2.(\forall V1B \in 2.(((p V0A) \vee (p V1B)) \Leftrightarrow ((p V1B) \vee (p V0A)))) \tag{11}$$

Assume the following.

$$(\forall V0r \in 2. (\forall V1p \in 2. (\forall V2q \in 2. (((p V1p) \wedge (p V2q) \Rightarrow (p V0r)) \Leftrightarrow ((p V1p) \Rightarrow ((p V2q) \Rightarrow (p V0r))))))) \quad (12)$$

Assume the following.

$$\forall A.27a.nonempty A.27a \Rightarrow \forall A.27b.nonempty A.27b \Rightarrow (\forall V0s \in (2^{A.27a}). (\forall V1t \in (2^{A.27b}). ((p (ap (ap (c.2Ecardinal.2Ecardeq A.27a A.27b) V0s) V1t)) \Leftrightarrow (p (ap (ap (c.2Ecardinal.2Ecardeq A.27b A.27a) V1t) V0s)))))) \quad (13)$$

Assume the following.

$$\forall A.27a.nonempty A.27a \Rightarrow (\forall V0s \in (2^{A.27a}). (\forall V1t \in (2^{A.27a}). (((p (ap (c.2Epred_set.2Ecountable A.27a) V1t)) \wedge (p (ap (ap (c.2Ecardinal.2Ecardeq A.27a A.27a) V0s) V1t))) \Rightarrow (p (ap (c.2Epred_set.2Ecountable A.27a) V0s)))))) \quad (14)$$

Assume the following.

$$(\neg (p (ap (c.2Epred_set.2Ecountable ty_2Erealax.2Ereal) (c.2Epred_set.2EUNIV ty_2Erealax.2Ereal)))) \quad (15)$$

Assume the following.

$$(\forall V0t \in 2. ((\neg(\neg(p V0t))) \Leftrightarrow (p V0t))) \quad (16)$$

Assume the following.

$$(\forall V0A \in 2. ((p V0A) \Rightarrow ((\neg(p V0A)) \Rightarrow False))) \quad (17)$$

Assume the following.

$$(\forall V0A \in 2. (\forall V1B \in 2. (((\neg((p V0A) \vee (p V1B))) \Rightarrow False) \Leftrightarrow (((p V0A) \Rightarrow False) \Rightarrow ((\neg(p V1B)) \Rightarrow False)))) \quad (18)$$

Assume the following.

$$(\forall V0A \in 2. (\forall V1B \in 2. (((\neg((\neg(p V0A)) \vee (p V1B))) \Rightarrow False) \Leftrightarrow ((p V0A) \Rightarrow ((\neg(p V1B)) \Rightarrow False)))) \quad (19)$$

Assume the following.

$$(\forall V0A \in 2. (((\neg(p V0A)) \Rightarrow False) \Rightarrow (((p V0A) \Rightarrow False) \Rightarrow False))) \quad (20)$$

Assume the following.

$$(\forall V0p \in 2. (\forall V1q \in 2. (\forall V2r \in 2. (((p V0p) \Leftrightarrow ((p V1q) \Leftrightarrow (p V2r))) \Leftrightarrow (((p V0p) \vee ((p V1q) \vee (p V2r))) \wedge (((p V0p) \vee ((\neg(p V2r)) \vee (\neg(p V1q)))) \wedge (((p V1q) \vee ((\neg(p V2r)) \vee (\neg(p V0p)))) \wedge ((p V2r) \vee ((\neg(p V1q)) \vee (\neg(p V0p)))))))))) \quad (21)$$

Assume the following.

$$\begin{aligned}
& (\forall V0p \in 2. (\forall V1q \in 2. (\forall V2r \in 2. (((p \ V0p) \Leftrightarrow (\\
& (p \ V1q) \wedge (p \ V2r))) \Leftrightarrow (((p \ V0p) \vee (\neg(p \ V1q)) \vee (\neg(p \ V2r)))) \wedge (((p \ V1q) \vee \\
& (\neg(p \ V0p))) \wedge ((p \ V2r) \vee (\neg(p \ V0p))))))))))
\end{aligned} \tag{22}$$

Assume the following.

$$\begin{aligned}
& (\forall V0p \in 2. (\forall V1q \in 2. (\forall V2r \in 2. (((p \ V0p) \Leftrightarrow (\\
& (p \ V1q) \vee (p \ V2r))) \Leftrightarrow (((p \ V0p) \vee (\neg(p \ V1q))) \wedge ((p \ V0p) \vee (\neg(p \ V2r)))) \wedge \\
& ((p \ V1q) \vee ((p \ V2r) \vee (\neg(p \ V0p))))))))))
\end{aligned} \tag{23}$$

Assume the following.

$$\begin{aligned}
& (\forall V0p \in 2. (\forall V1q \in 2. (\forall V2r \in 2. (((p \ V0p) \Leftrightarrow (\\
& (p \ V1q) \Rightarrow (p \ V2r))) \Leftrightarrow (((p \ V0p) \vee (p \ V1q)) \wedge (((p \ V0p) \vee (\neg(p \ V2r))) \wedge (\\
& \neg(p \ V1q)) \vee ((p \ V2r) \vee (\neg(p \ V0p))))))))))
\end{aligned} \tag{24}$$

Assume the following.

$$\begin{aligned}
& (\forall V0p \in 2. (\forall V1q \in 2. (((p \ V0p) \Leftrightarrow (\neg(p \ V1q))) \Leftrightarrow (((p \ V0p) \vee \\
& (p \ V1q)) \wedge ((\neg(p \ V1q)) \vee (\neg(p \ V0p))))))
\end{aligned} \tag{25}$$

Theorem 1

$$\begin{aligned}
& (\forall V0s \in (2^{ty_2Erealax_2Ereal}). ((p \ (ap \ (ap \ (c_2Ecardinal_2Ecardeq \\
& ty_2Erealax_2Ereal \ ty_2Erealax_2Ereal) \ V0s) \ (c_2Epred_set_2EUNIV \\
& ty_2Erealax_2Ereal))) \Rightarrow (\neg(p \ (ap \ (c_2Epred_set_2Ecountable \\
& ty_2Erealax_2Ereal) \ V0s))))))
\end{aligned}$$