

thm_2Ereal__topology_2ECBALL__SING (TM- MzMj5CeDoSYgDyNaJyHFPjtisfQPKFQA5)

October 26, 2020

Definition 1 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define $c_2Ebool_2E_7E$ to be $(ap (ap (c_2Emin_2E_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

Definition 3 We define $c_2Ebool_2E_21$ to be $\lambda A.\lambda a : \iota.(\lambda V0P \in (2^{A-27a}).(ap (ap (c_2Emin_2E_3D (2^{A-27a}))$

Definition 4 We define $c_2Ebool_2E_7E$ to be $(ap (c_2Ebool_2E_21 2) (\lambda V0t \in 2.V0t))$.

Definition 5 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p P \Rightarrow p Q)$ of type ι .

Definition 6 We define $c_2Ebool_2E_7E$ to be $(\lambda V0t \in 2.(ap (ap c_2Emin_2E_3D_3D_3E V0t) c_2Ebool_2E_7E))$

Definition 7 We define $c_2Ebool_2E_2F_5C$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21 2) (\lambda V2t \in 2.V2t))$

Let $c_2Enum_2EZERO_REP : \iota$ be given. Assume the following.

$$c_2Enum_2EZERO_REP \in \omega \tag{1}$$

Let $ty_2Enum_2Enum : \iota$ be given. Assume the following.

$$nonempty\ ty_2Enum_2Enum \tag{2}$$

Let $c_2Enum_2EABS_num : \iota$ be given. Assume the following.

$$c_2Enum_2EABS_num \in (ty_2Enum_2Enum^{\omega}) \tag{3}$$

Definition 8 We define c_2Enum_2E0 to be $(ap c_2Enum_2EABS_num c_2Enum_2EZERO_REP)$.

Let $ty_2Erealx_2Ereal : \iota$ be given. Assume the following.

$$nonempty\ ty_2Erealx_2Ereal \tag{4}$$

Let $c_2Ereal_2Ereal_of_num : \iota$ be given. Assume the following.

$$c_2Ereal_2Ereal_of_num \in (ty_2Erealx_2Ereal^{ty_2Enum_2Enum}) \tag{5}$$

Definition 9 We define $c_2Epred_set_2EEMPTY$ to be $\lambda A_27a : \iota. (\lambda V0x \in A_27a. c_2Ebool_2EF)$.

Definition 10 We define c_2Ebool_2EIN to be $\lambda A_27a : \iota. (\lambda V0x \in A_27a. (\lambda V1f \in (2^{A_27a}). (ap\ V1f\ V0x)))$

Definition 11 We define $c_2Ebool_2E_5C_2F$ to be $(\lambda V0t1 \in 2. (\lambda V1t2 \in 2. (ap\ (c_2Ebool_2E_21\ 2)\ (\lambda V2t \in 2. (c_2Ebool_2E_5C_2F\ V1t2\ V2t))))$

Let $ty_2Epair_2Eprod : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A0.nonempty\ A0 \Rightarrow \forall A1.nonempty\ A1 \Rightarrow nonempty\ (ty_2Epair_2Eprod\ A0\ A1) \quad (6)$$

Let $c_2Epair_2EABS_prod : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow \forall A_27b.nonempty\ A_27b \Rightarrow c_2Epair_2EABS_prod\ A_27a\ A_27b \in ((ty_2Epair_2Eprod\ A_27a\ A_27b)^{(2^{A_27b})^{A_27a}}) \quad (7)$$

Definition 12 We define $c_2Epair_2E_2C$ to be $\lambda A_27a : \iota. \lambda A_27b : \iota. \lambda V0x \in A_27a. \lambda V1y \in A_27b. (ap\ (c_2Ebool_2E_5C_2F\ V0x\ V1y))$

Let $c_2Epred_set_2EGSPEC : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow \forall A_27b.nonempty\ A_27b \Rightarrow c_2Epred_set_2EGSPEC\ A_27a\ A_27b \in ((2^{A_27a})^{(ty_2Epair_2Eprod\ A_27a\ 2)^{A_27b}}) \quad (8)$$

Definition 13 We define $c_2Epred_set_2EINSERT$ to be $\lambda A_27a : \iota. \lambda V0x \in A_27a. \lambda V1s \in (2^{A_27a}). (ap\ (c_2Ebool_2E_5C_2F\ V0x\ V1s))$

Let $c_2Ereal_topology_2Ecball : \iota$ be given. Assume the following.

$$c_2Ereal_topology_2Ecball \in ((2^{ty_2Erealax_2Ereal})^{(ty_2Epair_2Eprod\ ty_2Erealax_2Ereal\ ty_2Erealax_2Ereal)}) \quad (9)$$

Assume the following.

$$True \quad (10)$$

Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow (\forall V0t \in 2. ((\forall V1x \in A_27a. (p\ V0t)) \Leftrightarrow (p\ V0t))) \quad (11)$$

Assume the following.

$$\begin{aligned} & (\forall V0t \in 2. (((True \Rightarrow (p\ V0t)) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \Rightarrow True) \Leftrightarrow \\ & True) \wedge (((False \Rightarrow (p\ V0t)) \Leftrightarrow True) \wedge (((p\ V0t) \Rightarrow (p\ V0t)) \Leftrightarrow True) \wedge ((\\ & (p\ V0t) \Rightarrow False) \Leftrightarrow \neg (p\ V0t)))))) \end{aligned} \quad (12)$$

Assume the following.

$$\begin{aligned} & (\forall V0x \in ty_2Erealax_2Ereal. (\forall V1e \in ty_2Erealax_2Ereal. \\ & (((ap\ c_2Ereal_topology_2Ecball\ (ap\ (ap\ (c_2Epair_2E_2C\ ty_2Erealax_2Ereal\ ty_2Erealax_2Ereal)\ V0x)\ V1e)) = (ap\ (ap\ (c_2Epred_set_2EINSERT\ ty_2Erealax_2Ereal)\ V0x)\ (c_2Epred_set_2EEMPTY\ ty_2Erealax_2Ereal))) \Leftrightarrow \\ & (V1e = (ap\ c_2Ereal_2Ereal_of_num\ c_2Enum_2E0)))) \end{aligned} \quad (13)$$

Theorem 1

$$\begin{aligned} & (\forall V0x \in ty_2Erealax_2Ereal. (\forall V1e \in ty_2Erealax_2Ereal. \\ & ((V1e = (ap\ c_2Ereal_2Ereal_of_num\ c_2Enum_2E0)) \Rightarrow ((ap\ c_2Ereal_topology_2Ecball \\ & (ap\ (ap\ (c_2Epair_2E_2C\ ty_2Erealax_2Ereal\ ty_2Erealax_2Ereal) \\ & V0x)\ V1e)) = (ap\ (ap\ (c_2Epred_set_2EINSERT\ ty_2Erealax_2Ereal) \\ & V0x)\ (c_2Epred_set_2EEMPTY\ ty_2Erealax_2Ereal)))))) \end{aligned}$$