

thm\_2Ereal\_\_topology\_2ECLOSED\_\_SPHERE  
(TMLRaZ5BjtZtQPRF6Um75nUswvzxJ19heSc)

October 26, 2020

**Definition 1** We define  $c\_2Emin\_2E\_3D$  to be  $\lambda A.\lambda x \in A.\lambda y \in A.inj\_o (x = y)$  of type  $\iota \Rightarrow \iota$ .

**Definition 2** We define  $c\_2Ebool\_2E\_2T$  to be  $(ap (ap (c\_2Emin\_2E\_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

**Definition 3** We define  $c\_2Ebool\_2E\_21$  to be  $\lambda A\_27a : \iota.(\lambda V0P \in (2^{A\_27a}).(ap (ap (c\_2Emin\_2E\_3D (2^{A\_27a})))$

**Definition 4** We define  $c\_2Ebool\_2E\_2F$  to be  $(ap (c\_2Ebool\_2E\_21 2) (\lambda V0t \in 2.V0t))$ .

**Definition 5** We define  $c\_2Emin\_2E\_3D\_3D\_3E$  to be  $\lambda P \in 2.\lambda Q \in 2.inj\_o (p P \Rightarrow p Q)$  of type  $\iota$ .

**Definition 6** We define  $c\_2Ebool\_2E\_27E$  to be  $(\lambda V0t \in 2.(ap (ap c\_2Emin\_2E\_3D\_3D\_3E V0t) c\_2Ebool\_2E\_2F))$

**Definition 7** We define  $c\_2Ebool\_2E\_2F\_5C$  to be  $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c\_2Ebool\_2E\_21 2) (\lambda V2t \in 2.V2t)))$

Let  $ty\_2Erealax\_2Ereal : \iota$  be given. Assume the following.

$$nonempty\ ty\_2Erealax\_2Ereal \tag{1}$$

**Definition 8** We define  $c\_2Epred\_set\_2EUNIV$  to be  $\lambda A\_27a : \iota.(\lambda V0x \in A\_27a.c\_2Ebool\_2E\_2T)$ .

**Definition 9** We define  $c\_2Ebool\_2E\_2IN$  to be  $\lambda A\_27a : \iota.(\lambda V0x \in A\_27a.(\lambda V1f \in (2^{A\_27a}).(ap V1f V0x)))$

Let  $ty\_2Epair\_2Eprod : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A0.nonempty\ A0 \Rightarrow \forall A1.nonempty\ A1 \Rightarrow nonempty\ (ty\_2Epair\_2Eprod\ A0\ A1) \tag{2}$$

Let  $c\_2Epair\_2EABS\_prod : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A\_27a.nonempty\ A\_27a \Rightarrow \forall A\_27b.nonempty\ A\_27b \Rightarrow c\_2Epair\_2EABS\_prod\ A\_27a\ A\_27b \in ((ty\_2Epair\_2Eprod\ A\_27a\ A\_27b)^{(2^{A\_27b})^{A\_27a}}) \tag{3}$$

**Definition 10** We define  $c\_2Epair\_2E\_2C$  to be  $\lambda A\_27a : \iota.\lambda A\_27b : \iota.\lambda V0x \in A\_27a.\lambda V1y \in A\_27b.(ap (c\_2Ebool\_2E\_2IN$

Let  $c\_2Epred\_set\_2EGSPEC : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A\_27a.nonempty\ A\_27a \Rightarrow \forall A\_27b.nonempty\ A\_27b \Rightarrow c\_2Epred\_set\_2EGSPEC\ A\_27a\ A\_27b \in ((2^{A\_27a})^{(ty\_2Epair\_2Eprod\ A\_27a\ 2)^{A\_27b}})$$
(4)

**Definition 11** We define  $c\_2Epred\_set\_2EDIFF$  to be  $\lambda A\_27a : \iota.\lambda V0s \in (2^{A\_27a}).\lambda V1t \in (2^{A\_27a}).(ap\ (c\_2Epred\_set\_2EGSPEC\ A\_27a\ A\_27b)\ V0s\ V1t)$

Let  $c\_2Ereal\_topology\_2EDist : \iota$  be given. Assume the following.

$$c\_2Ereal\_topology\_2EDist \in (ty\_2Erealax\_2Ereal^{(ty\_2Epair\_2Eprod\ ty\_2Erealax\_2Ereal\ ty\_2Erealax\_2Ereal)})$$
(5)

Let  $ty\_2Ehreal\_2Ehreal : \iota$  be given. Assume the following.

$$nonempty\ ty\_2Ehreal\_2Ehreal$$
(6)

Let  $c\_2Erealax\_2Ereal\_REP\_CLASS : \iota$  be given. Assume the following.

$$c\_2Erealax\_2Ereal\_REP\_CLASS \in ((2^{(ty\_2Epair\_2Eprod\ ty\_2Ehreal\_2Ehreal\ ty\_2Ehreal\_2Ehreal)})^{ty\_2Erealax\_2Ereal})$$
(7)

**Definition 12** We define  $c\_2Emin\_2E\_40$  to be  $\lambda A.\lambda P \in 2^A$ .if  $(\exists x \in A.p\ (ap\ P\ x))$  then  $(the\ (\lambda x.x \in A \wedge P\ x))$  of type  $\iota \Rightarrow \iota$ .

**Definition 13** We define  $c\_2Erealax\_2Ereal\_REP$  to be  $\lambda V0a \in ty\_2Erealax\_2Ereal.(ap\ (c\_2Emin\_2E\_40\ (ap\ P\ x))\ V0a)$

Let  $c\_2Erealax\_2Etreall\_lt : \iota$  be given. Assume the following.

$$c\_2Erealax\_2Etreall\_lt \in ((2^{(ty\_2Epair\_2Eprod\ ty\_2Ehreal\_2Ehreal\ ty\_2Ehreal\_2Ehreal)})^{(ty\_2Epair\_2Eprod\ ty\_2Ehreal\_2Ehreal)})$$
(8)

**Definition 14** We define  $c\_2Erealax\_2Ereal\_lt$  to be  $\lambda V0T1 \in ty\_2Erealax\_2Ereal.\lambda V1T2 \in ty\_2Erealax\_2Ereal.(ap\ (c\_2Emin\_2E\_40\ (ap\ P\ x))\ V0T1\ V1T2)$

Let  $c\_2Enum\_2EZERO\_REP : \iota$  be given. Assume the following.

$$c\_2Enum\_2EZERO\_REP \in \omega$$
(9)

Let  $ty\_2Enum\_2Enum : \iota$  be given. Assume the following.

$$nonempty\ ty\_2Enum\_2Enum$$
(10)

Let  $c\_2Enum\_2EABS\_num : \iota$  be given. Assume the following.

$$c\_2Enum\_2EABS\_num \in (ty\_2Enum\_2Enum^{\omega})$$
(11)

**Definition 15** We define  $c\_2Enum\_2E0$  to be  $(ap\ c\_2Enum\_2EABS\_num\ c\_2Enum\_2EZERO\_REP)$ .

Let  $c\_2Ereal\_2Ereal\_of\_num : \iota$  be given. Assume the following.

$$c\_2Ereal\_2Ereal\_of\_num \in (ty\_2Erealax\_2Ereal^{ty\_2Enum\_2Enum})$$
(12)

**Definition 16** We define  $c\_2Ebool\_2E\_3F$  to be  $\lambda A\_27a : \iota.(\lambda V0P \in (2^{A\_27a}).(ap\ V0P\ (ap\ (c\_2Emin\_2E\_40\ (ap\ P\ x))\ V0a)\ V0P))$

**Definition 17** We define  $c\_Ereal\_topology\_2EOpen$  to be  $\lambda V0s \in (2^{ty\_2Erealax\_2Ereal}).(ap (c\_Ebool\_2E2$

**Definition 18** We define  $c\_Ereal\_topology\_2EClosed$  to be  $\lambda V0s \in (2^{ty\_2Erealax\_2Ereal}).(ap c\_Ereal\_topo$

Let  $c\_Ereal\_topology\_2Esphere : \iota$  be given. Assume the following.

$$c\_Ereal\_topology\_2Esphere \in ((2^{ty\_2Erealax\_2Ereal})^{(ty\_2Epair\_2Eprod ty\_2Erealax\_2Ereal ty\_2Erealax\_2Ereal)}) \quad (13)$$

Let  $c\_Eenum\_2EREP\_num : \iota$  be given. Assume the following.

$$c\_Eenum\_2EREP\_num \in (\omega^{ty\_2Eenum\_2Eenum}) \quad (14)$$

Let  $c\_Eenum\_2ESUC\_REP : \iota$  be given. Assume the following.

$$c\_Eenum\_2ESUC\_REP \in (\omega^{\omega}) \quad (15)$$

**Definition 19** We define  $c\_Eenum\_2ESUC$  to be  $\lambda V0m \in ty\_2Eenum\_2Eenum.(ap c\_Eenum\_2EABS\_num$

**Definition 20** We define  $c\_Eprim\_rec\_2E\_3C$  to be  $\lambda V0m \in ty\_2Eenum\_2Eenum.\lambda V1n \in ty\_2Eenum\_2Eenum$

**Definition 21** We define  $c\_Earithmic\_2E\_3E$  to be  $\lambda V0m \in ty\_2Eenum\_2Eenum.\lambda V1n \in ty\_2Eenum\_2Eenum$

**Definition 22** We define  $c\_Ebool\_2E\_5C\_2F$  to be  $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c\_Ebool\_2E\_21 2) (\lambda V2t \in$

**Definition 23** We define  $c\_Earithmic\_2E\_3E\_3D$  to be  $\lambda V0m \in ty\_2Eenum\_2Eenum.\lambda V1n \in ty\_2Eenum\_2Eenum$

Let  $ty\_2Ereal\_topology\_2Enet : \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A0.nonempty A0 \Rightarrow nonempty (ty\_2Ereal\_topology\_2Enet A0) \quad (16)$$

Let  $c\_Ereal\_topology\_2Emk\_net : \iota \Rightarrow \iota$  be given. Assume the following.

$$\begin{aligned} \forall A.27a.nonempty A.27a \Rightarrow c\_Ereal\_topology\_2Emk\_net \\ A.27a \in ((ty\_2Ereal\_topology\_2Enet A.27a)^{(2^{A.27a})^{A.27a}}) \end{aligned} \quad (17)$$

**Definition 24** We define  $c\_Ereal\_topology\_2Esequentially$  to be  $(ap (c\_Ereal\_topology\_2Emk\_net ty\_2E$

**Definition 25** We define  $c\_Ecombin\_2Eo$  to be  $\lambda A.27a : \iota.\lambda A.27b : \iota.\lambda A.27c : \iota.\lambda V0f \in (A.27b^{A.27c}).\lambda V1g \in$

Let  $c\_Ereal\_topology\_2Enetord : \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A.27a.nonempty A.27a \Rightarrow c\_Ereal\_topology\_2Enetord A.27a \in (((2^{A.27a})^{A.27a})^{(ty\_2Ereal\_topology\_2Enet A.27a)}) \quad (18)$$

**Definition 26** We define  $c\_Ereal\_topology\_2Etrivial\_limit$  to be  $\lambda A.27a : \iota.\lambda V0net \in (ty\_2Ereal\_topology$

**Definition 27** We define  $c\_Ereal\_topology\_2Eeventually$  to be  $\lambda A.27a : \iota.\lambda V0p \in (2^{A.27a}).\lambda V1net \in (ty\_2E$

**Definition 28** We define  $c\_Ereal\_topology\_2E\_2D\_2D\_3E$  to be  $\lambda A.27a : \iota.\lambda V0f \in (ty\_2Erealax\_2Ereal^A$

**Definition 29** We define  $c\_Ereal\_topology\_2Ecompact$  to be  $\lambda V0s \in (2^{ty\_2Erealax\_2Ereal}).(ap (c\_Ebool\_2E2$

Assume the following.

$$True \quad (19)$$

Assume the following.

$$\forall A_{27a}. nonempty\ A_{27a} \Rightarrow (\forall V0t \in 2. ((\forall V1x \in A_{27a}. (p\ V0t)) \Leftrightarrow (p\ V0t))) \quad (20)$$

Assume the following.

$$\begin{aligned} & (\forall V0t \in 2. (((True \Leftrightarrow (p\ V0t)) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \Leftrightarrow True) \Leftrightarrow \\ & (p\ V0t)) \wedge (((False \Leftrightarrow (p\ V0t)) \Leftrightarrow \neg(p\ V0t)) \wedge (((p\ V0t) \Leftrightarrow False) \Leftrightarrow \neg( \\ & p\ V0t)))))) \quad (21) \end{aligned}$$

Assume the following.

$$(\forall V0s \in (2^{ty\_2Erealax\_2Ereal}). ((p\ (ap\ c\_2Ereal\_topology\_2Ecompact\ V0s)) \Rightarrow (p\ (ap\ c\_2Ereal\_topology\_2EClosed\ V0s)))) \quad (22)$$

Assume the following.

$$\begin{aligned} & (\forall V0a \in ty\_2Erealax\_2Ereal. (\forall V1r \in ty\_2Erealax\_2Ereal. \\ & (p\ (ap\ c\_2Ereal\_topology\_2Ecompact\ (ap\ c\_2Ereal\_topology\_2Esphere \\ & (ap\ (ap\ (c\_2Epair\_2E\_2C\ ty\_2Erealax\_2Ereal\ ty\_2Erealax\_2Ereal) \\ & V0a\ V1r)))))) \quad (23) \end{aligned}$$

**Theorem 1**

$$\begin{aligned} & (\forall V0a \in ty\_2Erealax\_2Ereal. (\forall V1r \in ty\_2Erealax\_2Ereal. \\ & (p\ (ap\ c\_2Ereal\_topology\_2EClosed\ (ap\ c\_2Ereal\_topology\_2Esphere \\ & (ap\ (ap\ (c\_2Epair\_2E\_2C\ ty\_2Erealax\_2Ereal\ ty\_2Erealax\_2Ereal) \\ & V0a\ V1r)))))) \end{aligned}$$