

thm\_2Ereal\_\_topology\_2ECLOSED\_\_SUBSTANDARD  
(TMN-  
Wek3B52W5aWVRdCAM37qpgS7h3dUwesp)

October 26, 2020

Let  $c\_2Enum\_2EZERO\_REP : \iota$  be given. Assume the following.

$$c\_2Enum\_2EZERO\_REP \in \omega \tag{1}$$

Let  $ty\_2Enum\_2Enum : \iota$  be given. Assume the following.

$$nonempty\ ty\_2Enum\_2Enum \tag{2}$$

Let  $c\_2Enum\_2EABS\_num : \iota$  be given. Assume the following.

$$c\_2Enum\_2EABS\_num \in (ty\_2Enum\_2Enum^{\omega}) \tag{3}$$

**Definition 1** We define  $c\_2Emin\_2E3D$  to be  $\lambda A.\lambda x \in A.\lambda y \in A.inj\_o (x = y)$  of type  $\iota \Rightarrow \iota$ .

**Definition 2** We define  $c\_2Enum\_2E0$  to be  $(ap\ c\_2Enum\_2EABS\_num\ c\_2Enum\_2EZERO\_REP)$ .

Let  $ty\_2Erealx\_2Ereal : \iota$  be given. Assume the following.

$$nonempty\ ty\_2Erealx\_2Ereal \tag{4}$$

Let  $c\_2Ereal\_2Ereal\_of\_num : \iota$  be given. Assume the following.

$$c\_2Ereal\_2Ereal\_of\_num \in (ty\_2Erealx\_2Ereal^{ty\_2Enum\_2Enum}) \tag{5}$$

**Definition 3** We define  $c\_2Ebool\_2E2$  to be  $(ap\ (ap\ (c\_2Emin\_2E3D\ (2^2))\ (\lambda V0x \in 2.V0x))\ (\lambda V1x \in 2.V1x))$

**Definition 4** We define  $c\_2Ebool\_2E21$  to be  $\lambda A.27a : \iota.(\lambda V0P \in (2^{A-27a}).(ap\ (ap\ (c\_2Emin\_2E3D\ (2^{A-27a}))\ (\lambda V1Q \in 2.V1Q))\ (\lambda V2Q \in 2.V2Q))$

**Definition 5** We define  $c\_2Ebool\_2E2F$  to be  $(ap\ (c\_2Ebool\_2E21\ 2)\ (\lambda V0t \in 2.V0t))$ .

**Definition 6** We define  $c\_2Emin\_2E3D\_3D\_3E$  to be  $\lambda P \in 2.\lambda Q \in 2.inj\_o (p\ P \Rightarrow p\ Q)$  of type  $\iota$ .

**Definition 7** We define  $c\_2Ebool\_2E\_7E$  to be  $(\lambda V0t \in 2.(ap (ap c\_2Emin\_2E\_3D\_3D\_3E V0t) c\_2Ebool\_2EF)$

**Definition 8** We define  $c\_2Ebool\_2E\_2F\_5C$  to be  $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c\_2Ebool\_2E\_21 2) (\lambda V2t \in 2.$

Let  $ty\_2Epair\_2Eprod : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A0.nonempty A0 \Rightarrow \forall A1.nonempty A1 \Rightarrow nonempty (ty\_2Epair\_2Eprod A0 A1) \quad (6)$$

Let  $c\_2Epair\_2EABS\_prod : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A\_27a.nonempty A\_27a \Rightarrow \forall A\_27b.nonempty A\_27b \Rightarrow c\_2Epair\_2EABS\_prod A\_27a A\_27b \in ((ty\_2Epair\_2Eprod A\_27a A\_27b)^{(2^{A\_27b})^{A\_27a}}) \quad (7)$$

**Definition 9** We define  $c\_2Epair\_2E\_2C$  to be  $\lambda A\_27a : \iota.\lambda A\_27b : \iota.\lambda V0x \in A\_27a.\lambda V1y \in A\_27b.(ap (c\_2E$

Let  $c\_2Epred\_set\_2EGSPEC : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A\_27a.nonempty A\_27a \Rightarrow \forall A\_27b.nonempty A\_27b \Rightarrow c\_2Epred\_set\_2EGSPEC A\_27a A\_27b \in ((2^{A\_27a})^{(ty\_2Epair\_2Eprod A\_27a 2)^{A\_27b}}) \quad (8)$$

**Definition 10** We define  $c\_2Epred\_set\_2EEMPTY$  to be  $\lambda A\_27a : \iota.(\lambda V0x \in A\_27a.c\_2Ebool\_2EF)$ .

**Definition 11** We define  $c\_2Ebool\_2EIN$  to be  $\lambda A\_27a : \iota.(\lambda V0x \in A\_27a.(\lambda V1f \in (2^{A\_27a}).(ap V1f V0x))$

**Definition 12** We define  $c\_2Ebool\_2E\_5C\_2F$  to be  $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c\_2Ebool\_2E\_21 2) (\lambda V2t \in 2.$

**Definition 13** We define  $c\_2Epred\_set\_2EINSERT$  to be  $\lambda A\_27a : \iota.\lambda V0x \in A\_27a.\lambda V1s \in (2^{A\_27a}).(ap (c\_2E$

**Definition 14** We define  $c\_2Epred\_set\_2EUNIV$  to be  $\lambda A\_27a : \iota.(\lambda V0x \in A\_27a.c\_2Ebool\_2ET)$ .

**Definition 15** We define  $c\_2Epred\_set\_2EDIFF$  to be  $\lambda A\_27a : \iota.\lambda V0s \in (2^{A\_27a}).\lambda V1t \in (2^{A\_27a}).(ap (c\_2E$

Let  $c\_2Ereal\_topology\_2EDist : \iota$  be given. Assume the following.

$$c\_2Ereal\_topology\_2EDist \in (ty\_2Erealax\_2Ereal^{(ty\_2Epair\_2Eprod ty\_2Erealax\_2Ereal ty\_2Erealax\_2Ereal)}) \quad (9)$$

Let  $ty\_2Ehreal\_2Ehreal : \iota$  be given. Assume the following.

$$nonempty ty\_2Ehreal\_2Ehreal \quad (10)$$

Let  $c\_2Erealax\_2Ereal\_REP\_CLASS : \iota$  be given. Assume the following.

$$c\_2Erealax\_2Ereal\_REP\_CLASS \in ((2^{(ty\_2Epair\_2Eprod ty\_2Ehreal\_2Ehreal ty\_2Ehreal\_2Ehreal)})^{ty\_2Erealax\_2Ereal}) \quad (11)$$

**Definition 16** We define  $c\_2Emin\_2E\_40$  to be  $\lambda A.\lambda P \in 2^A.\mathbf{if} (\exists x \in A.p (ap P x)) \mathbf{then} (the (\lambda x.x \in A \wedge p x))$  of type  $\iota \Rightarrow \iota$ .

**Definition 17** We define  $c\_2Erealax\_2Ereal\_REP$  to be  $\lambda V0a \in ty\_2Erealax\_2Ereal.(ap (c\_2Emin\_2E\_40 (t$

Let  $c\_2Erealax\_2Etreall\_lt : \iota$  be given. Assume the following.

$$c\_2Erealax\_2Etreall\_lt \in ((2^{(ty\_2Epair\_2Eprod\ ty\_2Ehreal\_2Ehreal\ ty\_2Ehreal\_2Ehreal)})(ty\_2Epair\_2Eprod\ ty\_2Ehreal\_2Ehreal)) \quad (12)$$

**Definition 18** We define  $c\_2Erealax\_2Ereal\_lt$  to be  $\lambda V0T1 \in ty\_2Erealax\_2Ereal.\lambda V1T2 \in ty\_2Erealax\_2Ereal$

**Definition 19** We define  $c\_2Ebool\_2E\_3F$  to be  $\lambda A\_27a : \iota.(\lambda V0P \in (2^{A\_27a}).(ap\ V0P\ (ap\ (c\_2Emin\_2E\_40$

**Definition 20** We define  $c\_2Ereal\_topology\_2EOpen$  to be  $\lambda V0s \in (2^{ty\_2Erealax\_2Ereal}).(ap\ (c\_2Ebool\_2E\_3F$

**Definition 21** We define  $c\_2Ereal\_topology\_2EClosed$  to be  $\lambda V0s \in (2^{ty\_2Erealax\_2Ereal}).(ap\ c\_2Ereal\_topo$

Assume the following.

$$True \quad (13)$$

Assume the following.

$$\begin{aligned} & (\forall V0t \in 2.(((True \Leftrightarrow (p\ V0t)) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \Leftrightarrow True) \Leftrightarrow \\ & (p\ V0t)) \wedge (((False \Leftrightarrow (p\ V0t)) \Leftrightarrow \neg(p\ V0t)) \wedge (((p\ V0t) \Leftrightarrow False) \Leftrightarrow \neg( \\ & p\ V0t)))))) \end{aligned} \quad (14)$$

Assume the following.

$$\begin{aligned} & \forall A\_27a.nonempty\ A\_27a \Rightarrow (\forall V0y \in A\_27a.((ap\ (c\_2Epred\_set\_2EGSPEC \\ & A\_27a\ A\_27a)\ (\lambda V1x \in A\_27a.(ap\ (ap\ (c\_2Epair\_2E\_2C\ A\_27a\ 2) \\ & V1x)\ (ap\ (ap\ (c\_2Emin\_2E\_3D\ A\_27a)\ V1x)\ V0y)))) = (ap\ (ap\ (c\_2Epred\_set\_2EINSERT \\ & A\_27a)\ V0y)\ (c\_2Epred\_set\_2EEMPTY\ A\_27a)))) \end{aligned} \quad (15)$$

Assume the following.

$$\begin{aligned} & (\forall V0a \in ty\_2Erealax\_2Ereal.(p\ (ap\ c\_2Ereal\_topology\_2EClosed \\ & (ap\ (ap\ (c\_2Epred\_set\_2EINSERT\ ty\_2Erealax\_2Ereal)\ V0a)\ (c\_2Epred\_set\_2EEMPTY \\ & ty\_2Erealax\_2Ereal)))))) \end{aligned} \quad (16)$$

**Theorem 1**

$$\begin{aligned} & (p\ (ap\ c\_2Ereal\_topology\_2EClosed\ (ap\ (c\_2Epred\_set\_2EGSPEC \\ & ty\_2Erealax\_2Ereal\ ty\_2Erealax\_2Ereal)\ (\lambda V0x \in ty\_2Erealax\_2Ereal. \\ & (ap\ (ap\ (c\_2Epair\_2E\_2C\ ty\_2Erealax\_2Ereal\ 2)\ V0x)\ (ap\ (ap\ (c\_2Emin\_2E\_3D \\ & ty\_2Erealax\_2Ereal)\ V0x)\ (ap\ c\_2Ereal\_2Ereal\_of\_num\ c\_2Enum\_2E0)))))) \end{aligned}$$