

thm_2Ereal__topology_2ECLOSED__UNIV
 (TMaRBmeMuDaduuvJYY-
 BQm9JNLGwY7eCwtpB)

October 26, 2020

Definition 1 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define c_2Ebool_2ET to be $(ap (ap (c_2Emin_2E_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

Definition 3 We define $c_2Ebool_2E_21$ to be $\lambda A_27a : \iota.(\lambda V0P \in (2^{A_27a}).(ap (ap (c_2Emin_2E_3D (2^{A_27a}))$

Definition 4 We define c_2Ebool_2EF to be $(ap (c_2Ebool_2E_21 2) (\lambda V0t \in 2.V0t))$.

Definition 5 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p P \Rightarrow q Q)$ of type ι .

Definition 6 We define $c_2Ebool_2E_7E$ to be $(\lambda V0t \in 2.(ap (ap c_2Emin_2E_3D_3D_3E V0t) c_2Ebool_2EF$

Definition 7 We define $c_2Ebool_2E_2F_5C$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21 2) (\lambda V2t \in 2.V2t))$

Definition 8 We define $c_2Epred_set_2EUNIV$ to be $\lambda A_27a : \iota.(\lambda V0x \in A_27a.c_2Ebool_2ET)$.

Let $ty_2Erealx_2Ereal : \iota$ be given. Assume the following.

$$nonempty\ ty_2Erealx_2Ereal \tag{1}$$

Definition 9 We define c_2Ebool_2EIN to be $\lambda A_27a : \iota.(\lambda V0x \in A_27a.(\lambda V1f \in (2^{A_27a}).(ap V1f V0x)))$

Let $ty_2Epair_2Eprod : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A0.nonempty\ A0 \Rightarrow \forall A1.nonempty\ A1 \Rightarrow nonempty\ (ty_2Epair_2Eprod\ A0\ A1) \tag{2}$$

Let $c_2Epair_2EABS_prod : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow \forall A_27b.nonempty\ A_27b \Rightarrow c_2Epair_2EABS_prod\ A_27a\ A_27b \in ((ty_2Epair_2Eprod\ A_27a\ A_27b)^{(2^{A_27b})^{A_27a}}) \tag{3}$$

Definition 10 We define $c_2Epair_2E_2C$ to be $\lambda A.27a : \iota.\lambda A.27b : \iota.\lambda V0x \in A.27a.\lambda V1y \in A.27b.(ap (c_2Ereal_topology_2EDist : \iota$ be given. Assume the following.

$$c_2Ereal_topology_2EDist \in (ty_2Erealax_2Ereal^{(ty_2Epair_2Eprod\ ty_2Erealax_2Ereal\ ty_2Erealax_2Ereal)}) \quad (4)$$

Let $ty_2Ehreal_2Ehreal : \iota$ be given. Assume the following.

$$nonempty\ ty_2Ehreal_2Ehreal \quad (5)$$

Let $c_2Erealax_2Ereal_REP_CLASS : \iota$ be given. Assume the following.

$$c_2Erealax_2Ereal_REP_CLASS \in ((2^{(ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal\ ty_2Ehreal_2Ehreal)})^{ty_2Erealax}) \quad (6)$$

Definition 11 We define $c_2Emin_2E_40$ to be $\lambda A.\lambda P \in 2^A.$ **if** $(\exists x \in A.p (ap\ P\ x))$ **then** $(the (\lambda x.x \in A \wedge p\ x))$ of type $\iota \Rightarrow \iota$.

Definition 12 We define $c_2Erealax_2Ereal_REP$ to be $\lambda V0a \in ty_2Erealax_2Ereal.(ap (c_2Emin_2E_40 (t$

Let $c_2Erealax_2Etrealm_lt : \iota$ be given. Assume the following.

$$c_2Erealax_2Etrealm_lt \in ((2^{(ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal\ ty_2Ehreal_2Ehreal)})^{(ty_2Epair_2Eprod\ ty_2Ehreal)}) \quad (7)$$

Definition 13 We define $c_2Erealax_2Ereal_lt$ to be $\lambda V0T1 \in ty_2Erealax_2Ereal.\lambda V1T2 \in ty_2Erealax_2Ereal.$

Let $c_2Enum_2EZERO_REP : \iota$ be given. Assume the following.

$$c_2Enum_2EZERO_REP \in omega \quad (8)$$

Let $ty_2Enum_2Enum : \iota$ be given. Assume the following.

$$nonempty\ ty_2Enum_2Enum \quad (9)$$

Let $c_2Enum_2EABS_num : \iota$ be given. Assume the following.

$$c_2Enum_2EABS_num \in (ty_2Enum_2Enum^{omega}) \quad (10)$$

Definition 14 We define c_2Enum_2E0 to be $(ap\ c_2Enum_2EABS_num\ c_2Enum_2EZERO_REP)$.

Let $c_2Ereal_2Ereal_of_num : \iota$ be given. Assume the following.

$$c_2Ereal_2Ereal_of_num \in (ty_2Erealax_2Ereal^{ty_2Enum_2Enum}) \quad (11)$$

Definition 15 We define $c_2Ebool_2E_3F$ to be $\lambda A.27a : \iota.(\lambda V0P \in (2^{A.27a}).(ap\ V0P (ap (c_2Emin_2E_40 (t$

Definition 16 We define $c_2Ereal_topology_2EOpen$ to be $\lambda V0s \in (2^{ty_2Erealax_2Ereal}).(ap (c_2Ebool_2E_2$

Let $ty_2Etopology_2Etopology : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A0.nonempty\ A0 \Rightarrow nonempty\ (ty_2Etopology_2Etopology\ A0) \quad (12)$$

Let $c_2Etopology_2Etopology : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow c_2Etopology_2Etopology\ A_27a \in ((ty_2Etopology_2Etopology\ A_27a)^{(2^{2^A-27a}})) \quad (13)$$

Definition 17 We define $c_2Ereal_topology_2Euclidean$ to be $(ap\ (c_2Etopology_2Etopology\ ty_2Erealax_2Ereal))$.

Let $c_2Epred_set_2EGSPEC : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow \forall A_27b.nonempty\ A_27b \Rightarrow c_2Epred_set_2EGSPEC\ A_27a\ A_27b \in ((2^{A_27a})^{(ty_2Epair_2Eprod\ A_27a\ 2)^{A_27b}}) \quad (14)$$

Definition 18 We define $c_2Epred_set_2EDIFF$ to be $\lambda A_27a : \iota.\lambda V0s \in (2^{A_27a}).\lambda V1t \in (2^{A_27a}).(ap\ (c_2Epred_set_2EGSPEC\ A_27a\ A_27a))$.

Definition 19 We define $c_2Ereal_topology_2EClosed$ to be $\lambda V0s \in (2^{ty_2Erealax_2Ereal}).(ap\ c_2Ereal_topology_2Euclidean)$.

Let $c_2Etopology_2Eopen_in : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow c_2Etopology_2Eopen_in\ A_27a \in ((2^{(2^{A_27a})})^{(ty_2Etopology_2Etopology\ A_27a)}) \quad (15)$$

Definition 20 We define $c_2Epred_set_2EBIGUNION$ to be $\lambda A_27a : \iota.\lambda V0P \in (2^{(2^{A_27a})}).(ap\ (c_2Epred_set_2EGSPEC\ A_27a\ A_27a))$.

Definition 21 We define $c_2Etopology_2Etopspace$ to be $\lambda A_27a : \iota.\lambda V0top \in (ty_2Etopology_2Etopology\ A_27a)$.

Definition 22 We define $c_2Epred_set_2ESUBSET$ to be $\lambda A_27a : \iota.\lambda V0s \in (2^{A_27a}).\lambda V1t \in (2^{A_27a}).(ap\ (c_2Epred_set_2EDIFF\ A_27a\ A_27a))$.

Definition 23 We define $c_2Etopology_2Eclosed_in$ to be $\lambda A_27a : \iota.\lambda V0top \in (ty_2Etopology_2Etopology\ A_27a)$.

Assume the following.

$$True \quad (16)$$

Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow (\forall V0x \in A_27a.(\forall V1y \in A_27a.((V0x = V1y) \Leftrightarrow (V1y = V0x)))) \quad (17)$$

Assume the following.

$$(\forall V0t \in 2.(((True \Leftrightarrow (p\ V0t)) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \Leftrightarrow True) \Leftrightarrow (p\ V0t)) \wedge (((False \Leftrightarrow (p\ V0t)) \Leftrightarrow \neg(p\ V0t)) \wedge (((p\ V0t) \Leftrightarrow False) \Leftrightarrow \neg(p\ V0t)))))) \quad (18)$$

Assume the following.

$$((ap\ (c_2Etopology_2Etopspace\ ty_2Erealax_2Ereal)\ c_2Ereal_topology_2Euclidean) = (c_2Epred_set_2EUNIV\ ty_2Erealax_2Ereal)) \quad (19)$$

Assume the following.

$$(\forall V0s \in (2^{ty_2Erealax_2Ereal}).((p (ap c_2Ereal_topology_2EClosed V0s)) \Leftrightarrow (p (ap (ap (c_2Etopology_2Eclosed_in ty_2Erealax_2Ereal) c_2Ereal_topology_2Euclidean) V0s)))) \quad (20)$$

Assume the following.

$$\forall A_27a.nonempty A_27a \Rightarrow (\forall V0top \in (ty_2Etopology_2Etopology A_27a).(p (ap (ap (c_2Etopology_2Eclosed_in A_27a) V0top) (ap (c_2Etopology_2Etopspace A_27a) V0top)))) \quad (21)$$

Theorem 1

$$(p (ap c_2Ereal_topology_2EClosed (c_2Epred_set_2EUNIV ty_2Erealax_2Ereal)))$$