

thm\_2Ereal\_\_topology\_2ECLOSURE\_\_MINIMAL  
 (TMXQJnWohuRXhAKgfwD-  
 HVAoqK7ngiMzHJuv)

October 26, 2020

**Definition 1** We define  $c\_2Emin\_2E\_3D$  to be  $\lambda A.\lambda x \in A.\lambda y \in A.inj\_o (x = y)$  of type  $\iota \Rightarrow \iota$ .

**Definition 2** We define  $c\_2Ebool\_2E\_2T$  to be  $(ap (ap (c\_2Emin\_2E\_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

**Definition 3** We define  $c\_2Ebool\_2E\_21$  to be  $\lambda A\_27a : \iota.(\lambda V0P \in (2^{A\_27a}).(ap (ap (c\_2Emin\_2E\_3D (2^{A\_27a})))$

**Definition 4** We define  $c\_2Ebool\_2E\_2F$  to be  $(ap (c\_2Ebool\_2E\_21 2) (\lambda V0t \in 2.V0t))$ .

**Definition 5** We define  $c\_2Emin\_2E\_3D\_3D\_3E$  to be  $\lambda P \in 2.\lambda Q \in 2.inj\_o (p P \Rightarrow p Q)$  of type  $\iota$ .

**Definition 6** We define  $c\_2Ebool\_2E\_7E$  to be  $(\lambda V0t \in 2.(ap (ap c\_2Emin\_2E\_3D\_3D\_3E V0t) c\_2Ebool\_2E\_2F$

Let  $ty\_2Erealx\_2Ereal : \iota$  be given. Assume the following.

$$nonempty\ ty\_2Erealx\_2Ereal \tag{1}$$

**Definition 7** We define  $c\_2Epred\_set\_2EUNIV$  to be  $\lambda A\_27a : \iota.(\lambda V0x \in A\_27a.c\_2Ebool\_2E\_2T)$ .

**Definition 8** We define  $c\_2Ebool\_2E\_2IN$  to be  $\lambda A\_27a : \iota.(\lambda V0x \in A\_27a.(\lambda V1f \in (2^{A\_27a}).(ap V1f V0x)))$

**Definition 9** We define  $c\_2Ebool\_2E\_2F\_5C$  to be  $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c\_2Ebool\_2E\_21 2) (\lambda V2t \in 2.V2t)))$

Let  $ty\_2Epair\_2Eprod : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A0.nonempty\ A0 \Rightarrow \forall A1.nonempty\ A1 \Rightarrow nonempty\ (ty\_2Epair\_2Eprod\ A0\ A1) \tag{2}$$

Let  $c\_2Epair\_2EABS\_prod : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A\_27a.nonempty\ A\_27a \Rightarrow \forall A\_27b.nonempty\ A\_27b \Rightarrow c\_2Epair\_2EABS\_prod\ A\_27a\ A\_27b \in ((ty\_2Epair\_2Eprod\ A\_27a\ A\_27b)^{(2^{A\_27b})^{A\_27a}}) \tag{3}$$

**Definition 10** We define  $c\_2Epair\_2E\_2C$  to be  $\lambda A\_27a : \iota. \lambda A\_27b : \iota. \lambda V0x \in A\_27a. \lambda V1y \in A\_27b. (ap (c\_2Epred\_set\_2EGSPEC : \iota \Rightarrow \iota \Rightarrow \iota)$  be given. Assume the following.

$$\forall A\_27a.nonempty\ A\_27a \Rightarrow \forall A\_27b.nonempty\ A\_27b \Rightarrow c\_2Epred\_set\_2EGSPEC\ A\_27a\ A\_27b \in ((2^{A\_27a})^{(ty\_2Epair\_2Eprod\ A\_27a\ 2)^{A\_27b}})$$
(4)

**Definition 11** We define  $c\_2Epred\_set\_2EDIFF$  to be  $\lambda A\_27a : \iota. \lambda V0s \in (2^{A\_27a}). \lambda V1t \in (2^{A\_27a}). (ap (c\_2Ereal\_topology\_2EDist : \iota$  be given. Assume the following.

$$c\_2Ereal\_topology\_2EDist \in (ty\_2Erealax\_2Ereal^{(ty\_2Epair\_2Eprod\ ty\_2Erealax\_2Ereal\ ty\_2Erealax\_2Ereal)})$$
(5)

Let  $ty\_2Ehreal\_2Ehreal : \iota$  be given. Assume the following.

$$nonempty\ ty\_2Ehreal\_2Ehreal$$
(6)

Let  $c\_2Erealax\_2Ereal\_REP\_CLASS : \iota$  be given. Assume the following.

$$c\_2Erealax\_2Ereal\_REP\_CLASS \in ((2^{(ty\_2Epair\_2Eprod\ ty\_2Ehreal\_2Ehreal\ ty\_2Ehreal\_2Ehreal)})^{ty\_2Erealax\_2Ereal})$$
(7)

**Definition 12** We define  $c\_2Emin\_2E\_40$  to be  $\lambda A. \lambda P \in 2^A. \text{if } (\exists x \in A. p (ap\ P\ x)) \text{ then } (the\ (\lambda x. x \in A \wedge p\ x))$  of type  $\iota \Rightarrow \iota$ .

**Definition 13** We define  $c\_2Erealax\_2Ereal\_REP$  to be  $\lambda V0a \in ty\_2Erealax\_2Ereal. (ap (c\_2Emin\_2E\_40 (t$

Let  $c\_2Erealax\_2Etreall\_lt : \iota$  be given. Assume the following.

$$c\_2Erealax\_2Etreall\_lt \in ((2^{(ty\_2Epair\_2Eprod\ ty\_2Ehreal\_2Ehreal\ ty\_2Ehreal\_2Ehreal)})^{(ty\_2Epair\_2Eprod\ ty\_2Ehreal\_2Ehreal)})$$
(8)

**Definition 14** We define  $c\_2Erealax\_2Ereal\_lt$  to be  $\lambda V0T1 \in ty\_2Erealax\_2Ereal. \lambda V1T2 \in ty\_2Erealax\_2Ereal.$

Let  $c\_2Enum\_2EZERO\_REP : \iota$  be given. Assume the following.

$$c\_2Enum\_2EZERO\_REP \in \omega$$
(9)

Let  $ty\_2Enum\_2Enum : \iota$  be given. Assume the following.

$$nonempty\ ty\_2Enum\_2Enum$$
(10)

Let  $c\_2Enum\_2EABS\_num : \iota$  be given. Assume the following.

$$c\_2Enum\_2EABS\_num \in (ty\_2Enum\_2Enum^{\omega})$$
(11)

**Definition 15** We define  $c\_2Enum\_2E0$  to be  $(ap\ c\_2Enum\_2EABS\_num\ c\_2Enum\_2EZERO\_REP)$ .

Let  $c\_2Ereal\_2Ereal\_of\_num : \iota$  be given. Assume the following.

$$c\_2Ereal\_2Ereal\_of\_num \in (ty\_2Erealax\_2Ereal^{ty\_2Enum\_2Enum})$$
(12)

**Definition 16** We define  $c\_2Ebool\_2E\_3F$  to be  $\lambda A\_27a : \iota. (\lambda V0P \in (2^{A-27a}). (ap\ V0P\ (ap\ (c\_2Emin\_2E\_40$

**Definition 17** We define  $c\_2Ereal\_topology\_2EOpen$  to be  $\lambda V0s \in (2^{ty\_2Erealax\_2Ereal}). (ap\ (c\_2Ebool\_2E\_2$

**Definition 18** We define  $c\_2Ereal\_topology\_2EClosed$  to be  $\lambda V0s \in (2^{ty\_2Erealax\_2Ereal}). (ap\ c\_2Ereal\_topo$

**Definition 19** We define  $c\_2Ereal\_topology\_2Elimit\_point\_of$  to be  $\lambda V0x \in ty\_2Erealax\_2Ereal. \lambda V1s \in ($

**Definition 20** We define  $c\_2Ebool\_2E\_5C\_2F$  to be  $(\lambda V0t1 \in 2. (\lambda V1t2 \in 2. (ap\ (c\_2Ebool\_2E\_21\ 2)\ (\lambda V2t \in$

**Definition 21** We define  $c\_2Epred\_set\_2EUNION$  to be  $\lambda A\_27a : \iota. \lambda V0s \in (2^{A-27a}). \lambda V1t \in (2^{A-27a}). (ap\ (c$

**Definition 22** We define  $c\_2Ereal\_topology\_2Eclosure$  to be  $\lambda V0s \in (2^{ty\_2Erealax\_2Ereal}). (ap\ (ap\ (c\_2Epred\_s$

**Definition 23** We define  $c\_2Epred\_set\_2ESUBSET$  to be  $\lambda A\_27a : \iota. \lambda V0s \in (2^{A-27a}). \lambda V1t \in (2^{A-27a}). (ap\ ($

**Definition 24** We define  $c\_2Epred\_set\_2EBIGINTER$  to be  $\lambda A\_27a : \iota. \lambda V0P \in (2^{(2^{A-27a})}). (ap\ (c\_2Epred\_s$

**Definition 25** We define  $c\_2Etopology\_2Ehull$  to be  $\lambda A\_27a : \iota. \lambda V0P \in (2^{(2^{A-27a})}). \lambda V1s \in (2^{A-27a}). (ap\ (c$

Assume the following.

$$True \tag{13}$$

Assume the following.

$$\forall A\_27a. nonempty\ A\_27a \Rightarrow (\forall V0t \in 2. ((\forall V1x \in A\_27a. (p\ V0t) \Leftrightarrow (p\ V0t))) \tag{14}$$

Assume the following.

$$\begin{aligned} & (\forall V0t \in 2. (((True \Leftrightarrow (p\ V0t)) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \Leftrightarrow True) \Leftrightarrow \\ & (p\ V0t)) \wedge (((False \Leftrightarrow (p\ V0t)) \Leftrightarrow \neg(p\ V0t)) \wedge (((p\ V0t) \Leftrightarrow False) \Leftrightarrow \neg( \\ & p\ V0t)))))) \end{aligned} \tag{15}$$

Assume the following.

$$\begin{aligned} & (\forall V0s \in (2^{ty\_2Erealax\_2Ereal}). ((ap\ c\_2Ereal\_topology\_2Eclosure \\ & V0s) = (ap\ (ap\ (c\_2Etopology\_2Ehull\ ty\_2Erealax\_2Ereal)\ c\_2Ereal\_topology\_2EClosed \\ & V0s))) \end{aligned} \tag{16}$$

Assume the following.

$$\begin{aligned} & \forall A\_27a. nonempty\ A\_27a \Rightarrow (\forall V0P \in (2^{(2^{A-27a})}). (\forall V1s \in \\ & (2^{A-27a}). (\forall V2t \in (2^{A-27a}). (((p\ (ap\ (ap\ (c\_2Epred\_set\_2ESUBSET \\ & A\_27a)\ V1s)\ V2t)) \wedge (p\ (ap\ V0P\ V2t))) \Rightarrow (p\ (ap\ (ap\ (c\_2Epred\_set\_2ESUBSET \\ & A\_27a)\ (ap\ (ap\ (c\_2Etopology\_2Ehull\ A\_27a)\ V0P)\ V1s))\ V2t)))))) \end{aligned} \tag{17}$$

**Theorem 1**

$$\begin{aligned} & (\forall V0s \in (2^{ty\_2Erealax\_2Ereal}). (\forall V1t \in (2^{ty\_2Erealax\_2Ereal}). \\ & (((p\ (ap\ (ap\ (c\_2Epred\_set\_2ESUBSET\ ty\_2Erealax\_2Ereal)\ V0s) \\ & V1t)) \wedge (p\ (ap\ c\_2Ereal\_topology\_2EClosed\ V1t))) \Rightarrow (p\ (ap\ (ap\ (c\_2Epred\_set\_2ESUBSET \\ & ty\_2Erealax\_2Ereal)\ (ap\ c\_2Ereal\_topology\_2Eclosure\ V0s) \\ & V1t)))))) \end{aligned}$$