

thm_2Ereal__topology_2ECOMPONENTS__EMPTY
 (TMKPRYULjJAiQF-
 SRg1QfKE2jn5RiGqJyKYW)

October 26, 2020

Definition 1 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define $c_2Ebool_2E_ET$ to be $(ap (ap (c_2Emin_2E_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

Definition 3 We define $c_2Ebool_2E_21$ to be $\lambda A_27a : \iota.(\lambda V0P \in (2^{A_27a}).(ap (ap (c_2Emin_2E_3D (2^{A_27a}))$

Definition 4 We define $c_2Ebool_2E_EF$ to be $(ap (c_2Ebool_2E_21 2) (\lambda V0t \in 2.V0t))$.

Definition 5 We define $c_2Epred_set_2EEMPTY$ to be $\lambda A_27a : \iota.(\lambda V0x \in A_27a.c_2Ebool_2E_EF)$.

Let $ty_2Erealax_2Ereal : \iota$ be given. Assume the following.

$$nonempty\ ty_2Erealax_2Ereal \tag{1}$$

Definition 6 We define $c_2Ebool_2E_IN$ to be $\lambda A_27a : \iota.(\lambda V0x \in A_27a.(\lambda V1f \in (2^{A_27a}).(ap V1f V0x)))$

Definition 7 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p P \Rightarrow p Q)$ of type ι .

Definition 8 We define $c_2Ebool_2E_2F_5C$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21 2) (\lambda V2t \in 2.V2t)))$

Definition 9 We define $c_2Epred_set_2ESUBSET$ to be $\lambda A_27a : \iota.\lambda V0s \in (2^{A_27a}).\lambda V1t \in (2^{A_27a}).(ap (c_2Emin_2E_3D_3D_3E$

Let $ty_2Epair_2Eprod : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A0.nonempty\ A0 \Rightarrow \forall A1.nonempty\ A1 \Rightarrow nonempty\ (ty_2Epair_2Eprod\ A0\ A1) \tag{2}$$

Let $c_2Epair_2EABS_prod : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow \forall A_27b.nonempty\ A_27b \Rightarrow c_2Epair_2EABS_prod\ A_27a\ A_27b \in ((ty_2Epair_2Eprod\ A_27a\ A_27b)^{(2^{A_27b})^{A_27a}}) \tag{3}$$

Definition 10 We define $c_2Epair_2E_2C$ to be $\lambda A_27a : \iota. \lambda A_27b : \iota. \lambda V0x \in A_27a. \lambda V1y \in A_27b. (ap (c_2Epred_set_2EGSPEC : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A_27a. nonempty A_27a \Rightarrow \forall A_27b. nonempty A_27b \Rightarrow c_2Epred_set_2EGSPEC \\ A_27a A_27b \in ((2^{A_27a})^{(ty_2Epair_2Eprod A_27a 2)^{A_27b}})$$
(4)

Definition 11 We define $c_2Epred_set_2EINTER$ to be $\lambda A_27a : \iota. \lambda V0s \in (2^{A_27a}). \lambda V1t \in (2^{A_27a}). (ap (c_2Ebool_2E_7E$

Definition 12 We define $c_2Ebool_2E_7E$ to be $(\lambda V0t \in 2. (ap (ap c_2Emin_2E_3D_3D_3E V0t) c_2Ebool_2E_7E$

Definition 13 We define $c_2Ebool_2E_5C_2F$ to be $(\lambda V0t1 \in 2. (\lambda V1t2 \in 2. (ap (c_2Ebool_2E_21 2) (\lambda V2t \in 2$

Definition 14 We define $c_2Epred_set_2EUNION$ to be $\lambda A_27a : \iota. \lambda V0s \in (2^{A_27a}). \lambda V1t \in (2^{A_27a}). (ap (c_2Ebool_2E_7E$

Let $c_2Ereal_topology_2EDist : \iota$ be given. Assume the following.

$$c_2Ereal_topology_2EDist \in (ty_2Erealax_2Ereal^{(ty_2Epair_2Eprod ty_2Erealax_2Ereal ty_2Erealax_2Ereal)})$$
(5)

Let $ty_2Ehreal_2Ehreal : \iota$ be given. Assume the following.

$$nonempty ty_2Ehreal_2Ehreal$$
(6)

Let $c_2Erealax_2Ereal_REP_CLASS : \iota$ be given. Assume the following.

$$c_2Erealax_2Ereal_REP_CLASS \in ((2^{(ty_2Epair_2Eprod ty_2Ehreal_2Ehreal ty_2Ehreal_2Ehreal)})^{ty_2Erealax_2Ereal})$$
(7)

Definition 15 We define $c_2Emin_2E_40$ to be $\lambda A. \lambda P \in 2^A. \mathbf{if} (\exists x \in A. p (ap P x)) \mathbf{then} (the (\lambda x. x \in A \wedge P x))$ of type $\iota \Rightarrow \iota$.

Definition 16 We define $c_2Erealax_2Ereal_REP$ to be $\lambda V0a \in ty_2Erealax_2Ereal. (ap (c_2Emin_2E_40 (the (\lambda x. x \in A \wedge P x))))$

Let $c_2Erealax_2Etrealt_lt : \iota$ be given. Assume the following.

$$c_2Erealax_2Etrealt_lt \in ((2^{(ty_2Epair_2Eprod ty_2Ehreal_2Ehreal ty_2Ehreal_2Ehreal)})^{(ty_2Epair_2Eprod ty_2Ehreal_2Ehreal)})$$
(8)

Definition 17 We define $c_2Erealax_2Ereal_lt$ to be $\lambda V0T1 \in ty_2Erealax_2Ereal. \lambda V1T2 \in ty_2Erealax_2Ereal. (ap (c_2Emin_2E_40 (the (\lambda x. x \in A \wedge P x))))$

Let $c_2Enum_2EZERO_REP : \iota$ be given. Assume the following.

$$c_2Enum_2EZERO_REP \in \omega$$
(9)

Let $ty_2Enum_2Enum : \iota$ be given. Assume the following.

$$nonempty ty_2Enum_2Enum$$
(10)

Let $c_2Enum_2EABS_num : \iota$ be given. Assume the following.

$$c_2Enum_2EABS_num \in (ty_2Enum_2Enum^{\omega})$$
(11)

Definition 18 We define c_Enum_E0 to be $(ap\ c_Enum_EABS_num\ c_Enum_EZERO_REP)$.

Let $c_Ereal_Ereal_of_num : \iota$ be given. Assume the following.

$$c_Ereal_Ereal_of_num \in (ty_Erealax_Ereal^{ty_Eenum_Eenum}) \quad (12)$$

Definition 19 We define c_Ebool_E3F to be $\lambda A_27a : \iota. (\lambda V0P \in (2^{A_27a}). (ap\ V0P\ (ap\ (c_Emin_E40$

Definition 20 We define $c_Ereal_topology_EOpen$ to be $\lambda V0s \in (2^{ty_Erealax_Ereal}). (ap\ (c_Ebool_E2$

Definition 21 We define $c_Ereal_topology_Econnected$ to be $\lambda V0s \in (2^{ty_Erealax_Ereal}). (ap\ c_Ebool_E2$

Definition 22 We define $c_Ereal_topology_Econnected_component$ to be $\lambda V0s \in (2^{ty_Erealax_Ereal}). \lambda V$

Definition 23 We define $c_Ereal_topology_Ecomponents$ to be $\lambda V0s \in (2^{ty_Erealax_Ereal}). (ap\ (c_Epre$

Assume the following.

$$True \quad (13)$$

Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow (\forall V0x \in A_27a. ((V0x = V0x) \Leftrightarrow True)) \quad (14)$$

Assume the following.

$$(\forall V0s \in (2^{ty_Erealax_Ereal}). (((ap\ c_Ereal_topology_Ecomponents\ V0s) = (c_Epred_set_EEMPTY\ (2^{ty_Erealax_Ereal}))) \Leftrightarrow (V0s = (c_Epred_set_EEMPTY\ ty_Erealax_Ereal)))) \quad (15)$$

Theorem 1

$$((ap\ c_Ereal_topology_Ecomponents\ (c_Epred_set_EEMPTY\ ty_Erealax_Ereal)) = (c_Epred_set_EEMPTY\ (2^{ty_Erealax_Ereal})))$$