

thm_2Ereal__topology_2ECONTINUOUS__AT__AVOID (TMMJ841ujDYWzvpKUtBdzgbexqtgpoQ5XpM)

October 26, 2020

Definition 1 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define $c_2Ebool_2E_2ET$ to be $(ap (ap (c_2Emin_2E_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

Definition 3 We define $c_2Ebool_2E_2E21$ to be $\lambda A_27a : \iota.(\lambda V0P \in (2^{A_27a}).(ap (ap (c_2Emin_2E_3D (2^{A_27a}))$

Definition 4 We define $c_2Ebool_2E_2EF$ to be $(ap (c_2Ebool_2E_2E21 2) (\lambda V0t \in 2.V0t))$.

Definition 5 We define $c_2Epred_set_2EUNIV$ to be $\lambda A_27a : \iota.(\lambda V0x \in A_27a.c_2Ebool_2E_2ET)$.

Let $ty_2Ereal_topology_2E_2Enet : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A0.nonempty A0 \Rightarrow nonempty (ty_2Ereal_topology_2E_2Enet A0) \quad (1)$$

Definition 6 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p P \Rightarrow p Q)$ of type ι .

Definition 7 We define $c_2Ebool_2E_2E2F_5C$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_2E21 2) (\lambda V2t \in 2.V2t))$

Let $ty_2Epair_2E_2Eprod : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A0.nonempty A0 \Rightarrow \forall A1.nonempty A1 \Rightarrow nonempty (ty_2Epair_2E_2Eprod A0 A1) \quad (2)$$

Let $c_2Epair_2E_2EABS_prod : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A_27a.nonempty A_27a \Rightarrow \forall A_27b.nonempty A_27b \Rightarrow c_2Epair_2E_2EABS_prod A_27a A_27b \in ((ty_2Epair_2E_2Eprod A_27a A_27b)^{(2^{A_27b})^{A_27a}}) \quad (3)$$

Definition 8 We define $c_2Epair_2E_2E2C$ to be $\lambda A_27a : \iota.\lambda A_27b : \iota.\lambda V0x \in A_27a.\lambda V1y \in A_27b.(ap (c_2E$

Let $ty_2Erealax_2Ereal : \iota$ be given. Assume the following.

$$nonempty\ ty_2Erealax_2Ereal \quad (4)$$

Let $c_2Ereal_topology_2EDist : \iota$ be given. Assume the following.

$$c_2Ereal_topology_2EDist \in (ty_2Erealax_2Ereal^{(ty_2Epair_2Eprod\ ty_2Erealax_2Ereal\ ty_2Erealax_2Ereal)}) \quad (5)$$

Let $c_2Enum_2EZERO_REP : \iota$ be given. Assume the following.

$$c_2Enum_2EZERO_REP \in \omega \quad (6)$$

Let $ty_2Enum_2Enum : \iota$ be given. Assume the following.

$$nonempty\ ty_2Enum_2Enum \quad (7)$$

Let $c_2Enum_2EABS_num : \iota$ be given. Assume the following.

$$c_2Enum_2EABS_num \in (ty_2Enum_2Enum^{\omega}) \quad (8)$$

Definition 9 We define c_2Enum_2E0 to be $(ap\ c_2Enum_2EABS_num\ c_2Enum_2EZERO_REP)$.

Let $c_2Ereal_2Ereal_of_num : \iota$ be given. Assume the following.

$$c_2Ereal_2Ereal_of_num \in (ty_2Erealax_2Ereal^{ty_2Enum_2Enum}) \quad (9)$$

Let $ty_2Ehreal_2Ehreal : \iota$ be given. Assume the following.

$$nonempty\ ty_2Ehreal_2Ehreal \quad (10)$$

Let $c_2Erealax_2Ereal_REP_CLASS : \iota$ be given. Assume the following.

$$c_2Erealax_2Ereal_REP_CLASS \in ((2^{(ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal\ ty_2Ehreal_2Ehreal)})^{ty_2Erealax_2Ereal}) \quad (11)$$

Definition 10 We define c_2Emin_2E40 to be $\lambda A.\lambda P \in 2^A.\text{if } (\exists x \in A.p\ (ap\ P\ x)) \text{ then } (the\ (\lambda x.x \in A \wedge p\ x)) \text{ of type } \iota \Rightarrow \iota$.

Definition 11 We define $c_2Erealax_2Ereal_REP$ to be $\lambda V0a \in ty_2Erealax_2Ereal.(ap\ (c_2Emin_2E40\ (ap\ V0a)))$

Let $c_2Erealax_2Etreall_lt : \iota$ be given. Assume the following.

$$c_2Erealax_2Etreall_lt \in ((2^{(ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal\ ty_2Ehreal_2Ehreal)})^{(ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal)}) \quad (12)$$

Definition 12 We define $c_2Erealax_2Ereal_lt$ to be $\lambda V0T1 \in ty_2Erealax_2Ereal.\lambda V1T2 \in ty_2Erealax_2Ereal.$

Definition 13 We define c_2Ebool_2E3F to be $\lambda A.\lambda 27a : \iota.(\lambda V0P \in (2^{A-27a}).(ap\ V0P\ (ap\ (c_2Emin_2E40\ (ap\ V0a))))$

Definition 14 We define c_2Ebool_2E7E to be $(\lambda V0t \in 2.(ap\ (ap\ c_2Emin_2E3D_3D_3E\ V0t)\ c_2Ebool_2E3F))$

Definition 15 We define c_2Ebool_2EIN to be $\lambda A_27a : \iota. (\lambda V0x \in A_27a. (\lambda V1f \in (2^{A_27a}). (ap\ V1f\ V0x)))$

Definition 16 We define $c_2Ereal_2Ereal_lte$ to be $\lambda V0x \in ty_2Erealax_2Ereal. \lambda V1y \in ty_2Erealax_2Ereal$

Let $c_2Ereal_topology_2Emk_net : \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned} \forall A_27a. nonempty\ A_27a \Rightarrow c_2Ereal_topology_2Emk_net \\ A_27a \in ((ty_2Ereal_topology_2Enet\ A_27a)^{(2^{A_27a})^{A_27a}}) \end{aligned} \quad (13)$$

Definition 17 We define $c_2Ereal_topology_2Eat$ to be $\lambda V0a \in ty_2Erealax_2Ereal. (ap\ (c_2Ereal_topology_2Eat\ V0a))$

Let $c_2Ereal_topology_2Enetord : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A_27a. nonempty\ A_27a \Rightarrow c_2Ereal_topology_2Enetord\ A_27a \in ((2^{A_27a})^{A_27a})^{(ty_2Ereal_topology_2Enet\ A_27a)} \quad (14)$$

Definition 18 We define $c_2Ereal_topology_2Ewithin$ to be $\lambda A_27a : \iota. \lambda V0net \in (ty_2Ereal_topology_2Ewithin\ A_27a\ V0net)$

Definition 19 We define $c_2Ereal_topology_2Eenetlimit$ to be $\lambda A_27a : \iota. \lambda V0net \in (ty_2Ereal_topology_2Eenetlimit\ A_27a\ V0net)$

Definition 20 We define $c_2Ebool_2E5C_2F$ to be $(\lambda V0t1 \in 2. (\lambda V1t2 \in 2. (ap\ (c_2Ebool_2E21\ 2)\ (\lambda V2t \in 2. (c_2Ebool_2E5C_2F\ V0t1\ V1t2\ V2t))))$

Definition 21 We define $c_2Ereal_topology_2Etrivial_limit$ to be $\lambda A_27a : \iota. \lambda V0net \in (ty_2Ereal_topology_2Etrivial_limit\ A_27a\ V0net)$

Definition 22 We define $c_2Ereal_topology_2Eeventually$ to be $\lambda A_27a : \iota. \lambda V0p \in (2^{A_27a}). \lambda V1net \in (ty_2Ereal_topology_2Eeventually\ A_27a\ V0p\ V1net)$

Definition 23 We define $c_2Ereal_topology_2E2D_2D_3E$ to be $\lambda A_27a : \iota. \lambda V0f \in (ty_2Erealax_2Ereal^A\ A_27a\ V0f)$

Definition 24 We define $c_2Ereal_topology_2Econtinuous$ to be $\lambda A_27a : \iota. \lambda V0f \in (ty_2Erealax_2Ereal^A\ A_27a\ V0f)$

Assume the following.

$$True \quad (15)$$

Assume the following.

$$\begin{aligned} (\forall V0t \in 2. (((True \wedge (p\ V0t)) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \wedge True) \Leftrightarrow \\ (p\ V0t)) \wedge (((False \wedge (p\ V0t)) \Leftrightarrow False) \wedge (((p\ V0t) \wedge False) \Leftrightarrow False) \wedge \\ (((p\ V0t) \wedge (p\ V0t)) \Leftrightarrow (p\ V0t)))))) \end{aligned} \quad (16)$$

Assume the following.

$$\begin{aligned} (\forall V0t \in 2. (((True \Rightarrow (p\ V0t)) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \Rightarrow True) \Leftrightarrow \\ True) \wedge (((False \Rightarrow (p\ V0t)) \Leftrightarrow True) \wedge (((p\ V0t) \Rightarrow (p\ V0t)) \Leftrightarrow True) \wedge ((\\ (p\ V0t) \Rightarrow False) \Leftrightarrow (\neg (p\ V0t)))))) \end{aligned} \quad (17)$$

Assume the following.

$$\begin{aligned} (\forall V0t \in 2. (((True \Leftrightarrow (p\ V0t)) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \Leftrightarrow True) \Leftrightarrow \\ (p\ V0t)) \wedge (((False \Leftrightarrow (p\ V0t)) \Leftrightarrow (\neg (p\ V0t))) \wedge (((p\ V0t) \Leftrightarrow False) \Leftrightarrow (\neg \\ (p\ V0t)))))) \end{aligned} \quad (18)$$

Assume the following.

$$\forall A.27a.nonempty A.27a \Rightarrow (\forall V0x \in A.27a. (p (ap (ap (c.2Ebool.2EIN A.27a) V0x) (c.2Epred_set.2EUNIV A.27a)))) \quad (19)$$

Assume the following.

$$(\forall V0x \in ty.2Erealax.2Ereal. ((ap (ap (c.2Ereal_topology.2Ewithin ty.2Erealax.2Ereal) (ap c.2Ereal_topology.2Eat V0x)) (c.2Epred_set.2EUNIV ty.2Erealax.2Ereal)) = (ap c.2Ereal_topology.2Eat V0x))) \quad (20)$$

Assume the following.

$$\begin{aligned} & (\forall V0f \in (ty.2Erealax.2Ereal^{ty.2Erealax.2Ereal}). (\forall V1x \in \\ & \quad ty.2Erealax.2Ereal. (\forall V2s \in (2^{ty.2Erealax.2Ereal}). (\\ \forall V3a \in ty.2Erealax.2Ereal. ((p (ap (ap (c.2Ereal_topology.2Econtinuous \\ & \quad ty.2Erealax.2Ereal) V0f) (ap (ap (c.2Ereal_topology.2Ewithin \\ & \quad ty.2Erealax.2Ereal) (ap c.2Ereal_topology.2Eat V1x)) V2s))) \wedge \\ & \quad ((p (ap (ap (c.2Ebool.2EIN ty.2Erealax.2Ereal) V1x) V2s)) \wedge (\neg \\ & \quad (ap V0f V1x) = V3a)))) \Rightarrow (\exists V4e \in ty.2Erealax.2Ereal. ((p (ap \\ & \quad (ap c.2Erealax.2Ereal_lt (ap c.2Ereal.2Ereal_of_num c.2Enum.2E0)) \\ & \quad V4e)) \wedge (\forall V5y \in ty.2Erealax.2Ereal. ((p (ap (ap (c.2Ebool.2EIN \\ & \quad ty.2Erealax.2Ereal) V5y) V2s)) \wedge (p (ap (ap c.2Erealax.2Ereal_lt \\ & \quad (ap c.2Ereal_topology.2EDist (ap (ap (c.2Epair.2E.2C ty.2Erealax.2Ereal \\ & \quad ty.2Erealax.2Ereal) V1x) V5y))) V4e))) \Rightarrow (\neg ((ap V0f V5y) = V3a)))))) \quad (21) \end{aligned}$$

Theorem 1

$$\begin{aligned} & (\forall V0f \in (ty.2Erealax.2Ereal^{ty.2Erealax.2Ereal}). (\forall V1x \in \\ & \quad ty.2Erealax.2Ereal. (\forall V2a \in ty.2Erealax.2Ereal. ((p (\\ & \quad ap (ap (c.2Ereal_topology.2Econtinuous ty.2Erealax.2Ereal) \\ & \quad V0f) (ap c.2Ereal_topology.2Eat V1x))) \wedge (\neg ((ap V0f V1x) = V2a))) \Rightarrow \\ & \quad (\exists V3e \in ty.2Erealax.2Ereal. ((p (ap (ap c.2Erealax.2Ereal_lt \\ & \quad (ap c.2Ereal.2Ereal_of_num c.2Enum.2E0)) V3e)) \wedge (\forall V4y \in \\ & \quad ty.2Erealax.2Ereal. ((p (ap (ap c.2Erealax.2Ereal_lt (ap c.2Ereal_topology.2EDist \\ & \quad (ap (ap (c.2Epair.2E.2C ty.2Erealax.2Ereal ty.2Erealax.2Ereal) \\ & \quad V1x) V4y))) V3e))) \Rightarrow (\neg ((ap V0f V4y) = V2a)))))) \end{aligned}$$