

# thm\_2Ereal\_topology\_2ECONTINUOUS\_NEG (TMVL515UXrgvzd7fLFz6q8rfaFMUfiKr9br)

October 26, 2020

**Definition 1** We define  $c\_2Emin\_2E\_3D$  to be  $\lambda A.\lambda x \in A.\lambda y \in A.inj\_o (x = y)$  of type  $\iota \Rightarrow \iota$ .

**Definition 2** We define  $c\_2Ebool\_2E\_ET$  to be  $(ap (ap (c\_2Emin\_2E\_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

**Definition 3** We define  $c\_2Ebool\_2E\_21$  to be  $\lambda A.\lambda P \in 2^A.(ap (ap (c\_2Emin\_2E\_3D (2^{A-27a})) (\lambda V0t \in 2.V0t)) (\lambda V1t \in 2.V1t))$

**Definition 4** We define  $c\_2Ebool\_2E\_EF$  to be  $(ap (c\_2Ebool\_2E\_21 2) (\lambda V0t \in 2.V0t))$ .

**Definition 5** We define  $c\_2Emin\_2E\_3D\_3D\_3E$  to be  $\lambda P \in 2.\lambda Q \in 2.inj\_o (p P \Rightarrow p Q)$  of type  $\iota$ .

**Definition 6** We define  $c\_2Ebool\_2E\_7E$  to be  $(\lambda V0t \in 2.(ap (ap c\_2Emin\_2E\_3D\_3D\_3E V0t) c\_2Ebool\_2E\_EF))$

**Definition 7** We define  $c\_2Ebool\_2E\_2F\_5C$  to be  $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c\_2Ebool\_2E\_21 2) (\lambda V2t \in 2.V2t)) (\lambda V3t \in 2.V3t)))$

Let  $ty\_2Ehreal\_2Ehreal : \iota$  be given. Assume the following.

$$nonempty\ ty\_2Ehreal\_2Ehreal \tag{1}$$

Let  $ty\_2Epair\_2Eprod : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A0.nonempty\ A0 \Rightarrow \forall A1.nonempty\ A1 \Rightarrow nonempty\ (ty\_2Epair\_2Eprod\ A0\ A1) \tag{2}$$

Let  $ty\_2Erealax\_2Ereal : \iota$  be given. Assume the following.

$$nonempty\ ty\_2Erealax\_2Ereal \tag{3}$$

Let  $c\_2Erealax\_2Ereal\_REP\_CLASS : \iota$  be given. Assume the following.

$$c\_2Erealax\_2Ereal\_REP\_CLASS \in ((2^{(ty\_2Epair\_2Eprod\ ty\_2Ehreal\_2Ehreal\ ty\_2Ehreal\_2Ehreal)})\ ty\_2Erealax\_2Ereal) \tag{4}$$

**Definition 8** We define  $c\_2Emin\_2E\_40$  to be  $\lambda A.\lambda P \in 2^A.if (\exists x \in A.p (ap P x)) \mathbf{then} (the (\lambda x.x \in A \wedge p x))$  of type  $\iota \Rightarrow \iota$ .

**Definition 9** We define  $c\_2Erealax\_2Ereal\_REP$  to be  $\lambda V0a \in ty\_2Erealax\_2Ereal.(ap (c\_2Emin\_2E40 (ty$

Let  $c\_2Erealax\_2Etrealm\_neg : \iota$  be given. Assume the following.

$$c\_2Erealax\_2Etrealm\_neg \in ((ty\_2Epair\_2Eprod ty\_2Ehreal\_2Ehreal ty\_2Ehreal\_2Ehreal)(ty\_2Epair\_2Eprod ty\_2Ehreal\_2Ehreal ty\_2Ehreal\_2Ehreal)) \quad (5)$$

Let  $c\_2Erealax\_2Etrealm\_eq : \iota$  be given. Assume the following.

$$c\_2Erealax\_2Etrealm\_eq \in ((2^{(ty\_2Epair\_2Eprod ty\_2Ehreal\_2Ehreal ty\_2Ehreal\_2Ehreal)})(ty\_2Epair\_2Eprod ty\_2Ehreal\_2Ehreal ty\_2Ehreal\_2Ehreal)) \quad (6)$$

Let  $c\_2Erealax\_2Ereal\_ABS\_CLASS : \iota$  be given. Assume the following.

$$c\_2Erealax\_2Ereal\_ABS\_CLASS \in (ty\_2Erealax\_2Ereal^{(2^{(ty\_2Epair\_2Eprod ty\_2Ehreal\_2Ehreal ty\_2Ehreal\_2Ehreal)})(ty\_2Epair\_2Eprod ty\_2Ehreal\_2Ehreal ty\_2Ehreal\_2Ehreal)}) \quad (7)$$

**Definition 10** We define  $c\_2Erealax\_2Ereal\_ABS$  to be  $\lambda V0r \in (ty\_2Epair\_2Eprod ty\_2Ehreal\_2Ehreal ty\_2Ehreal\_2Ehreal)$

**Definition 11** We define  $c\_2Erealax\_2Ereal\_neg$  to be  $\lambda V0T1 \in ty\_2Erealax\_2Ereal.(ap c\_2Erealax\_2Ereal$

Let  $ty\_2Ereal\_topology\_2Enet : \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A0.nonempty A0 \Rightarrow nonempty (ty\_2Ereal\_topology\_2Enet A0) \quad (8)$$

Let  $c\_2Ereal\_topology\_2Enetord : \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A\_27a.nonempty A\_27a \Rightarrow c\_2Ereal\_topology\_2Enetord A\_27a \in (((2^{A\_27a})^{A\_27a})(ty\_2Ereal\_topology\_2Enet A\_27a)) \quad (9)$$

**Definition 12** We define  $c\_2Ereal\_topology\_2Enetlimit$  to be  $\lambda A\_27a : \iota.\lambda V0net \in (ty\_2Ereal\_topology\_2E$

Let  $c\_2Epair\_2EABS\_prod : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A\_27a.nonempty A\_27a \Rightarrow \forall A\_27b.nonempty A\_27b \Rightarrow c\_2Epair\_2EABS\_prod A\_27a A\_27b \in ((ty\_2Epair\_2Eprod A\_27a A\_27b)^{(2^{A\_27b})^{A\_27a}}) \quad (10)$$

**Definition 13** We define  $c\_2Epair\_2E2C$  to be  $\lambda A\_27a : \iota.\lambda A\_27b : \iota.\lambda V0x \in A\_27a.\lambda V1y \in A\_27b.(ap (c\_2$

Let  $c\_2Ereal\_topology\_2EDist : \iota$  be given. Assume the following.

$$c\_2Ereal\_topology\_2EDist \in (ty\_2Erealax\_2Ereal^{(ty\_2Epair\_2Eprod ty\_2Erealax\_2Ereal ty\_2Erealax\_2Ereal)}) \quad (11)$$

Let  $c\_2Erealax\_2Etrealm\_lt : \iota$  be given. Assume the following.

$$c\_2Erealax\_2Etrealm\_lt \in ((2^{(ty\_2Epair\_2Eprod ty\_2Ehreal\_2Ehreal ty\_2Ehreal\_2Ehreal)})(ty\_2Epair\_2Eprod ty\_2Ehreal\_2Ehreal ty\_2Ehreal\_2Ehreal)) \quad (12)$$

**Definition 14** We define  $c\_2Erealax\_2Ereal\_lt$  to be  $\lambda V0T1 \in ty\_2Erealax\_2Ereal.\lambda V1T2 \in ty\_2Erealax\_2E$

**Definition 15** We define  $c\_2Ebool\_2E3F$  to be  $\lambda A\_27a : \iota.(\lambda V0P \in (2^{A\_27a}).(ap V0P (ap (c\_2Emin\_2E40$

**Definition 16** We define  $c\_Ebool\_2E\_5C\_2F$  to be  $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c\_Ebool\_2E\_21) 2) (\lambda V2t \in$

**Definition 17** We define  $c\_Ereal\_topology\_2Etrivial\_limit$  to be  $\lambda A\_27a : \iota.\lambda V0net \in (ty\_2Ereal\_topology$

**Definition 18** We define  $c\_Ereal\_topology\_2Eeventually$  to be  $\lambda A\_27a : \iota.\lambda V0p \in (2^{A\_27a}).\lambda V1net \in (ty\_2$

Let  $c\_2Enum\_2EZERO\_REP : \iota$  be given. Assume the following.

$$c\_2Enum\_2EZERO\_REP \in \omega \tag{13}$$

Let  $ty\_2Enum\_2Enum : \iota$  be given. Assume the following.

$$nonempty\ ty\_2Enum\_2Enum \tag{14}$$

Let  $c\_2Enum\_2EABS\_num : \iota$  be given. Assume the following.

$$c\_2Enum\_2EABS\_num \in (ty\_2Enum\_2Enum^{\omega}) \tag{15}$$

**Definition 19** We define  $c\_2Enum\_2E0$  to be  $(ap\ c\_2Enum\_2EABS\_num\ c\_2Enum\_2EZERO\_REP)$ .

Let  $c\_2Ereal\_2Ereal\_of\_num : \iota$  be given. Assume the following.

$$c\_2Ereal\_2Ereal\_of\_num \in (ty\_2Erealax\_2Ereal^{ty\_2Enum\_2Enum}) \tag{16}$$

**Definition 20** We define  $c\_2Ereal\_topology\_2E\_2D\_2D\_3E$  to be  $\lambda A\_27a : \iota.\lambda V0f \in (ty\_2Erealax\_2Ereal^A$

**Definition 21** We define  $c\_2Ereal\_topology\_2Econtinuous$  to be  $\lambda A\_27a : \iota.\lambda V0f \in (ty\_2Erealax\_2Ereal^A$

Assume the following.

$$True \tag{17}$$

Assume the following.

$$\forall A\_27a.nonempty\ A\_27a \Rightarrow (\forall V0t \in 2.((\forall V1x \in A\_27a.(p\ V0t)) \Leftrightarrow (p\ V0t))) \tag{18}$$

Assume the following.

$$\begin{aligned} & (\forall V0t \in 2.(((True \Rightarrow (p\ V0t)) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \Rightarrow True) \Leftrightarrow \\ & True) \wedge (((False \Rightarrow (p\ V0t)) \Leftrightarrow True) \wedge (((p\ V0t) \Rightarrow (p\ V0t)) \Leftrightarrow True) \wedge (( \\ & (p\ V0t) \Rightarrow False) \Leftrightarrow (\neg (p\ V0t)))))) \end{aligned} \tag{19}$$

Assume the following.

$$\forall A\_27a.nonempty\ A\_27a \Rightarrow (\forall V0x \in A\_27a.(\forall V1y \in A\_27a.((V0x = V1y) \Leftrightarrow (V1y = V0x)))) \tag{20}$$

Assume the following.

$$\begin{aligned} & (\forall V0t \in 2.(((True \Leftrightarrow (p\ V0t)) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \Leftrightarrow True) \Leftrightarrow \\ & (p\ V0t)) \wedge (((False \Leftrightarrow (p\ V0t)) \Leftrightarrow (\neg (p\ V0t))) \wedge (((p\ V0t) \Leftrightarrow False) \Leftrightarrow (\neg ( \\ & p\ V0t)))))) \end{aligned} \tag{21}$$

Assume the following.

$$(\forall V0t1 \in 2.(\forall V1t2 \in 2.(\forall V2t3 \in 2.(((p V0t1) \Rightarrow ((p V1t2) \Rightarrow (p V2t3))) \Leftrightarrow (((p V0t1) \wedge (p V1t2)) \Rightarrow (p V2t3)))))) \quad (22)$$

Assume the following.

$$(\forall V0x \in 2.(\forall V1x_{.27} \in 2.(\forall V2y \in 2.(\forall V3y_{.27} \in 2.(((p V0x) \Leftrightarrow (p V1x_{.27})) \wedge ((p V1x_{.27}) \Rightarrow ((p V2y) \Leftrightarrow (p V3y_{.27})))))) \Rightarrow ((p V0x) \Rightarrow (p V2y)) \Leftrightarrow ((p V1x_{.27}) \Rightarrow (p V3y_{.27})))))) \quad (23)$$

Assume the following.

$$\begin{aligned} \forall A_{.27a}.nonempty A_{.27a} \Rightarrow (\forall V0net \in (ty\_2Ereal\_topology\_2Enet \\ A_{.27a}).(\forall V1f \in (ty\_2Erealax\_2Ereal^{A_{.27a}}).(\forall V2l \in \\ ty\_2Erealax\_2Ereal.((p (ap (ap (ap (c\_2Ereal\_topology\_2E\_2D\_2D\_3E \\ A_{.27a}) V1f) V2l) V0net)) \Rightarrow (p (ap (ap (ap (c\_2Ereal\_topology\_2E\_2D\_2D\_3E \\ A_{.27a}) (\lambda V3x \in A_{.27a}.(ap c\_2Erealax\_2Ereal\_neg (ap V1f V3x)))) \\ (ap c\_2Erealax\_2Ereal\_neg V2l)) V0net)))))) \end{aligned} \quad (24)$$

### Theorem 1

$$\begin{aligned} \forall A_{.27a}.nonempty A_{.27a} \Rightarrow (\forall V0f \in (ty\_2Erealax\_2Ereal^{A_{.27a}}). \\ (\forall V1net \in (ty\_2Ereal\_topology\_2Enet A_{.27a}).((p (ap (ap \\ (c\_2Ereal\_topology\_2Econtinuous A_{.27a}) V0f) V1net)) \Rightarrow (p (ap \\ (ap (c\_2Ereal\_topology\_2Econtinuous A_{.27a}) (\lambda V2x \in A_{.27a}. \\ (ap c\_2Erealax\_2Ereal\_neg (ap V0f V2x)))) V1net)))) \end{aligned}$$