

thm_2Ereal__topology_2ECONTINUOUS__ON__EQ__CONTINUOU (TMcv8mJrKQs3KNFmZHWqZNP8eKEAtYxP9Dd)

October 26, 2020

Definition 1 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define $c_2Ebool_2E_2T$ to be $(ap (ap (c_2Emin_2E_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

Let $ty_2Erealax_2Ereal : \iota$ be given. Assume the following.

$$nonempty\ ty_2Erealax_2Ereal \tag{1}$$

Definition 3 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p \Rightarrow p \Rightarrow Q)$ of type ι .

Definition 4 We define $c_2Ebool_2E_21$ to be $\lambda A_27a : \iota.(\lambda V0P \in (2^{A_27a}).(ap (ap (c_2Emin_2E_3D (2^{A_27a}))$

Definition 5 We define $c_2Ebool_2E_2F_5C$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21 2) (\lambda V2t \in 2.V2t)))$

Let $ty_2Epair_2Eprod : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A0.nonempty\ A0 \Rightarrow \forall A1.nonempty\ A1 \Rightarrow nonempty\ (ty_2Epair_2Eprod\ A0\ A1) \tag{2}$$

Let $c_2Epair_2EABS_prod : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow \forall A_27b.nonempty\ A_27b \Rightarrow c_2Epair_2EABS_prod\ A_27a\ A_27b \in ((ty_2Epair_2Eprod\ A_27a\ A_27b)^{(2^{A_27b})^{A_27a}}) \tag{3}$$

Definition 6 We define $c_2Epair_2E_2C$ to be $\lambda A_27a : \iota.\lambda A_27b : \iota.\lambda V0x \in A_27a.\lambda V1y \in A_27b.(ap (c_2Emin_2E_3D (2^{A_27a}))$

Let $c_2Ereal_topology_2EDist : \iota$ be given. Assume the following.

$$c_2Ereal_topology_2EDist \in (ty_2Erealax_2Ereal^{(ty_2Epair_2Eprod\ ty_2Erealax_2Ereal\ ty_2Erealax_2Ereal)}) \tag{4}$$

Let $ty_2Ehreal_2Ehreal : \iota$ be given. Assume the following.

$$nonempty\ ty_2Ehreal_2Ehreal \quad (5)$$

Let $c_2Erealax_2Ereal_REP_CLASS : \iota$ be given. Assume the following.

$$c_2Erealax_2Ereal_REP_CLASS \in ((2^{(ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal\ ty_2Ehreal_2Ehreal)})^{ty_2Erealax}) \quad (6)$$

Definition 7 We define c_2Emin_2E40 to be $\lambda A.\lambda P \in 2^A.$ if $(\exists x \in A.p (ap\ P\ x))$ then (the $(\lambda x.x \in A \wedge p$ of type $\iota \Rightarrow \iota$).

Definition 8 We define $c_2Erealax_2Ereal_REP$ to be $\lambda V0a \in ty_2Erealax_2Ereal.(ap\ (c_2Emin_2E40\ (ty$

Let $c_2Erealax_2Etrealt_lt : \iota$ be given. Assume the following.

$$c_2Erealax_2Etrealt_lt \in ((2^{(ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal\ ty_2Ehreal_2Ehreal)})^{(ty_2Epair_2Eprod\ ty_2Ehreal)}) \quad (7)$$

Definition 9 We define $c_2Erealax_2Ereal_lt$ to be $\lambda V0T1 \in ty_2Erealax_2Ereal.\lambda V1T2 \in ty_2Erealax_2Ereal$

Definition 10 We define c_2Ebool_2E21 to be $(ap\ (c_2Ebool_2E21\ 2)) (\lambda V0t \in 2.V0t)$.

Definition 11 We define c_2Ebool_2E7E to be $(\lambda V0t \in 2.(ap\ (ap\ c_2Emin_2E3D_3D_3E\ V0t)\ c_2Ebool_2E21))$

Definition 12 We define $c_2Ereal_2Ereal_lte$ to be $\lambda V0x \in ty_2Erealax_2Ereal.\lambda V1y \in ty_2Erealax_2Ereal$

Let $c_2Enum_2EZERO_REP : \iota$ be given. Assume the following.

$$c_2Enum_2EZERO_REP \in \omega \quad (8)$$

Let $ty_2Enum_2Enum : \iota$ be given. Assume the following.

$$nonempty\ ty_2Enum_2Enum \quad (9)$$

Let $c_2Enum_2EABS_num : \iota$ be given. Assume the following.

$$c_2Enum_2EABS_num \in (ty_2Enum_2Enum^{\omega}) \quad (10)$$

Definition 13 We define c_2Enum_2E0 to be $(ap\ c_2Enum_2EABS_num\ c_2Enum_2EZERO_REP)$.

Let $c_2Ereal_2Ereal_of_num : \iota$ be given. Assume the following.

$$c_2Ereal_2Ereal_of_num \in (ty_2Erealax_2Ereal^{ty_2Enum_2Enum}) \quad (11)$$

Let $ty_2Ereal_topology_2Enet : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A0.nonempty\ A0 \Rightarrow nonempty\ (ty_2Ereal_topology_2Enet\ A0) \quad (12)$$

Let $c_2Ereal_topology_2Emk_net : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A.27a.nonempty\ A.27a \Rightarrow c_2Ereal_topology_2Emk_net\ A.27a \in ((ty_2Ereal_topology_2Enet\ A.27a)^{(2^{A.27a})^{A.27a}}) \quad (13)$$

Definition 14 We define $c_2Ereal_topology_2Eat$ to be $\lambda V0a \in ty_2Erealax_2Ereal.(ap (c_2Ereal_topology_2Eat) V0a)$

Definition 15 We define c_2Ebool_2EIN to be $\lambda A_27a : \iota.(\lambda V0x \in A_27a.(\lambda V1f \in (2^{A_27a}).(ap V1f V0x)))$

Let $c_2Ereal_topology_2Eenetord : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A_27a.nonempty A_27a \Rightarrow c_2Ereal_topology_2Eenetord A_27a \in (((2^{A_27a})^{A_27a})^{(ty_2Ereal_topology_2Eenet A_27a)}) \quad (14)$$

Definition 16 We define $c_2Ereal_topology_2Ewithin$ to be $\lambda A_27a : \iota.\lambda V0net \in (ty_2Ereal_topology_2Ewithin A_27a V0net)$

Definition 17 We define $c_2Ereal_topology_2Eenlimit$ to be $\lambda A_27a : \iota.\lambda V0net \in (ty_2Ereal_topology_2Eenlimit A_27a V0net)$

Definition 18 We define $c_2Ebool_2E_3F$ to be $\lambda A_27a : \iota.(\lambda V0P \in (2^{A_27a}).(ap V0P (ap (c_2Emin_2E_40) V0P)))$

Definition 19 We define $c_2Ebool_2E_5C_2F$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21) V0t1) (\lambda V2t \in 2.V2t)))$

Definition 20 We define $c_2Ereal_topology_2Etrivial_limit$ to be $\lambda A_27a : \iota.\lambda V0net \in (ty_2Ereal_topology_2Etrivial_limit A_27a V0net)$

Definition 21 We define $c_2Ereal_topology_2Eeventually$ to be $\lambda A_27a : \iota.\lambda V0p \in (2^{A_27a}).\lambda V1net \in (ty_2Ereal_topology_2Eeventually A_27a V0p V1net)$

Definition 22 We define $c_2Ereal_topology_2E_2D_2D_3E$ to be $\lambda A_27a : \iota.\lambda V0f \in (ty_2Erealax_2Ereal^{A_27a} V0f)$

Definition 23 We define $c_2Ereal_topology_2Econtinuous$ to be $\lambda A_27a : \iota.\lambda V0f \in (ty_2Erealax_2Ereal^{A_27a} V0f)$

Definition 24 We define $c_2Ereal_topology_2Econtinuous_on$ to be $\lambda V0f \in (ty_2Erealax_2Ereal^{ty_2Erealax_2Ereal} V0f)$

Assume the following.

$$True \quad (15)$$

Assume the following.

$$\forall A_27a.nonempty A_27a \Rightarrow (\forall V0t \in 2.((\forall V1x \in A_27a.(p V0t)) \Leftrightarrow (p V0t))) \quad (16)$$

Assume the following.

$$\forall A_27a.nonempty A_27a \Rightarrow (\forall V0x \in A_27a.((V0x = V0x) \Leftrightarrow True)) \quad (17)$$

Assume the following.

$$\begin{aligned} & (\forall V0f \in (ty_2Erealax_2Ereal^{ty_2Erealax_2Ereal} V0f).(\forall V1x \in ty_2Erealax_2Ereal.(\forall V2s \in (2^{ty_2Erealax_2Ereal}).(p (ap (ap (c_2Ereal_topology_2Econtinuous ty_2Erealax_2Ereal) V0f) (ap (ap (c_2Ereal_topology_2Ewithin ty_2Erealax_2Ereal) (ap c_2Ereal_topology_2Eat V1x)) V2s))) \Leftrightarrow (\forall V3e \in ty_2Erealax_2Ereal.((p (ap (ap c_2Erealax_2Ereal_lt (ap c_2Ereal_2Ereal_of_num c_2Enum_2E0)) V3e)) \Rightarrow (\exists V4d \in ty_2Erealax_2Ereal.((p (ap (ap c_2Erealax_2Ereal_lt (ap c_2Ereal_2Ereal_of_num c_2Enum_2E0)) V4d)) \wedge (\forall V5x_27 \in ty_2Erealax_2Ereal.(((p (ap (ap (c_2Ebool_2EIN ty_2Erealax_2Ereal) V5x_27) V2s)) \wedge (p (ap (ap c_2Erealax_2Ereal_lt (ap c_2Ereal_topology_2EDist (ap (ap (c_2Epair_2E_2C ty_2Erealax_2Ereal ty_2Erealax_2Ereal) V5x_27) V1x))) V4d)))) \Rightarrow (p (ap (ap c_2Erealax_2Ereal_lt (ap c_2Ereal_topology_2EDist (ap (ap (c_2Epair_2E_2C ty_2Erealax_2Ereal ty_2Erealax_2Ereal) (ap V0f V5x_27)) (ap V0f V1x)))) V3e))))))))))))) \end{aligned} \quad (18)$$

Theorem 1

$$\begin{aligned} & (\forall V0f \in (ty_2Erealax_2Ereal^{ty_2Erealax_2Ereal}).(\forall V1s \in \\ & (2^{ty_2Erealax_2Ereal}).((p (ap (ap (c_2Ereal_topology_2Econtinuous_on \\ & V0f) V1s)) \Leftrightarrow (\forall V2x \in ty_2Erealax_2Ereal.((p (ap (ap (c_2Ebool_2EIN \\ & ty_2Erealax_2Ereal) V2x) V1s))) \Rightarrow (p (ap (ap (c_2Ereal_topology_2Econtinuous \\ & ty_2Erealax_2Ereal) V0f) (ap (ap (c_2Ereal_topology_2Ewithin \\ & ty_2Erealax_2Ereal) (ap c_2Ereal_topology_2Eat V2x)) V1s)))))))))) \end{aligned}$$