

thm\_2Ereal\_\_topology\_2ECONTINUOUS\_\_ON\_\_INV  
(TM-  
RzQMN1TNC4pPddRhs69p8ypWYGm3tEgoL)

October 26, 2020

**Definition 1** We define  $c\_2Emin\_2E\_3D$  to be  $\lambda A.\lambda x \in A.\lambda y \in A.inj\_o (x = y)$  of type  $\iota \Rightarrow \iota$ .

**Definition 2** We define  $c\_2Ebool\_2E\_2T$  to be  $(ap (ap (c\_2Emin\_2E\_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

**Definition 3** We define  $c\_2Ebool\_2E\_21$  to be  $\lambda A\_27a : \iota.(\lambda V0P \in (2^{A\_27a}).(ap (ap (c\_2Emin\_2E\_3D (2^{A\_27a}))$

**Definition 4** We define  $c\_2Ebool\_2E\_2F$  to be  $(ap (c\_2Ebool\_2E\_21 2) (\lambda V0t \in 2.V0t))$ .

**Definition 5** We define  $c\_2Ebool\_2E\_2IN$  to be  $\lambda A\_27a : \iota.(\lambda V0x \in A\_27a.(\lambda V1f \in (2^{A\_27a}).(ap V1f V0x)))$

Let  $ty\_2Erealx\_2Ereal : \iota$  be given. Assume the following.

$$nonempty\ ty\_2Erealx\_2Ereal \tag{1}$$

**Definition 6** We define  $c\_2Emin\_2E\_3D\_3D\_3E$  to be  $\lambda P \in 2.\lambda Q \in 2.inj\_o (p P \Rightarrow p Q)$  of type  $\iota$ .

**Definition 7** We define  $c\_2Ebool\_2E\_2F\_5C$  to be  $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c\_2Ebool\_2E\_21 2) (\lambda V2t \in 2.V2t)))$

Let  $ty\_2Epair\_2Eprod : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A0.nonempty\ A0 \Rightarrow \forall A1.nonempty\ A1 \Rightarrow nonempty\ (ty\_2Epair\_2Eprod\ A0\ A1) \tag{2}$$

Let  $c\_2Epair\_2EABS\_prod : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A\_27a.nonempty\ A\_27a \Rightarrow \forall A\_27b.nonempty\ A\_27b \Rightarrow c\_2Epair\_2EABS\_prod\ A\_27a\ A\_27b \in ((ty\_2Epair\_2Eprod\ A\_27a\ A\_27b)^{(2^{A\_27b})^{A\_27a}}) \tag{3}$$

**Definition 8** We define  $c\_2Epair\_2E\_2C$  to be  $\lambda A\_27a : \iota.\lambda A\_27b : \iota.\lambda V0x \in A\_27a.\lambda V1y \in A\_27b.(ap (c\_2E$

Let  $c\_2Ereal\_topology\_2EDist : \iota$  be given. Assume the following.

$$c\_2Ereal\_topology\_2EDist \in (ty\_2Erealax\_2Ereal^{(ty\_2Epair\_2Eprod\ ty\_2Erealax\_2Ereal\ ty\_2Erealax\_2Ereal)}) \quad (4)$$

Let  $ty\_2Ehreal\_2Ehreal : \iota$  be given. Assume the following.

$$nonempty\ ty\_2Ehreal\_2Ehreal \quad (5)$$

Let  $c\_2Erealax\_2Ereal\_REP\_CLASS : \iota$  be given. Assume the following.

$$c\_2Erealax\_2Ereal\_REP\_CLASS \in ((2^{(ty\_2Epair\_2Eprod\ ty\_2Ehreal\_2Ehreal\ ty\_2Ehreal\_2Ehreal)})^{ty\_2Erealax}) \quad (6)$$

**Definition 9** We define  $c\_2Emin\_2E40$  to be  $\lambda A.\lambda P \in 2^A$ . **if**  $(\exists x \in A.p\ (ap\ P\ x))$  **then** (the  $(\lambda x.x \in A \wedge p)$  of type  $\iota \Rightarrow \iota$ ).

**Definition 10** We define  $c\_2Erealax\_2Ereal\_REP$  to be  $\lambda V0a \in ty\_2Erealax\_2Ereal.(ap\ (c\_2Emin\_2E40\ ($

Let  $c\_2Erealax\_2Etrealm\_lt : \iota$  be given. Assume the following.

$$c\_2Erealax\_2Etrealm\_lt \in ((2^{(ty\_2Epair\_2Eprod\ ty\_2Ehreal\_2Ehreal\ ty\_2Ehreal\_2Ehreal)})^{(ty\_2Epair\_2Eprod\ ty\_2Ehreal)}) \quad (7)$$

**Definition 11** We define  $c\_2Erealax\_2Ereal\_lt$  to be  $\lambda V0T1 \in ty\_2Erealax\_2Ereal.\lambda V1T2 \in ty\_2Erealax\_2Ereal$

Let  $c\_2Enum\_2EZERO\_REP : \iota$  be given. Assume the following.

$$c\_2Enum\_2EZERO\_REP \in \omega \quad (8)$$

Let  $ty\_2Enum\_2Enum : \iota$  be given. Assume the following.

$$nonempty\ ty\_2Enum\_2Enum \quad (9)$$

Let  $c\_2Enum\_2EABS\_num : \iota$  be given. Assume the following.

$$c\_2Enum\_2EABS\_num \in (ty\_2Enum\_2Enum^{\omega}) \quad (10)$$

**Definition 12** We define  $c\_2Enum\_2E0$  to be  $(ap\ c\_2Enum\_2EABS\_num\ c\_2Enum\_2EZERO\_REP)$ .

Let  $c\_2Ereal\_2Ereal\_of\_num : \iota$  be given. Assume the following.

$$c\_2Ereal\_2Ereal\_of\_num \in (ty\_2Erealax\_2Ereal^{ty\_2Enum\_2Enum}) \quad (11)$$

**Definition 13** We define  $c\_2Ebool\_2E3F$  to be  $\lambda A.\lambda 27a : \iota.(\lambda V0P \in (2^{A-27a}).(ap\ V0P\ (ap\ (c\_2Emin\_2E40\ ($

**Definition 14** We define  $c\_2Ereal\_topology\_2Econtinuous\_on$  to be  $\lambda V0f \in (ty\_2Erealax\_2Ereal^{ty\_2Erealax})$

Let  $c\_2Erealax\_2Etrealm\_inv : \iota$  be given. Assume the following.

$$c\_2Erealax\_2Etrealm\_inv \in ((ty\_2Epair\_2Eprod\ ty\_2Ehreal\_2Ehreal\ ty\_2Ehreal\_2Ehreal)^{ty\_2Epair\_2Eprod\ ty\_2Ehreal\_2Ehreal\ ty\_2Ehreal\_2Ehreal}) \quad (12)$$



Assume the following.

$$True \quad (18)$$

Assume the following.

$$(\forall V0t1 \in 2.(\forall V1t2 \in 2.(((p \ V0t1) \Rightarrow (p \ V1t2)) \Rightarrow (((p \ V1t2) \Rightarrow (p \ V0t1)) \Rightarrow ((p \ V0t1) \Leftrightarrow (p \ V1t2)))))) \quad (19)$$

Assume the following.

$$(\forall V0t \in 2.(False \Rightarrow (p \ V0t))) \quad (20)$$

Assume the following.

$$\forall A\_27a.nonempty \ A\_27a \Rightarrow (\forall V0t \in 2.((\forall V1x \in A\_27a.(p \ V0t)) \Leftrightarrow (p \ V0t))) \quad (21)$$

Assume the following.

$$(\forall V0t \in 2.(((True \wedge (p \ V0t)) \Leftrightarrow (p \ V0t)) \wedge (((p \ V0t) \wedge True) \Leftrightarrow (p \ V0t)) \wedge (((False \wedge (p \ V0t)) \Leftrightarrow False) \wedge (((p \ V0t) \wedge False) \Leftrightarrow False) \wedge (((p \ V0t) \wedge (p \ V0t)) \Leftrightarrow (p \ V0t)))))) \quad (22)$$

Assume the following.

$$(\forall V0t \in 2.(((True \Rightarrow (p \ V0t)) \Leftrightarrow (p \ V0t)) \wedge (((p \ V0t) \Rightarrow True) \Leftrightarrow True) \wedge (((False \Rightarrow (p \ V0t)) \Leftrightarrow True) \wedge (((p \ V0t) \Rightarrow (p \ V0t)) \Leftrightarrow True) \wedge ((p \ V0t) \Rightarrow False) \Leftrightarrow (\neg (p \ V0t)))))) \quad (23)$$

Assume the following.

$$((\forall V0t \in 2.((\neg(\neg(p \ V0t))) \Leftrightarrow (p \ V0t)) \wedge ((\neg True) \Leftrightarrow False) \wedge ((\neg False) \Leftrightarrow True))) \quad (24)$$

Assume the following.

$$\forall A\_27a.nonempty \ A\_27a \Rightarrow (\forall V0x \in A\_27a.(\forall V1y \in A\_27a.((V0x = V1y) \Leftrightarrow (V1y = V0x)))) \quad (25)$$

Assume the following.

$$(\forall V0t \in 2.(((True \Leftrightarrow (p \ V0t)) \Leftrightarrow (p \ V0t)) \wedge (((p \ V0t) \Leftrightarrow True) \Leftrightarrow (p \ V0t)) \wedge (((False \Leftrightarrow (p \ V0t)) \Leftrightarrow (\neg (p \ V0t))) \wedge (((p \ V0t) \Leftrightarrow False) \Leftrightarrow (\neg (p \ V0t)))))) \quad (26)$$

Assume the following.

$$(\forall V0t1 \in 2.(\forall V1t2 \in 2.(\forall V2t3 \in 2.(((p \ V0t1) \Rightarrow ((p \ V1t2) \Rightarrow (p \ V2t3))) \Leftrightarrow (((p \ V0t1) \wedge (p \ V1t2)) \Rightarrow (p \ V2t3)))))) \quad (27)$$

Assume the following.

$$2.(((p \ V0x) \Leftrightarrow (p \ V1x\_27)) \wedge ((p \ V1x\_27) \Rightarrow ((p \ V2y) \Leftrightarrow (p \ V3y\_27)))) \Rightarrow \quad (28)$$

$$(((p \ V0x) \Rightarrow (p \ V2y)) \Leftrightarrow ((p \ V1x\_27) \Rightarrow (p \ V3y\_27))))$$

Assume the following.

$$(\forall V0f \in (ty\_2Erealax\_2Ereal^{ty\_2Erealax\_2Ereal}).(\forall V1s \in$$

$$(2^{ty\_2Erealax\_2Ereal}).((p \ (ap \ (ap \ c\_2Ereal\_topology\_2Econtinuous\_on$$

$$V0f) \ V1s)) \Leftrightarrow (\forall V2x \in ty\_2Erealax\_2Ereal.((p \ (ap \ (ap \ (c\_2Ebool\_2EIN$$

$$ty\_2Erealax\_2Ereal) \ V2x) \ V1s)) \Rightarrow (p \ (ap \ (ap \ (c\_2Ereal\_topology\_2Econtinuous$$

$$ty\_2Erealax\_2Ereal) \ V0f) \ (ap \ (ap \ (c\_2Ereal\_topology\_2Ewithin$$

$$ty\_2Erealax\_2Ereal) \ (ap \ c\_2Ereal\_topology\_2Eat \ V2x)) \ V1s)))))) \quad (29)$$

Assume the following.

$$(\forall V0f \in (ty\_2Erealax\_2Ereal^{ty\_2Erealax\_2Ereal}).(\forall V1s \in$$

$$(2^{ty\_2Erealax\_2Ereal}).(\forall V2a \in ty\_2Erealax\_2Ereal.(($$

$$((p \ (ap \ (ap \ (c\_2Ereal\_topology\_2Econtinuous \ ty\_2Erealax\_2Ereal)$$

$$V0f) \ (ap \ (ap \ (c\_2Ereal\_topology\_2Ewithin \ ty\_2Erealax\_2Ereal)$$

$$(ap \ c\_2Ereal\_topology\_2Eat \ V2a)) \ V1s))) \wedge (\neg((ap \ V0f \ V2a) = (ap$$

$$c\_2Ereal\_2Ereal\_of\_num \ c\_2Enum\_2E0)))) \Rightarrow (p \ (ap \ (ap \ (c\_2Ereal\_topology\_2Econtinuous$$

$$ty\_2Erealax\_2Ereal) \ (ap \ (ap \ (c\_2Ecombin\_2Eo \ ty\_2Erealax\_2Ereal$$

$$ty\_2Erealax\_2Ereal \ ty\_2Erealax\_2Ereal) \ c\_2Erealax\_2Einv) \ V0f))$$

$$(ap \ (ap \ (c\_2Ereal\_topology\_2Ewithin \ ty\_2Erealax\_2Ereal) \ (ap$$

$$c\_2Ereal\_topology\_2Eat \ V2a)) \ V1s)))))) \quad (30)$$

### Theorem 1

$$(\forall V0f \in (ty\_2Erealax\_2Ereal^{ty\_2Erealax\_2Ereal}).(\forall V1s \in$$

$$(2^{ty\_2Erealax\_2Ereal}).(((p \ (ap \ (ap \ c\_2Ereal\_topology\_2Econtinuous\_on$$

$$V0f) \ V1s)) \wedge (\forall V2x \in ty\_2Erealax\_2Ereal.((p \ (ap \ (ap \ (c\_2Ebool\_2EIN$$

$$ty\_2Erealax\_2Ereal) \ V2x) \ V1s)) \Rightarrow (\neg((ap \ V0f \ V2x) = (ap \ c\_2Ereal\_2Ereal\_of\_num$$

$$c\_2Enum\_2E0)))))) \Rightarrow (p \ (ap \ (ap \ c\_2Ereal\_topology\_2Econtinuous\_on$$

$$(ap \ (ap \ (c\_2Ecombin\_2Eo \ ty\_2Erealax\_2Ereal \ ty\_2Erealax\_2Ereal$$

$$ty\_2Erealax\_2Ereal) \ c\_2Erealax\_2Einv) \ V0f)) \ V1s))))$$