

# thm\_2Ereal\_\_topology\_2ECONTINUOUS\_\_ON\_\_SUM (TMG18moxcg1a9GGRNQeBqGbbYUVG5cvc4vM)

October 26, 2020

**Definition 1** We define  $c\_2Emin\_2E\_3D$  to be  $\lambda A.\lambda x \in A.\lambda y \in A.inj\_o (x = y)$  of type  $\iota \Rightarrow \iota$ .

**Definition 2** We define  $c\_2Ebool\_2E\_2T$  to be  $(ap (ap (c\_2Emin\_2E\_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

**Definition 3** We define  $c\_2Ebool\_2E\_21$  to be  $\lambda A\_27a : \iota.(\lambda V0P \in (2^{A\_27a}).(ap (ap (c\_2Emin\_2E\_3D (2^{A\_27a}))$

**Definition 4** We define  $c\_2Ebool\_2E\_2F$  to be  $(ap (c\_2Ebool\_2E\_21 2) (\lambda V0t \in 2.V0t))$ .

**Definition 5** We define  $c\_2Emin\_2E\_3D\_3D\_3E$  to be  $\lambda P \in 2.\lambda Q \in 2.inj\_o (p P \Rightarrow p Q)$  of type  $\iota$ .

**Definition 6** We define  $c\_2Ebool\_2E\_7E$  to be  $(\lambda V0t \in 2.(ap (ap c\_2Emin\_2E\_3D\_3D\_3E V0t) c\_2Ebool\_2E\_2F$

Let  $ty\_2Erealx\_2Ereal : \iota$  be given. Assume the following.

$$nonempty\ ty\_2Erealx\_2Ereal \tag{1}$$

**Definition 7** We define  $c\_2Ebool\_2E\_2F\_5C$  to be  $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c\_2Ebool\_2E\_21 2) (\lambda V2t \in 2.V2t))$

Let  $ty\_2Epair\_2Eprod : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A0.nonempty\ A0 \Rightarrow \forall A1.nonempty\ A1 \Rightarrow nonempty\ (ty\_2Epair\_2Eprod\ A0\ A1) \tag{2}$$

Let  $c\_2Epair\_2EABS\_prod : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A\_27a.nonempty\ A\_27a \Rightarrow \forall A\_27b.nonempty\ A\_27b \Rightarrow c\_2Epair\_2EABS\_prod\ A\_27a\ A\_27b \in ((ty\_2Epair\_2Eprod\ A\_27a\ A\_27b)^{(2^{A\_27b})^{A\_27a}}) \tag{3}$$

**Definition 8** We define  $c\_2Epair\_2E\_2C$  to be  $\lambda A\_27a : \iota.\lambda A\_27b : \iota.\lambda V0x \in A\_27a.\lambda V1y \in A\_27b.(ap (c\_2E$

Let  $c\_2Ereal\_topology\_2EDist : \iota$  be given. Assume the following.

$$c\_2Ereal\_topology\_2EDist \in (ty\_2Erealax\_2Ereal^{(ty\_2Epair\_2Eprod\ ty\_2Erealax\_2Ereal\ ty\_2Erealax\_2Ereal)}) \quad (4)$$

Let  $ty\_2Ehreal\_2Ehreal : \iota$  be given. Assume the following.

$$nonempty\ ty\_2Ehreal\_2Ehreal \quad (5)$$

Let  $c\_2Erealax\_2Ereal\_REP\_CLASS : \iota$  be given. Assume the following.

$$c\_2Erealax\_2Ereal\_REP\_CLASS \in ((2^{(ty\_2Epair\_2Eprod\ ty\_2Ehreal\_2Ehreal\ ty\_2Ehreal\_2Ehreal)})^{ty\_2Erealax}) \quad (6)$$

**Definition 9** We define  $c\_2Emin\_2E40$  to be  $\lambda A.\lambda P \in 2^A$ .if  $(\exists x \in A.p\ (ap\ P\ x))$  then (the  $(\lambda x.x \in A \wedge p$  of type  $\iota \Rightarrow \iota$ ).

**Definition 10** We define  $c\_2Erealax\_2Ereal\_REP$  to be  $\lambda V0a \in ty\_2Erealax\_2Ereal.(ap\ (c\_2Emin\_2E40\ (t$

Let  $c\_2Erealax\_2Etreall\_lt : \iota$  be given. Assume the following.

$$c\_2Erealax\_2Etreall\_lt \in ((2^{(ty\_2Epair\_2Eprod\ ty\_2Ehreal\_2Ehreal\ ty\_2Ehreal\_2Ehreal)})^{(ty\_2Epair\_2Eprod\ ty\_2Ehreal)}) \quad (7)$$

**Definition 11** We define  $c\_2Erealax\_2Ereal\_lt$  to be  $\lambda V0T1 \in ty\_2Erealax\_2Ereal.\lambda V1T2 \in ty\_2Erealax\_2Ereal$

**Definition 12** We define  $c\_2Ereal\_2Ereal\_lte$  to be  $\lambda V0x \in ty\_2Erealax\_2Ereal.\lambda V1y \in ty\_2Erealax\_2Ereal$

Let  $c\_2Enum\_2EZERO\_REP : \iota$  be given. Assume the following.

$$c\_2Enum\_2EZERO\_REP \in omega \quad (8)$$

Let  $ty\_2Enum\_2Enum : \iota$  be given. Assume the following.

$$nonempty\ ty\_2Enum\_2Enum \quad (9)$$

Let  $c\_2Enum\_2EABS\_num : \iota$  be given. Assume the following.

$$c\_2Enum\_2EABS\_num \in (ty\_2Enum\_2Enum^{omega}) \quad (10)$$

**Definition 13** We define  $c\_2Enum\_2E0$  to be  $(ap\ c\_2Enum\_2EABS\_num\ c\_2Enum\_2EZERO\_REP)$ .

Let  $c\_2Ereal\_2Ereal\_of\_num : \iota$  be given. Assume the following.

$$c\_2Ereal\_2Ereal\_of\_num \in (ty\_2Erealax\_2Ereal^{ty\_2Enum\_2Enum}) \quad (11)$$

Let  $ty\_2Ereal\_topology\_2Enet : \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A0.nonempty\ A0 \Rightarrow nonempty\ (ty\_2Ereal\_topology\_2Enet\ A0) \quad (12)$$

Let  $c\_2Ereal\_topology\_2Emk\_net : \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A.27a.nonempty\ A.27a \Rightarrow c\_2Ereal\_topology\_2Emk\_net\ A.27a \in ((ty\_2Ereal\_topology\_2Enet\ A.27a)^{(2^{A.27a})^{A.27a}}) \quad (13)$$



**Definition 27** We define  $c\_Epred\_set\_EFINITE$  to be  $\lambda A\_27a : \iota.\lambda V0s \in (2^{A\_27a}).(ap (c\_Ebool\_2E\_21 (2$

**Definition 28** We define  $c\_Eiterate\_2EITSET$  to be  $\lambda A\_27a : \iota.\lambda A\_27b : \iota.\lambda V0f \in ((A\_27a^{A\_27a})^{A\_27b}).\lambda V$

**Definition 29** We define  $c\_Eiterate\_2Eiterate$  to be  $\lambda A\_27a : \iota.\lambda A\_27b : \iota.\lambda V0op \in ((A\_27b^{A\_27b})^{A\_27b}).\lambda V$

**Definition 30** We define  $c\_Eiterate\_2ESum$  to be  $\lambda A\_27a : \iota.(ap (c\_Eiterate\_2Eiterate A\_27a ty\_2Erealax$

**Definition 31** We define  $c\_Ereal\_topology\_2Eenlimit$  to be  $\lambda A\_27a : \iota.\lambda V0net \in (ty\_2Ereal\_topology\_2E$

**Definition 32** We define  $c\_Ereal\_topology\_2Etrivial\_limit$  to be  $\lambda A\_27a : \iota.\lambda V0net \in (ty\_2Ereal\_topology$

**Definition 33** We define  $c\_Ereal\_topology\_2Eeventually$  to be  $\lambda A\_27a : \iota.\lambda V0p \in (2^{A\_27a}).\lambda V1net \in (ty\_2$

**Definition 34** We define  $c\_Ereal\_topology\_2E\_2D\_2D\_3E$  to be  $\lambda A\_27a : \iota.\lambda V0f \in (ty\_2Erealax\_2Ereal^A$

**Definition 35** We define  $c\_Ereal\_topology\_2Econtinuous$  to be  $\lambda A\_27a : \iota.\lambda V0f \in (ty\_2Erealax\_2Ereal^A$

Assume the following.

$$True \tag{19}$$

Assume the following.

$$\forall A\_27a.nonempty A\_27a \Rightarrow (\forall V0t \in 2. ((\forall V1x \in A\_27a.(p V0t) \Leftrightarrow (p V0t)))) \tag{20}$$

Assume the following.

$$\begin{aligned} & (\forall V0t \in 2. (((True \wedge (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \wedge True) \Leftrightarrow \\ & (p V0t)) \wedge (((False \wedge (p V0t)) \Leftrightarrow False) \wedge (((p V0t) \wedge False) \Leftrightarrow False) \wedge \\ & (((p V0t) \wedge (p V0t)) \Leftrightarrow (p V0t)))))) \end{aligned} \tag{21}$$

Assume the following.

$$\begin{aligned} & (\forall V0t \in 2. (((True \Rightarrow (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \Rightarrow True) \Leftrightarrow \\ & True) \wedge (((False \Rightarrow (p V0t)) \Leftrightarrow True) \wedge (((p V0t) \Rightarrow (p V0t)) \Leftrightarrow True) \wedge (( \\ & (p V0t) \Rightarrow False) \Leftrightarrow \neg (p V0t)))))) \end{aligned} \tag{22}$$

Assume the following.

$$\forall A\_27a.nonempty A\_27a \Rightarrow (\forall V0x \in A\_27a. (\forall V1y \in A\_27a. ((V0x = V1y) \Leftrightarrow (V1y = V0x)))) \tag{23}$$

Assume the following.

$$\begin{aligned} & (\forall V0t \in 2. (((True \Leftrightarrow (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \Leftrightarrow True) \Leftrightarrow \\ & (p V0t)) \wedge (((False \Leftrightarrow (p V0t)) \Leftrightarrow \neg (p V0t)) \wedge (((p V0t) \Leftrightarrow False) \Leftrightarrow \neg( \\ & p V0t)))))) \end{aligned} \tag{24}$$

Assume the following.

$$(\forall V0t1 \in 2.(\forall V1t2 \in 2.(\forall V2t3 \in 2.(((p V0t1) \Rightarrow (p V1t2) \Rightarrow (p V2t3))) \Leftrightarrow (((p V0t1) \wedge (p V1t2)) \Rightarrow (p V2t3)))))) \quad (25)$$

Assume the following.

$$(\forall V0x \in 2.(\forall V1x_{.27} \in 2.(\forall V2y \in 2.(\forall V3y_{.27} \in 2.(((p V0x) \Leftrightarrow (p V1x_{.27})) \wedge ((p V1x_{.27}) \Rightarrow ((p V2y) \Leftrightarrow (p V3y_{.27})))))) \Rightarrow ((p V0x) \Rightarrow (p V2y)) \Leftrightarrow ((p V1x_{.27}) \Rightarrow (p V3y_{.27})))))) \quad (26)$$

Assume the following.

$$(\forall V0f \in (ty\_2Erealax\_2Ereal^{ty\_2Erealax\_2Ereal}).(\forall V1s \in (2^{ty\_2Erealax\_2Ereal}).((p (ap (ap c\_2Ereal\_topology\_2Econtinuous\_on V0f) V1s)) \Leftrightarrow (\forall V2x \in ty\_2Erealax\_2Ereal.((p (ap (ap (c\_2Ebool\_2EIN ty\_2Erealax\_2Ereal) V2x) V1s)) \Rightarrow (p (ap (ap (c\_2Ereal\_topology\_2Econtinuous ty\_2Erealax\_2Ereal) V0f) (ap (ap (c\_2Ereal\_topology\_2Ewithin ty\_2Erealax\_2Ereal) (ap c\_2Ereal\_topology\_2Eat V2x)) V1s)))))))))) \quad (27)$$

Assume the following.

$$\forall A\_27a.nonempty A\_27a \Rightarrow \forall A\_27b.nonempty A\_27b \Rightarrow (\forall V0net \in (ty\_2Ereal\_topology\_2Enet A\_27a).(\forall V1f \in ((ty\_2Erealax\_2Ereal^{A\_27a})^{A\_27b}).(\forall V2s \in (2^{A\_27b}).(((p (ap (c\_2Epred\_set\_2EFINITE A\_27b) V2s)) \wedge (\forall V3a \in A\_27b.((p (ap (ap (c\_2Ebool\_2EIN A\_27b) V3a) V2s)) \Rightarrow (p (ap (ap (c\_2Ereal\_topology\_2Econtinuous A\_27a) (ap V1f V3a)) V0net)))))) \Rightarrow (p (ap (ap (c\_2Ereal\_topology\_2Econtinuous A\_27a) (\lambda V4x \in A\_27a.(ap (ap (c\_2Eiterate\_2ESum A\_27b) V2s) (\lambda V5a \in A\_27b.(ap (ap V1f V5a) V4x)))))) V0net)))))) \quad (28)$$

### Theorem 1

$$\forall A\_27a.nonempty A\_27a \Rightarrow (\forall V0t \in (2^{ty\_2Erealax\_2Ereal}).(\forall V1f \in ((ty\_2Erealax\_2Ereal^{ty\_2Erealax\_2Ereal})^{A\_27a}).(\forall V2s \in (2^{A\_27a}).(((p (ap (c\_2Epred\_set\_2EFINITE A\_27a) V2s)) \wedge (\forall V3a \in A\_27a.((p (ap (ap (c\_2Ebool\_2EIN A\_27a) V3a) V2s)) \Rightarrow (p (ap (ap c\_2Ereal\_topology\_2Econtinuous\_on (ap V1f V3a)) V0t)))))) \Rightarrow (p (ap (ap c\_2Ereal\_topology\_2Econtinuous\_on (\lambda V4x \in ty\_2Erealax\_2Ereal.(ap (ap (c\_2Eiterate\_2ESum A\_27a) V2s) (\lambda V5a \in A\_27a.(ap (ap V1f V5a) V4x)))))) V0t))))))$$