

thm_2Ereal_topology_2EHOMEOMORPHIC_BALLS (TMQLh2WGbDxJYgtgyiMjVWLD5nUULAAH7p2)

October 26, 2020

Definition 1 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define $c_2Ebool_2E_2T$ to be $(ap (ap (c_2Emin_2E_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

Definition 3 We define $c_2Ebool_2E_21$ to be $\lambda A.\lambda a : \iota.(\lambda V0P \in (2^{A \cdot 27a}).(ap (ap (c_2Emin_2E_3D (2^{A \cdot 27a}))$

Definition 4 We define $c_2Ebool_2E_2F$ to be $(ap (c_2Ebool_2E_21 2) (\lambda V0t \in 2.V0t))$.

Definition 5 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p P \Rightarrow p Q)$ of type ι .

Definition 6 We define $c_2Ebool_2E_27E$ to be $(\lambda V0t \in 2.(ap (ap c_2Emin_2E_3D_3D_3E V0t) c_2Ebool_2E_2F$

Let $ty_2Erealx_2Ereal : \iota$ be given. Assume the following.

$$nonempty\ ty_2Erealx_2Ereal \tag{1}$$

Let $ty_2Epair_2Eprod : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A0.nonempty\ A0 \Rightarrow \forall A1.nonempty\ A1 \Rightarrow nonempty\ (ty_2Epair_2Eprod\ A0\ A1) \tag{2}$$

Let $c_2Ereal_topology_2Esphere : \iota$ be given. Assume the following.

$$c_2Ereal_topology_2Esphere \in ((2^{ty_2Erealx_2Ereal})(ty_2Epair_2Eprod\ ty_2Erealx_2Ereal\ ty_2Erealx_2Ereal)) \tag{3}$$

Let $c_2Ereal_topology_2Ecball : \iota$ be given. Assume the following.

$$c_2Ereal_topology_2Ecball \in ((2^{ty_2Erealx_2Ereal})(ty_2Epair_2Eprod\ ty_2Erealx_2Ereal\ ty_2Erealx_2Ereal)) \tag{4}$$

Definition 7 We define $c_2Ebool_2E_2F_5C$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21 2) (\lambda V2t \in 2.V2t))$

Let $c_2Epair_2EABS_prod : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow \forall A_27b.nonempty\ A_27b \Rightarrow c_2Epair_2EABS_prod\ A_27a\ A_27b \in ((ty_2Epair_2Eprod\ A_27a\ A_27b)^{(2^{A_27b})^{A_27a}})$$
(5)

Definition 8 We define $c_2Epair_2E_2C$ to be $\lambda A_27a : \iota.\lambda A_27b : \iota.\lambda V0x \in A_27a.\lambda V1y \in A_27b.(ap\ (c_2E$

Let $c_2Ereal_topology_2Eball : \iota$ be given. Assume the following.

$$c_2Ereal_topology_2Eball \in ((2^{ty_2Erealax_2Ereal})(ty_2Epair_2Eprod\ ty_2Erealax_2Ereal\ ty_2Erealax_2Ereal))$$
(6)

Let $c_2Ereal_topology_2Ehomeomorphism : \iota$ be given. Assume the following.

$$c_2Ereal_topology_2Ehomeomorphism \in ((2^{(ty_2Epair_2Eprod\ (ty_2Erealax_2Ereal)^{ty_2Erealax_2Ereal})\ (ty_2Erealax_2Ereal))})$$
(7)

Definition 9 We define $c_2Emin_2E_40$ to be $\lambda A.\lambda P \in 2^A$.if $(\exists x \in A.p\ (ap\ P\ x))$ then (the $(\lambda x.x \in A \wedge p\ x)$ of type $\iota \Rightarrow \iota$).

Definition 10 We define $c_2Ebool_2E_3F$ to be $\lambda A_27a : \iota.(\lambda V0P \in (2^{A_27a}).(ap\ V0P\ (ap\ (c_2Emin_2E_40$

Definition 11 We define $c_2Ereal_topology_2Ehomeomorphic$ to be $\lambda V0s \in (2^{ty_2Erealax_2Ereal}).\lambda V1t \in (2^{ty_2Erealax_2Ereal})$

Let $c_2Enum_2EZERO_REP : \iota$ be given. Assume the following.

$$c_2Enum_2EZERO_REP \in \omega$$
(8)

Let $ty_2Enum_2Enum : \iota$ be given. Assume the following.

$$nonempty\ ty_2Enum_2Enum$$
(9)

Let $c_2Enum_2EABS_num : \iota$ be given. Assume the following.

$$c_2Enum_2EABS_num \in (ty_2Enum_2Enum^{\omega})$$
(10)

Definition 12 We define c_2Enum_2E0 to be $(ap\ c_2Enum_2EABS_num\ c_2Enum_2EZERO_REP)$.

Let $c_2Ereal_2Ereal_of_num : \iota$ be given. Assume the following.

$$c_2Ereal_2Ereal_of_num \in (ty_2Erealax_2Ereal^{ty_2Enum_2Enum})$$
(11)

Let $ty_2Ehreal_2Ehreal : \iota$ be given. Assume the following.

$$nonempty\ ty_2Ehreal_2Ehreal$$
(12)

Let $c_2Erealax_2Ereal_REP_CLASS : \iota$ be given. Assume the following.

$$c_2Erealax_2Ereal_REP_CLASS \in ((2^{(ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal\ ty_2Ehreal_2Ehreal)})^{ty_2Erealax_2Ereal})$$
(13)

Definition 13 We define $c_2Erealax_2Ereal_REP$ to be $\lambda V0a \in ty_2Erealax_2Ereal.(ap (c_2Emin_2E.40 (t$
Let $c_2Erealax_2Etrealt_lt : \iota$ be given. Assume the following.

$$c_2Erealax_2Etrealt_lt \in ((2^{(ty_2Epair_2Eprod ty_2Ehreal_2Ehreal ty_2Ehreal_2Ehreal)})(ty_2Epair_2Eprod ty_2Ehreal_2Ehreal)) \quad (14)$$

Definition 14 We define $c_2Erealax_2Ereal_lt$ to be $\lambda V0T1 \in ty_2Erealax_2Ereal.\lambda V1T2 \in ty_2Erealax_2Ereal$
Assume the following.

$$True \quad (15)$$

Assume the following.

$$\forall A_27a.nonempty A_27a \Rightarrow (\forall V0t \in 2.((\forall V1x \in A_27a.(p V0t)) \Leftrightarrow (p V0t))) \quad (16)$$

Assume the following.

$$(\forall V0t \in 2.(((True \Leftrightarrow (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \Leftrightarrow True) \Leftrightarrow (p V0t)) \wedge (((False \Leftrightarrow (p V0t)) \Leftrightarrow \neg(p V0t)) \wedge (((p V0t) \Leftrightarrow False) \Leftrightarrow \neg(p V0t)))))) \quad (17)$$

Assume the following.

$$\begin{aligned} & ((\forall V0a \in ty_2Erealax_2Ereal.(\forall V1b \in ty_2Erealax_2Ereal. \\ & (\forall V2d \in ty_2Erealax_2Ereal.(\forall V3e \in ty_2Erealax_2Ereal. \\ & (((p (ap (ap c_2Erealax_2Ereal_lt (ap c_2Ereal_2Ereal_of_num \\ c_2Enum_2E0)) V2d)) \wedge (p (ap (ap c_2Erealax_2Ereal_lt (ap c_2Ereal_2Ereal_of_num \\ c_2Enum_2E0)) V3e))) \Rightarrow (p (ap (ap c_2Ereal_topology_2Ehomeomorphic \\ (ap c_2Ereal_topology_2Eball (ap (ap (c_2Epair_2E.2C ty_2Erealax_2Ereal \\ ty_2Erealax_2Ereal) V0a) V2d))) (ap c_2Ereal_topology_2Eball \\ (ap (ap (c_2Epair_2E.2C ty_2Erealax_2Ereal ty_2Erealax_2Ereal) \\ V1b) V3e)))))) \wedge ((\forall V4a \in ty_2Erealax_2Ereal.(\forall V5b \in \\ ty_2Erealax_2Ereal.(\forall V6d \in ty_2Erealax_2Ereal.(\forall V7e \in \\ ty_2Erealax_2Ereal.(((p (ap (ap c_2Erealax_2Ereal_lt (ap c_2Ereal_2Ereal_of_num \\ c_2Enum_2E0)) V6d)) \wedge (p (ap (ap c_2Erealax_2Ereal_lt (ap c_2Ereal_2Ereal_of_num \\ c_2Enum_2E0)) V7e))) \Rightarrow (p (ap (ap c_2Ereal_topology_2Ehomeomorphic \\ (ap c_2Ereal_topology_2Eball (ap (ap (c_2Epair_2E.2C ty_2Erealax_2Ereal \\ ty_2Erealax_2Ereal) V4a) V6d))) (ap c_2Ereal_topology_2Eball \\ (ap (ap (c_2Epair_2E.2C ty_2Erealax_2Ereal ty_2Erealax_2Ereal) \\ V5b) V7e)))))) \wedge ((\forall V8a \in ty_2Erealax_2Ereal.(\forall V9b \in \\ ty_2Erealax_2Ereal.(\forall V10d \in ty_2Erealax_2Ereal.(\forall V11e \in \\ ty_2Erealax_2Ereal.(((p (ap (ap c_2Erealax_2Ereal_lt (ap c_2Ereal_2Ereal_of_num \\ c_2Enum_2E0)) V10d)) \wedge (p (ap (ap c_2Erealax_2Ereal_lt (ap c_2Ereal_2Ereal_of_num \\ c_2Enum_2E0)) V11e))) \Rightarrow (p (ap (ap c_2Ereal_topology_2Ehomeomorphic \\ (ap c_2Ereal_topology_2Esphere (ap (ap (c_2Epair_2E.2C ty_2Erealax_2Ereal \\ ty_2Erealax_2Ereal) V8a) V10d))) (ap c_2Ereal_topology_2Esphere \\ (ap (ap (c_2Epair_2E.2C ty_2Erealax_2Ereal ty_2Erealax_2Ereal) \\ V9b) V11e))))))))) \quad (18) \end{aligned}$$

Theorem 1

$$\begin{aligned} & (\forall V0a \in ty_2Erealax_2Ereal. (\forall V1b \in ty_2Erealax_2Ereal. \\ & (\forall V2d \in ty_2Erealax_2Ereal. (\forall V3e \in ty_2Erealax_2Ereal. \\ & (((p (ap (ap c_2Erealax_2Ereal_lt (ap c_2Ereal_2Ereal_of_num \\ c_2Enum_2E0)) V2d)) \wedge (p (ap (ap c_2Erealax_2Ereal_lt (ap c_2Ereal_2Ereal_of_num \\ c_2Enum_2E0)) V3e))) \Rightarrow (p (ap (ap c_2Ereal_topology_2Ehomeomorphic \\ (ap c_2Ereal_topology_2Eball (ap (ap (c_2Epair_2E_2C ty_2Erealax_2Ereal \\ ty_2Erealax_2Ereal) V0a) V2d))) (ap c_2Ereal_topology_2Eball \\ (ap (ap (c_2Epair_2E_2C ty_2Erealax_2Ereal ty_2Erealax_2Ereal) \\ V1b) V3e)))))))))) \end{aligned}$$