

thm_2Ereal__topology_2EHOMEOMORPHIC__REFL (TMS6N1xQjJpNfmw1jt9s3RnVuXFJd9eNz3P)

October 26, 2020

Definition 1 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define $c_2Ebool_2E_2T$ to be $(ap (ap (c_2Emin_2E_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

Definition 3 We define $c_2Emin_2E_40$ to be $\lambda A.\lambda P \in 2^A.if (\exists x \in A.p (ap P x))$ then (the $(\lambda x.x \in A \wedge p (ap P x))$) of type $\iota \Rightarrow \iota$.

Definition 4 We define $c_2Ebool_2E_3F$ to be $\lambda A.27a : \iota.(\lambda V0P \in (2^{A-27a}).(ap V0P (ap (c_2Emin_2E_40 A$

Let $ty_2Erealx_2Ereal : \iota$ be given. Assume the following.

$$nonempty\ ty_2Erealx_2Ereal \tag{1}$$

Definition 5 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p P \Rightarrow p Q)$ of type ι .

Definition 6 We define $c_2Ebool_2E_21$ to be $\lambda A.27a : \iota.(\lambda V0P \in (2^{A-27a}).(ap (ap (c_2Emin_2E_3D (2^{A-27a}$

Definition 7 We define $c_2Ebool_2E_2F_5C$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21 2) (\lambda V2t \in 2$

Let $ty_2Epair_2Eprod : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A0.nonempty\ A0 \Rightarrow \forall A1.nonempty\ A1 \Rightarrow nonempty\ (ty_2Epair_2Eprod\ A0\ A1) \tag{2}$$

Let $c_2Epair_2EABS_prod : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A.27a.nonempty\ A.27a \Rightarrow \forall A.27b.nonempty\ A.27b \Rightarrow c_2Epair_2EABS_prod\ A.27a\ A.27b \in ((ty_2Epair_2Eprod\ A.27a\ A.27b)^{(2^{A-27b})^{A-27a}}) \tag{3}$$

Definition 8 We define $c_2Epair_2E_2C$ to be $\lambda A.27a : \iota.\lambda A.27b : \iota.\lambda V0x \in A.27a.\lambda V1y \in A.27b.(ap (c_2E$

Let $c_2Ereal_topology_2Ehomeomorphism : \iota$ be given. Assume the following.

$$c_2Ereal_topology_2Ehomeomorphism \in ((2^{ty_2Epair_2Eprod} (ty_2Erealax_2Ereal^{ty_2Erealax_2Ereal}) (ty_2Erealax_2Ereal))) \quad (4)$$

Definition 9 We define $c_2Ereal_topology_2Ehomeomorphic$ to be $\lambda V0s \in (2^{ty_2Erealax_2Ereal}).\lambda V1t \in (2^{ty_2Erealax_2Ereal})$.

Definition 10 We define c_2Ebool_2EF to be $(ap (c_2Ebool_2E_21) 2) (\lambda V0t \in 2.V0t)$.

Definition 11 We define $c_2Ebool_2E_5C_2F$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21) 2) (\lambda V2t \in 2.V2t)))$.

Definition 12 We define $c_2Ebool_2E_7E$ to be $(\lambda V0t \in 2.(ap (ap c_2Emin_2E_3D_3D_3E V0t) c_2Ebool_2E_21))$.

Assume the following.

$$True \quad (5)$$

Assume the following.

$$(\forall V0t1 \in 2.(\forall V1t2 \in 2.(((p V0t1) \Rightarrow (p V1t2)) \Rightarrow (((p V1t2) \Rightarrow (p V0t1)) \Rightarrow ((p V0t1) \Leftrightarrow (p V1t2)))))) \quad (6)$$

Assume the following.

$$(\forall V0t \in 2.(((p V0t) \Rightarrow False) \Rightarrow (\neg(p V0t)))) \quad (7)$$

Assume the following.

$$(\forall V0t \in 2.((\neg(p V0t)) \Rightarrow ((p V0t) \Rightarrow False))) \quad (8)$$

Assume the following.

$$(\forall V0t \in 2.(((True) \Rightarrow (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \Rightarrow True) \Leftrightarrow True) \wedge (((False) \Rightarrow (p V0t)) \Leftrightarrow True) \wedge (((p V0t) \Rightarrow (p V0t)) \Leftrightarrow True) \wedge (((p V0t) \Rightarrow False) \Leftrightarrow (\neg(p V0t)))) \quad (9)$$

Assume the following.

$$((\forall V0t \in 2.((\neg(\neg(p V0t))) \Leftrightarrow (p V0t))) \wedge (((\neg True) \Leftrightarrow False) \wedge ((\neg False) \Leftrightarrow True))) \quad (10)$$

Assume the following.

$$(\forall V0t \in 2.(((True) \Leftrightarrow (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \Leftrightarrow True) \Leftrightarrow (p V0t)) \wedge (((False) \Leftrightarrow (p V0t)) \Leftrightarrow (\neg(p V0t))) \wedge (((p V0t) \Leftrightarrow False) \Leftrightarrow (\neg(p V0t)))) \quad (11)$$

Assume the following.

$$\forall A.27a.nonempty A.27a \Rightarrow (\forall V0P \in (2^{A-27a}).(\neg(\forall V1x \in A.27a.(p (ap V0P V1x)))) \Leftrightarrow (\exists V2x \in A.27a.(\neg(p (ap V0P V2x))))) \quad (12)$$

Assume the following.

$$\forall A.27a.nonempty A.27a \Rightarrow (\forall V0P \in (2^{A.27a}).((\neg(\exists V1x \in A.27a.(p (ap V0P V1x)))) \Leftrightarrow (\forall V2x \in A.27a.(\neg(p (ap V0P V2x)))))) \quad (13)$$

Assume the following.

$$\begin{aligned} & (\forall V0s \in (2^{ty_2Erealax_2Ereal}).(p (ap (ap c_2Ereal_topology_2Ehomeomorphism \\ & \quad (ap (ap (c_2Epair_2E_2C (2^{ty_2Erealax_2Ereal}) (2^{ty_2Erealax_2Ereal})) \\ & \quad V0s) V0s)) (ap (ap (c_2Epair_2E_2C (ty_2Erealax_2Ereal^{ty_2Erealax_2Ereal}) \\ & \quad (ty_2Erealax_2Ereal^{ty_2Erealax_2Ereal})) (\lambda V1x \in ty_2Erealax_2Ereal. \\ & \quad V1x)) (\lambda V2x \in ty_2Erealax_2Ereal.V2x)))))) \end{aligned} \quad (14)$$

Assume the following.

$$(\forall V0t \in 2.((\neg(\neg(p V0t))) \Leftrightarrow (p V0t))) \quad (15)$$

Assume the following.

$$(\forall V0A \in 2.((p V0A) \Rightarrow ((\neg(p V0A)) \Rightarrow False))) \quad (16)$$

Assume the following.

$$(\forall V0A \in 2.(\forall V1B \in 2.(((\neg((p V0A) \vee (p V1B))) \Rightarrow False) \Leftrightarrow ((p V0A) \Rightarrow False) \Rightarrow ((\neg(p V1B)) \Rightarrow False)))) \quad (17)$$

Assume the following.

$$(\forall V0A \in 2.(\forall V1B \in 2.(((\neg((\neg(p V0A)) \vee (p V1B))) \Rightarrow False) \Leftrightarrow ((p V0A) \Rightarrow ((\neg(p V1B)) \Rightarrow False)))) \quad (18)$$

Assume the following.

$$(\forall V0A \in 2.(((\neg(p V0A)) \Rightarrow False) \Rightarrow (((p V0A) \Rightarrow False) \Rightarrow False))) \quad (19)$$

Assume the following.

$$\begin{aligned} & (\forall V0p \in 2.(\forall V1q \in 2.(\forall V2r \in 2.(((p V0p) \Leftrightarrow (\\ & \quad (p V1q) \Leftrightarrow (p V2r))) \Leftrightarrow (((p V0p) \vee ((p V1q) \vee (p V2r))) \wedge (((p V0p) \vee ((\neg \\ & \quad p V2r)) \vee (\neg(p V1q)))) \wedge (((p V1q) \vee ((\neg(p V2r)) \vee (\neg(p V0p)))) \wedge ((p V2r) \vee \\ & \quad ((\neg(p V1q)) \vee (\neg(p V0p)))))))))) \end{aligned} \quad (20)$$

Assume the following.

$$\begin{aligned} & (\forall V0p \in 2.(\forall V1q \in 2.(\forall V2r \in 2.(((p V0p) \Leftrightarrow (\\ & \quad (p V1q) \Rightarrow (p V2r))) \Leftrightarrow (((p V0p) \vee (p V1q)) \wedge (((p V0p) \vee (\neg(p V2r))) \wedge ((\\ & \quad \neg(p V1q)) \vee ((p V2r) \vee (\neg(p V0p)))))))))) \end{aligned} \quad (21)$$

Assume the following.

$$(\forall V0p \in 2.(\forall V1q \in 2.(((p V0p) \Leftrightarrow (\neg(p V1q))) \Leftrightarrow (((p V0p) \vee (p V1q)) \wedge ((\neg(p V1q)) \vee (\neg(p V0p)))))) \quad (22)$$

Theorem 1

$$(\forall V0s \in (2^{ty_2Erealax_2Ereal}).(p (ap (ap c_2Ereal_topology_2Ehomeomorphic V0s) V0s)))$$