

thm_2Ereal__topology_2EIN__BALL__0
(TMR7UQedeGjktKRTzuJgAaJHoLTXHP86iJU)

October 26, 2020

Definition 1 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define c_2Ebool_2ET to be $(ap (ap (c_2Emin_2E_3D (2^2))) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

Let $c_2Enum_2EZERO_REP : \iota$ be given. Assume the following.

$$c_2Enum_2EZERO_REP \in \omega \tag{1}$$

Let $ty_2Enum_2Enum : \iota$ be given. Assume the following.

$$nonempty\ ty_2Enum_2Enum \tag{2}$$

Let $c_2Enum_2EABS_num : \iota$ be given. Assume the following.

$$c_2Enum_2EABS_num \in (ty_2Enum_2Enum^{\omega}) \tag{3}$$

Definition 3 We define c_2Enum_2E0 to be $(ap\ c_2Enum_2EABS_num\ c_2Enum_2EZERO_REP)$.

Let $ty_2Erealx_2Ereal : \iota$ be given. Assume the following.

$$nonempty\ ty_2Erealx_2Ereal \tag{4}$$

Let $c_2Ereal_2Ereal_of_num : \iota$ be given. Assume the following.

$$c_2Ereal_2Ereal_of_num \in (ty_2Erealx_2Ereal^{ty_2Enum_2Enum}) \tag{5}$$

Let $ty_2Ehreal_2Ehreal : \iota$ be given. Assume the following.

$$nonempty\ ty_2Ehreal_2Ehreal \tag{6}$$

Let $ty_2Epair_2Eprod : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A0.nonempty\ A0 \Rightarrow \forall A1.nonempty\ A1 \Rightarrow nonempty\ (ty_2Epair_2Eprod\ A0\ A1) \tag{7}$$

Let $c_2Erealx_2Ereal_REP_CLASS : \iota$ be given. Assume the following.

$$c_2Erealx_2Ereal_REP_CLASS \in ((2^{(ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal\ ty_2Ehreal_2Ehreal)})^{ty_2Erealx_2Ereal}) \tag{8}$$

Definition 4 We define $c_2Emin_2E_40$ to be $\lambda A.\lambda P \in 2^A.\mathbf{if} (\exists x \in A.p (ap P x)) \mathbf{then}$ (the $(\lambda x.x \in A \wedge p)$ of type $\iota \Rightarrow \iota$).

Definition 5 We define $c_2Ebool_2E_21$ to be $\lambda A.27a : \iota.(\lambda V0P \in (2^{A-27a}).(ap (ap (c_2Emin_2E_3D (2^{A-27a})))$

Definition 6 We define $c_2Erealax_2Ereal_REP$ to be $\lambda V0a \in ty_2Erealax_2Ereal.(ap (c_2Emin_2E_40 (ty$

Let $c_2Erealax_2Etrealm_neg : \iota$ be given. Assume the following.

$$c_2Erealax_2Etrealm_neg \in ((ty_2Epair_2Eprod ty_2Ehreal_2Ehreal ty_2Ehreal_2Ehreal)_{(ty_2Epair_2Eprod ty_2Ehreal_2Ehreal ty_2Ehreal_2Ehreal)}) \quad (9)$$

Let $c_2Erealax_2Etrealm_eq : \iota$ be given. Assume the following.

$$c_2Erealax_2Etrealm_eq \in ((2^{(ty_2Epair_2Eprod ty_2Ehreal_2Ehreal ty_2Ehreal_2Ehreal)})_{(ty_2Epair_2Eprod ty_2Ehreal_2Ehreal)}) \quad (10)$$

Let $c_2Erealax_2Ereal_ABS_CLASS : \iota$ be given. Assume the following.

$$c_2Erealax_2Ereal_ABS_CLASS \in (ty_2Erealax_2Ereal)^{(2^{(ty_2Epair_2Eprod ty_2Ehreal_2Ehreal ty_2Ehreal_2Ehreal)})} \quad (11)$$

Definition 7 We define $c_2Erealax_2Ereal_ABS$ to be $\lambda V0r \in (ty_2Epair_2Eprod ty_2Ehreal_2Ehreal ty$

Definition 8 We define $c_2Erealax_2Ereal_neg$ to be $\lambda V0T1 \in ty_2Erealax_2Ereal.(ap c_2Erealax_2Ereal$

Let $c_2Erealax_2Etrealm_add : \iota$ be given. Assume the following.

$$c_2Erealax_2Etrealm_add \in (((ty_2Epair_2Eprod ty_2Ehreal_2Ehreal ty_2Ehreal_2Ehreal)_{(ty_2Epair_2Eprod ty_2Ehreal_2Ehreal ty_2Ehreal_2Ehreal)})_{(ty_2Epair_2Eprod ty_2Ehreal_2Ehreal ty_2Ehreal_2Ehreal)}) \quad (12)$$

Definition 9 We define $c_2Erealax_2Ereal_add$ to be $\lambda V0T1 \in ty_2Erealax_2Ereal.\lambda V1T2 \in ty_2Erealax$

Definition 10 We define $c_2Ereal_2Ereal_sub$ to be $\lambda V0x \in ty_2Erealax_2Ereal.\lambda V1y \in ty_2Erealax_2Ereal$

Let $c_2Erealax_2Etrealm_lt : \iota$ be given. Assume the following.

$$c_2Erealax_2Etrealm_lt \in ((2^{(ty_2Epair_2Eprod ty_2Ehreal_2Ehreal ty_2Ehreal_2Ehreal)})_{(ty_2Epair_2Eprod ty_2Ehreal_2Ehreal ty_2Ehreal_2Ehreal)}) \quad (13)$$

Definition 11 We define $c_2Erealax_2Ereal_lt$ to be $\lambda V0T1 \in ty_2Erealax_2Ereal.\lambda V1T2 \in ty_2Erealax$

Definition 12 We define c_2Ebool_2E2EF to be $(ap (c_2Ebool_2E21 2) (\lambda V0t \in 2.V0t))$.

Definition 13 We define $c_2Emin_2E3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p P \Rightarrow p Q)$ of type ι .

Definition 14 We define c_2Ebool_2E7E to be $(\lambda V0t \in 2.(ap (ap c_2Emin_2E3D_3D_3E V0t) c_2Ebool_2E21))$

Definition 15 We define $c_2Ereal_2Ereal_lte$ to be $\lambda V0x \in ty_2Erealax_2Ereal.\lambda V1y \in ty_2Erealax_2Ereal$

Definition 16 We define $c_Ebool_2E_2F_5C$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_Ebool_2E_21 2) (\lambda V2t \in$

Definition 17 We define c_Ebool_2ECOND to be $\lambda A_27a : \iota.(\lambda V0t \in 2.(\lambda V1t1 \in A_27a.(\lambda V2t2 \in A_27a.($

Definition 18 We define c_Ereal_2Eabs to be $\lambda V0x \in ty_2Erealax_2Ereal.(ap (ap (ap (c_Ebool_2ECOND$

Let $c_2Ereal_topology_2EDist : \iota$ be given. Assume the following.

$$c_2Ereal_topology_2EDist \in (ty_2Erealax_2Ereal^{(ty_2Epair_2Eprod ty_2Erealax_2Ereal ty_2Erealax_2Ereal)}) \quad (14)$$

Let $c_2Epair_2EABS_prod : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A_27a.nonempty A_27a \Rightarrow \forall A_27b.nonempty A_27b \Rightarrow c_2Epair_2EABS_prod A_27a A_27b \in ((ty_2Epair_2Eprod A_27a A_27b)^{(2^{A_27b})^{A_27a}}) \quad (15)$$

Definition 19 We define $c_2Epair_2E_2C$ to be $\lambda A_27a : \iota.\lambda A_27b : \iota.\lambda V0x \in A_27a.\lambda V1y \in A_27b.(ap (c_2$

Let $c_2Ereal_topology_2Eball : \iota$ be given. Assume the following.

$$c_2Ereal_topology_2Eball \in ((2^{ty_2Erealax_2Ereal})^{(ty_2Epair_2Eprod ty_2Erealax_2Ereal ty_2Erealax_2Ereal)}) \quad (16)$$

Definition 20 We define c_Ebool_2EIN to be $\lambda A_27a : \iota.(\lambda V0x \in A_27a.(\lambda V1f \in (2^{A_27a}).(ap V1f V0x)))$

Assume the following.

$$True \quad (17)$$

Assume the following.

$$\forall A_27a.nonempty A_27a \Rightarrow (\forall V0t \in 2.((\forall V1x \in A_27a.(p V0t)) \Leftrightarrow (p V0t))) \quad (18)$$

Assume the following.

$$\forall A_27a.nonempty A_27a \Rightarrow (\forall V0x \in A_27a.((V0x = V0x) \Leftrightarrow True)) \quad (19)$$

Assume the following.

$$(\forall V0x \in ty_2Erealax_2Ereal.((ap (ap c_2Ereal_2Ereal_sub (ap c_2Ereal_2Ereal_of_num c_2Eenum_2E0) V0x) = (ap c_2Erealax_2Ereal_neg V0x)))) \quad (20)$$

Assume the following.

$$(\forall V0x \in ty_2Erealax_2Ereal.((ap c_2Ereal_2Eabs (ap c_2Erealax_2Ereal_neg V0x) = (ap c_2Ereal_2Eabs V0x)))) \quad (21)$$

Assume the following.

$$\begin{aligned}
& (\forall V0x \in ty_2Erealax_2Ereal. (\forall V1y \in ty_2Erealax_2Ereal. \\
& ((ap\ c_2Ereal_topology_2EDist\ (ap\ (ap\ (c_2Epair_2E_2C\ ty_2Erealax_2Ereal \\
& ty_2Erealax_2Ereal)\ V0x)\ V1y)) = (ap\ c_2Ereal_2Eabs\ (ap\ (ap\ c_2Ereal_2Ereal_sub \\
& V0x)\ V1y))))))
\end{aligned} \tag{22}$$

Assume the following.

$$\begin{aligned}
& (\forall V0x \in ty_2Erealax_2Ereal. (\forall V1y \in ty_2Erealax_2Ereal. \\
& (\forall V2e \in ty_2Erealax_2Ereal. ((p\ (ap\ (ap\ (c_2Ebool_2EIN\ ty_2Erealax_2Ereal) \\
& V1y)\ (ap\ c_2Ereal_topology_2Eball\ (ap\ (ap\ (c_2Epair_2E_2C\ ty_2Erealax_2Ereal \\
& ty_2Erealax_2Ereal)\ V0x)\ V2e)))) \Leftrightarrow (p\ (ap\ (ap\ c_2Erealax_2Ereal_lt \\
& (ap\ c_2Ereal_topology_2EDist\ (ap\ (ap\ (c_2Epair_2E_2C\ ty_2Erealax_2Ereal \\
& ty_2Erealax_2Ereal)\ V0x)\ V1y)))\ V2e))))))
\end{aligned} \tag{23}$$

Theorem 1

$$\begin{aligned}
& (\forall V0x \in ty_2Erealax_2Ereal. (\forall V1e \in ty_2Erealax_2Ereal. \\
& ((p\ (ap\ (ap\ (c_2Ebool_2EIN\ ty_2Erealax_2Ereal)\ V0x)\ (ap\ c_2Ereal_topology_2Eball \\
& (ap\ (ap\ (c_2Epair_2E_2C\ ty_2Erealax_2Ereal\ ty_2Erealax_2Ereal) \\
& (ap\ c_2Ereal_2Ereal_of_num\ c_2Enum_2E0))\ V1e)))) \Leftrightarrow (p\ (ap\ (ap \\
& c_2Erealax_2Ereal_lt\ (ap\ c_2Ereal_2Eabs\ V0x))\ V1e))))))
\end{aligned}$$