

thm_2Ereal__topology_2EIS__INTERVAL__POINTWISE (TMdq8VzHjjs3FsyZUDdELkhmF9kmsvKcJKW)

October 26, 2020

Definition 1 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define c_2Ebool_2EIN to be $\lambda A.\lambda a : \iota.(\lambda V0x \in A.27a.(\lambda V1f \in (2^{A-27a}).(ap V1f V0x)))$

Definition 3 We define $c_2Emin_2E_40$ to be $\lambda A.\lambda P \in 2^A.if (\exists x \in A.p (ap P x))$ then (the $(\lambda x.x \in A \wedge p$ of type $\iota \Rightarrow \iota$).

Definition 4 We define $c_2Ebool_2E_3F$ to be $\lambda A.\lambda a : \iota.(\lambda V0P \in (2^{A-27a}).(ap V0P (ap (c_2Emin_2E_40 A$

Let $ty_2Erealax_2Ereal : \iota$ be given. Assume the following.

$$nonempty\ ty_2Erealax_2Ereal \tag{1}$$

Let $ty_2Ehreal_2Ehreal : \iota$ be given. Assume the following.

$$nonempty\ ty_2Ehreal_2Ehreal \tag{2}$$

Let $ty_2Epair_2Eprod : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A0.nonempty\ A0 \Rightarrow \forall A1.nonempty\ A1 \Rightarrow nonempty\ (ty_2Epair_2Eprod\ A0\ A1) \tag{3}$$

Let $c_2Erealax_2Ereal_REP_CLASS : \iota$ be given. Assume the following.

$$c_2Erealax_2Ereal_REP_CLASS \in ((2^{(ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal\ ty_2Ehreal_2Ehreal)})\ ty_2Erealax) \tag{4}$$

Definition 5 We define c_2Ebool_2EET to be $(ap (ap (c_2Emin_2E_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

Definition 6 We define $c_2Ebool_2E_21$ to be $\lambda A.\lambda a : \iota.(\lambda V0P \in (2^{A-27a}).(ap (ap (c_2Emin_2E_3D (2^{A-27a}.$

Definition 7 We define $c_2Erealax_2Ereal_REP$ to be $\lambda V0a \in ty_2Erealax_2Ereal.(ap (c_2Emin_2E_40 (ty$

Let $c_2Erealax_2Etreall_lt : \iota$ be given. Assume the following.

$$c_2Erealax_2Etreall_lt \in ((2^{(ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal\ ty_2Ehreal_2Ehreal)})^{(ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal)}) \quad (5)$$

Definition 8 We define $c_2Erealax_2Ereal_lt$ to be $\lambda V0T1 \in ty_2Erealax_2Ereal.\lambda V1T2 \in ty_2Erealax_2Ereal$.

Definition 9 We define c_2Ebool_2EF to be $(ap\ (c_2Ebool_2E_21\ 2)\ (\lambda V0t \in 2.V0t))$.

Definition 10 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o\ (p\ P \Rightarrow p\ Q)$ of type ι .

Definition 11 We define $c_2Ebool_2E_7E$ to be $(\lambda V0t \in 2.(ap\ (ap\ c_2Emin_2E_3D_3D_3E\ V0t)\ c_2Ebool_2E_21))$.

Definition 12 We define $c_2Ereal_2Ereal_lte$ to be $\lambda V0x \in ty_2Erealax_2Ereal.\lambda V1y \in ty_2Erealax_2Ereal$.

Definition 13 We define $c_2Ebool_2E_2F_5C$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap\ (c_2Ebool_2E_21\ 2)\ (\lambda V2t \in 2.V2t))))$.

Definition 14 We define $c_2Ebool_2E_5C_2F$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap\ (c_2Ebool_2E_21\ 2)\ (\lambda V2t \in 2.V2t))))$.

Definition 15 We define $c_2Ereal_topology_2Eis_interval$ to be $\lambda V0s \in (2^{ty_2Erealax_2Ereal}).(ap\ (c_2Ebool_2E_21))$.

Assume the following.

$$True \quad (6)$$

Assume the following.

$$\begin{aligned} & (\forall V0t \in 2.(((True \Rightarrow (p\ V0t)) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \Rightarrow True) \Leftrightarrow \\ & True) \wedge (((False \Rightarrow (p\ V0t)) \Leftrightarrow True) \wedge (((p\ V0t) \Rightarrow (p\ V0t)) \Leftrightarrow True) \wedge ((\\ & (p\ V0t) \Rightarrow False) \Leftrightarrow (\neg(p\ V0t)))))) \end{aligned} \quad (7)$$

Assume the following.

$$\begin{aligned} & ((\forall V0t \in 2.((\neg(\neg(p\ V0t))) \Leftrightarrow (p\ V0t))) \wedge ((\neg True) \Leftrightarrow False) \wedge \\ & ((\neg False) \Leftrightarrow True)) \end{aligned} \quad (8)$$

Assume the following.

$$\begin{aligned} & (\forall V0t \in 2.(((True \Leftrightarrow (p\ V0t)) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \Leftrightarrow True) \Leftrightarrow \\ & (p\ V0t)) \wedge (((False \Leftrightarrow (p\ V0t)) \Leftrightarrow (\neg(p\ V0t))) \wedge (((p\ V0t) \Leftrightarrow False) \Leftrightarrow (\neg(\\ & p\ V0t)))))) \end{aligned} \quad (9)$$

Assume the following.

$$(\forall V0t \in 2.((\neg(\neg(p\ V0t))) \Leftrightarrow (p\ V0t))) \quad (10)$$

Assume the following.

$$(\forall V0A \in 2.((p\ V0A) \Rightarrow ((\neg(p\ V0A)) \Rightarrow False))) \quad (11)$$

Assume the following.

$$(\forall V0A \in 2.(\forall V1B \in 2.(((\neg((p V0A) \vee (p V1B))) \Rightarrow False) \Leftrightarrow ((p V0A) \Rightarrow False) \Rightarrow ((\neg(p V1B)) \Rightarrow False)))) \quad (12)$$

Assume the following.

$$(\forall V0A \in 2.(\forall V1B \in 2.(((\neg((\neg(p V0A)) \vee (p V1B))) \Rightarrow False) \Leftrightarrow ((p V0A) \Rightarrow ((\neg(p V1B)) \Rightarrow False)))) \quad (13)$$

Assume the following.

$$(\forall V0A \in 2.(((\neg(p V0A)) \Rightarrow False) \Rightarrow ((p V0A) \Rightarrow False) \Rightarrow False)) \quad (14)$$

Assume the following.

$$(\forall V0p \in 2.(\forall V1q \in 2.(\forall V2r \in 2.(((p V0p) \Leftrightarrow (p V1q) \Leftrightarrow (p V2r)) \Leftrightarrow (((p V0p) \vee (p V1q) \vee (p V2r)) \wedge (((p V0p) \vee ((\neg(p V2r)) \vee (\neg(p V1q)))) \wedge (((p V1q) \vee ((\neg(p V2r)) \vee (\neg(p V0p)))) \wedge ((p V2r) \vee ((\neg(p V1q)) \vee (\neg(p V0p)))))))))) \quad (15)$$

Assume the following.

$$(\forall V0p \in 2.(\forall V1q \in 2.(\forall V2r \in 2.(((p V0p) \Leftrightarrow (p V1q) \Rightarrow (p V2r)) \Leftrightarrow (((p V0p) \vee (p V1q)) \wedge (((p V0p) \vee (\neg(p V2r))) \wedge ((\neg(p V1q)) \vee ((p V2r) \vee (\neg(p V0p)))))))) \quad (16)$$

Assume the following.

$$(\forall V0p \in 2.(\forall V1q \in 2.(((\neg((p V0p) \Rightarrow (p V1q))) \Rightarrow (p V0p)))) \quad (17)$$

Assume the following.

$$(\forall V0p \in 2.(\forall V1q \in 2.(((\neg((p V0p) \Rightarrow (p V1q))) \Rightarrow (\neg(p V1q)))) \quad (18)$$

Assume the following.

$$(\forall V0p \in 2.(\forall V1q \in 2.(((\neg((p V0p) \vee (p V1q))) \Rightarrow (\neg(p V0p)))) \quad (19)$$

Assume the following.

$$(\forall V0p \in 2.(\forall V1q \in 2.(((\neg((p V0p) \vee (p V1q))) \Rightarrow (\neg(p V1q)))) \quad (20)$$

Assume the following.

$$(\forall V0p \in 2.(((\neg(\neg(p V0p))) \Rightarrow (p V0p))) \quad (21)$$

Theorem 1

$$(\forall V0s \in (2^{ty_2Erealax_2Ereal}).(\forall V1x \in ty_2Erealax_2Ereal. ((p (ap c_2Ereal_topology_2Eis_interval V0s)) \Rightarrow ((\exists V2a \in ty_2Erealax_2Ereal. ((p (ap (ap (c_2Ebool_2EIN ty_2Erealax_2Ereal) V2a) V0s)) \wedge (V2a = V1x))) \Rightarrow (p (ap (ap (c_2Ebool_2EIN ty_2Erealax_2Ereal) V1x) V0s))))))$$