

thm\_2Ereal\_topology\_2EIS\_INTERVAL\_SING  
(TMTK6cHfeoUdm6AHv7mBWtZbVNLCqivU66L)

October 26, 2020

**Definition 1** We define  $c\_2Emin\_2E\_3D$  to be  $\lambda A.\lambda x \in A.\lambda y \in A.inj\_o (x = y)$  of type  $\iota \Rightarrow \iota$ .

**Definition 2** We define  $c\_2Ebool\_2E\_2E$  to be  $(ap (ap (c\_2Emin\_2E\_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

**Definition 3** We define  $c\_2Ebool\_2E\_2E1$  to be  $\lambda A\_27a : \iota.(\lambda V0P \in (2^{A-27a}).(ap (ap (c\_2Emin\_2E\_3D (2^{A-27a})) (\lambda V1x \in 2.V1x)) (\lambda V2x \in 2.V2x))$

**Definition 4** We define  $c\_2Ebool\_2E\_2EF$  to be  $(ap (c\_2Ebool\_2E\_2E1 2) (\lambda V0t \in 2.V0t))$ .

**Definition 5** We define  $c\_2Emin\_2E\_3D\_3D\_3E$  to be  $\lambda P \in 2.\lambda Q \in 2.inj\_o (p P \Rightarrow p Q)$  of type  $\iota$ .

**Definition 6** We define  $c\_2Ebool\_2E\_2E7E$  to be  $(\lambda V0t \in 2.(ap (ap c\_2Emin\_2E\_3D\_3D\_3E V0t) c\_2Ebool\_2E\_2EF))$

**Definition 7** We define  $c\_2Epred\_set\_2EEMPTY$  to be  $\lambda A\_27a : \iota.(\lambda V0x \in A\_27a.c\_2Ebool\_2E\_2EF)$ .

**Definition 8** We define  $c\_2Ebool\_2E\_2EIN$  to be  $\lambda A\_27a : \iota.(\lambda V0x \in A\_27a.(\lambda V1f \in (2^{A-27a}).(ap V1f V0x)))$

**Definition 9** We define  $c\_2Ebool\_2E\_2E5C\_2EF$  to be  $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c\_2Ebool\_2E\_2E1 2) (\lambda V2t \in 2.V2t))$

**Definition 10** We define  $c\_2Ebool\_2E\_2E2F\_5C$  to be  $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c\_2Ebool\_2E\_2E1 2) (\lambda V2t \in 2.V2t))$

Let  $ty\_2Epair\_2Eprod : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A0.nonempty A0 \Rightarrow \forall A1.nonempty A1 \Rightarrow nonempty (ty\_2Epair\_2Eprod A0 A1) \quad (1)$$

Let  $c\_2Epair\_2EABS\_prod : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A\_27a.nonempty A\_27a \Rightarrow \forall A\_27b.nonempty A\_27b \Rightarrow c\_2Epair\_2EABS\_prod A\_27a A\_27b \in ((ty\_2Epair\_2Eprod A\_27a A\_27b)^{(2^{A-27b})^{A-27a}}) \quad (2)$$

**Definition 11** We define  $c\_2Epair\_2E\_2E2C$  to be  $\lambda A\_27a : \iota.\lambda A\_27b : \iota.\lambda V0x \in A\_27a.\lambda V1y \in A\_27b.(ap (c\_2Ebool\_2E\_2E1 2) (c\_2Epair\_2EABS\_prod A\_27a A\_27b))$

Let  $c\_2Epred\_set\_2EGSPEC : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\begin{aligned} \forall A\_27a.nonempty\ A\_27a \Rightarrow \forall A\_27b.nonempty\ A\_27b \Rightarrow c\_2Epred\_set\_2EGSPEC \\ A\_27a\ A\_27b \in ((2^{A\_27a})^{((ty\_2Epair\_2Eprod\ A\_27a\ 2)^{A\_27b})}) \end{aligned} \quad (3)$$

**Definition 12** We define  $c\_2Epred\_set\_2EINSERT$  to be  $\lambda A\_27a : \iota.\lambda V0x \in A\_27a.\lambda V1s \in (2^{A\_27a}).(ap\ (c\_2Epred\_set\_2EINSERT\ s\ x))$

Let  $ty\_2Ehreal\_2Ehreal : \iota$  be given. Assume the following.

$$nonempty\ ty\_2Ehreal\_2Ehreal \quad (4)$$

Let  $ty\_2Erealax\_2Ereal : \iota$  be given. Assume the following.

$$nonempty\ ty\_2Erealax\_2Ereal \quad (5)$$

Let  $c\_2Erealax\_2Ereal\_REP\_CLASS : \iota$  be given. Assume the following.

$$c\_2Erealax\_2Ereal\_REP\_CLASS \in ((2^{(ty\_2Epair\_2Eprod\ ty\_2Ehreal\_2Ehreal\ ty\_2Ehreal\_2Ehreal)})^{ty\_2Erealax\_2Ereal}) \quad (6)$$

**Definition 13** We define  $c\_2Emin\_2E40$  to be  $\lambda A.\lambda P \in 2^A.$ if  $(\exists x \in A.p\ (ap\ P\ x))$  then  $(the\ (\lambda x.x \in A)\ P)$  of type  $\iota \Rightarrow \iota$ .

**Definition 14** We define  $c\_2Erealax\_2Ereal\_REP$  to be  $\lambda V0a \in ty\_2Erealax\_2Ereal.(ap\ (c\_2Emin\_2E40\ a))$

Let  $c\_2Erealax\_2Etreallt : \iota$  be given. Assume the following.

$$c\_2Erealax\_2Etreallt \in ((2^{(ty\_2Epair\_2Eprod\ ty\_2Ehreal\_2Ehreal\ ty\_2Ehreal\_2Ehreal)})^{(ty\_2Epair\_2Eprod\ ty\_2Ehreal\_2Ehreal)}) \quad (7)$$

**Definition 15** We define  $c\_2Erealax\_2Ereal\_lt$  to be  $\lambda V0T1 \in ty\_2Erealax\_2Ereal.\lambda V1T2 \in ty\_2Erealax\_2Ereal.$

**Definition 16** We define  $c\_2Ereal\_2Ereal\_lte$  to be  $\lambda V0x \in ty\_2Erealax\_2Ereal.\lambda V1y \in ty\_2Erealax\_2Ereal.$

**Definition 17** We define  $c\_2Ereal\_topology\_2Eis\_interval$  to be  $\lambda V0s \in (2^{ty\_2Erealax\_2Ereal}).(ap\ (c\_2Ebool\_2Eis\_interval\ s))$

Assume the following.

$$True \quad (8)$$

Assume the following.

$$\forall A\_27a.nonempty\ A\_27a \Rightarrow (\forall V0t \in 2.((\forall V1x \in A\_27a.(p\ V0t)) \Leftrightarrow (p\ V0t))) \quad (9)$$

Assume the following.

$$\begin{aligned} (\forall V0t \in 2.(((True \vee (p\ V0t)) \Leftrightarrow True) \wedge (((p\ V0t) \vee True) \Leftrightarrow True) \wedge \\ (((False \vee (p\ V0t)) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \vee False) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \vee \\ (p\ V0t)) \Leftrightarrow (p\ V0t)))))) \end{aligned} \quad (10)$$

Assume the following.

$$\begin{aligned} & (\forall V0t \in 2.(((True \Rightarrow (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \Rightarrow True) \Leftrightarrow \\ & True) \wedge (((False \Rightarrow (p V0t)) \Leftrightarrow True) \wedge (((p V0t) \Rightarrow (p V0t)) \Leftrightarrow True) \wedge (( \\ & (p V0t) \Rightarrow False) \Leftrightarrow \neg(p V0t)))))) \end{aligned} \quad (11)$$

Assume the following.

$$\forall A\_27a.nonempty \ A\_27a \Rightarrow (\forall V0x \in A\_27a.((V0x = V0x) \Leftrightarrow True)) \quad (12)$$

Assume the following.

$$\forall A\_27a.nonempty \ A\_27a \Rightarrow (\forall V0x \in A\_27a.(\forall V1y \in A\_27a.((V0x = V1y) \Leftrightarrow (V1y = V0x)))) \quad (13)$$

Assume the following.

$$\begin{aligned} & (\forall V0t1 \in 2.(\forall V1t2 \in 2.(\forall V2t3 \in 2.(((p V0t1) \Rightarrow \\ & ((p V1t2) \Rightarrow (p V2t3))) \Leftrightarrow (((p V0t1) \wedge (p V1t2)) \Rightarrow (p V2t3)))))) \end{aligned} \quad (14)$$

Assume the following.

$$\begin{aligned} & (\forall V0x \in 2.(\forall V1x\_27 \in 2.(\forall V2y \in 2.(\forall V3y\_27 \in \\ & 2.(((p V0x) \Leftrightarrow (p V1x\_27)) \wedge ((p V1x\_27) \Rightarrow ((p V2y) \Leftrightarrow (p V3y\_27)))) \Rightarrow \\ & (((p V0x) \Rightarrow (p V2y)) \Leftrightarrow ((p V1x\_27) \Rightarrow (p V3y\_27)))))) \end{aligned} \quad (15)$$

Assume the following.

$$\begin{aligned} & (\forall V0r \in 2.(\forall V1p \in 2.(\forall V2q \in 2.(((p V1p) \wedge \\ & (p V2q)) \Rightarrow (p V0r)) \Leftrightarrow ((p V1p) \Rightarrow ((p V2q) \Rightarrow (p V0r)))))) \end{aligned} \quad (16)$$

Assume the following.

$$\begin{aligned} & \forall A\_27a.nonempty \ A\_27a \Rightarrow (\forall V0x \in A\_27a.(\forall V1y \in \\ & A\_27a.((p (ap (ap (c\_2Ebool\_2EIN A\_27a) V0x) (ap (ap (c\_2Epred\_set\_2EINSERT \\ & A\_27a) V1y) (c\_2Epred\_set\_2EEMPTY A\_27a)))) \Leftrightarrow (V0x = V1y)))) \end{aligned} \quad (17)$$

Assume the following.

$$\begin{aligned} & (\forall V0x \in ty\_2Erealx\_2Ereal.(\forall V1y \in ty\_2Erealx\_2Ereal. \\ & (((p (ap (ap c\_2Ereal\_2Ereal\_lte V0x) V1y)) \wedge (p (ap (ap c\_2Ereal\_2Ereal\_lte \\ & V1y) V0x)))) \Leftrightarrow (V0x = V1y)))) \end{aligned} \quad (18)$$

### Theorem 1

$$\begin{aligned} & (\forall V0a \in ty\_2Erealx\_2Ereal.(p (ap c\_2Ereal\_topology\_2Eis\_interval \\ & (ap (ap (c\_2Epred\_set\_2EINSERT ty\_2Erealx\_2Ereal) V0a) (c\_2Epred\_set\_2EEMPTY \\ & ty\_2Erealx\_2Ereal)))))) \end{aligned}$$