

thm_2Ereal__topology_2EIS__INTERVAL__UNIV
 (TM-
 Gyuj9v8Fh3PHGvzsaYCUFTgwY4rgW5VXo)

October 26, 2020

Definition 1 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define c_2Ebool_2ET to be $(ap (ap (c_2Emin_2E_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

Definition 3 We define $c_2Ebool_2E_21$ to be $\lambda A_27a : \iota.(\lambda V0P \in (2^{A_27a}).(ap (ap (c_2Emin_2E_3D (2^{A_27a}))$

Definition 4 We define c_2Ebool_2EF to be $(ap (c_2Ebool_2E_21 2) (\lambda V0t \in 2.V0t))$.

Definition 5 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p \Rightarrow q)$ of type ι .

Definition 6 We define $c_2Ebool_2E_7E$ to be $(\lambda V0t \in 2.(ap (ap c_2Emin_2E_3D_3D_3E V0t) c_2Ebool_2EF$

Definition 7 We define $c_2Epred_set_2EUNIV$ to be $\lambda A_27a : \iota.(\lambda V0x \in A_27a.c_2Ebool_2ET)$.

Let $ty_2Ehreal_2Ehreal : \iota$ be given. Assume the following.

$$nonempty\ ty_2Ehreal_2Ehreal \tag{1}$$

Let $ty_2Epair_2Eprod : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A0.nonempty\ A0 \Rightarrow \forall A1.nonempty\ A1 \Rightarrow nonempty\ (ty_2Epair_2Eprod\ A0\ A1) \tag{2}$$

Let $ty_2Erealax_2Ereal : \iota$ be given. Assume the following.

$$nonempty\ ty_2Erealax_2Ereal \tag{3}$$

Let $c_2Erealax_2Ereal_REP_CLASS : \iota$ be given. Assume the following.

$$c_2Erealax_2Ereal_REP_CLASS \in ((2^{(ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal\ ty_2Ehreal_2Ehreal)})\ ty_2Erealax_2Ereal) \tag{4}$$

Definition 8 We define $c_2Emin_2E.40$ to be $\lambda A.\lambda P \in 2^A.$ **if** $(\exists x \in A.p (ap P x))$ **then** $(the (\lambda x.x \in A \wedge p$
of type $\iota \Rightarrow \iota$.

Definition 9 We define $c_2Erealax_2Ereal_REP$ to be $\lambda V0a \in ty_2Erealax_2Ereal.(ap (c_2Emin_2E.40 (ty$

Let $c_2Erealax_2Etreall_lt : \iota$ be given. Assume the following.

$$c_2Erealax_2Etreall_lt \in ((2^{(ty_2Epair_2Eprod ty_2Ehreal_2Ehreal ty_2Ehreal_2Ehreal)}(ty_2Epair_2Eprod ty_2Ehreal_2Ehreal)) \quad (5)$$

Definition 10 We define $c_2Erealax_2Ereal_lt$ to be $\lambda V0T1 \in ty_2Erealax_2Ereal.\lambda V1T2 \in ty_2Erealax_2Ereal$

Definition 11 We define $c_2Ereal_2Ereal_lte$ to be $\lambda V0x \in ty_2Erealax_2Ereal.\lambda V1y \in ty_2Erealax_2Ereal$

Definition 12 We define $c_2Ebool_2E.5C.2F$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E.21 2) (\lambda V2t \in 2$

Definition 13 We define c_2Ebool_2EIN to be $\lambda A.27a : \iota.(\lambda V0x \in A.27a.(\lambda V1f \in (2^{A-27a}).(ap V1f V0x))$

Definition 14 We define $c_2Ebool_2E.2F.5C$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E.21 2) (\lambda V2t \in 2$

Definition 15 We define $c_2Ereal_topology_2Eis_interval$ to be $\lambda V0s \in (2^{ty_2Erealax_2Ereal}).(ap (c_2Ebool_2E.5C.2F$

Assume the following.

$$True \quad (6)$$

Assume the following.

$$\forall A.27a.nonempty A.27a \Rightarrow (\forall V0t \in 2.((\forall V1x \in A.27a.(p V0t)) \Leftrightarrow (p V0t))) \quad (7)$$

Assume the following.

$$\begin{aligned} & (\forall V0t \in 2.(((True \wedge (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \wedge True) \Leftrightarrow \\ & (p V0t)) \wedge (((False \wedge (p V0t)) \Leftrightarrow False) \wedge (((p V0t) \wedge False) \Leftrightarrow False) \wedge \\ & (((p V0t) \wedge (p V0t)) \Leftrightarrow (p V0t)))))) \end{aligned} \quad (8)$$

Assume the following.

$$\begin{aligned} & (\forall V0t \in 2.(((True \Rightarrow (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \Rightarrow True) \Leftrightarrow \\ & True) \wedge (((False \Rightarrow (p V0t)) \Leftrightarrow True) \wedge (((p V0t) \Rightarrow (p V0t)) \Leftrightarrow True) \wedge ((\\ & (p V0t) \Rightarrow False) \Leftrightarrow (\neg (p V0t)))))) \end{aligned} \quad (9)$$

Assume the following.

$$\begin{aligned} & (\forall V0t \in 2.(((True \Leftrightarrow (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \Leftrightarrow True) \Leftrightarrow \\ & (p V0t)) \wedge (((False \Leftrightarrow (p V0t)) \Leftrightarrow (\neg (p V0t))) \wedge (((p V0t) \Leftrightarrow False) \Leftrightarrow (\neg (\\ & p V0t)))))) \end{aligned} \quad (10)$$

Assume the following.

$$\forall A.27a.nonempty A.27a \Rightarrow (\forall V0x \in A.27a.(p (ap (ap (c_2Ebool_2EIN A.27a) V0x) (c_2Epred_set_2EUNIV A.27a)))) \quad (11)$$

Theorem 1

$$(p (ap c_2Ereal_topology_2Eis_interval (c_2Epred_set_2EUNIV ty_2Erealax_2Ereal)))$$