

thm_2Ereal_topology_2ELE_ADD
 (TMFA1Vky6M2K3hvdRS5g9LmE7dHZSwxBUpG)

October 26, 2020

Let $ty_2Enum_2Enum : \iota$ be given. Assume the following.

$$nonempty\ ty_2Enum_2Enum \quad (1)$$

Let $c_2Earithmetic_2E_2B : \iota$ be given. Assume the following.

$$c_2Earithmetic_2E_2B \in ((ty_2Enum_2Enum^{ty_2Enum_2Enum})^{ty_2Enum_2Enum}) \quad (2)$$

Definition 1 We define $c_2Emin_2E_3D$ to be $\lambda A. \lambda x \in A. \lambda y \in A. inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define c_2Ebool_2ET to be $(ap (ap (c_2Emin_2E_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

Definition 3 We define $c_2Ebool_2E_21$ to be $\lambda A_27a : \iota. (\lambda V0P \in (2^{A_27a}). (ap (ap (c_2Emin_2E_3D (2^{A_27a})) (\lambda V1P \in 2.V1P)) (\lambda V2P \in 2.V2P)))$

Definition 4 We define c_2Ebool_2EF to be $(ap (c_2Ebool_2E_21 2) (\lambda V0t \in 2.V0t))$.

Definition 5 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2. \lambda Q \in 2. inj_o (p \ P \Rightarrow p \ Q)$ of type ι .

Definition 6 We define $c_2Ebool_2E_7E$ to be $(\lambda V0t \in 2. (ap (ap c_2Emin_2E_3D_3D_3E V0t) c_2Ebool_2EF))$

Definition 7 We define $c_2Ebool_2E_2F_5C$ to be $(\lambda V0t1 \in 2. (\lambda V1t2 \in 2. (ap (c_2Ebool_2E_21 2) (\lambda V2t \in 2. inj_o (V0t1 = V1t2))))$

Let $c_2Enum_2EREP_num : \iota$ be given. Assume the following.

$$c_2Enum_2EREP_num \in (\omega^{ty_2Enum_2Enum}) \quad (3)$$

Let $c_2Enum_2ESUC_REP : \iota$ be given. Assume the following.

$$c_2Enum_2ESUC_REP \in (\omega^{\omega}) \quad (4)$$

Let $c_2Enum_2EABS_num : \iota$ be given. Assume the following.

$$c_2Enum_2EABS_num \in (ty_2Enum_2Enum^{\omega}) \quad (5)$$

Definition 8 We define c_2Enum_2ESUC to be $\lambda V0m \in ty_2Enum_2Enum.(ap\ c_2Enum_2EABS_num\ ($

Definition 9 We define $c_2Emin_2E_40$ to be $\lambda A.\lambda P \in 2^A.\text{if } (\exists x \in A.p\ (ap\ P\ x)) \text{ then } (\text{the } (\lambda x.x \in A \wedge p\ of\ type\ i \Rightarrow \iota).$

Definition 10 We define $c_2Ebool_2E_3F$ to be $\lambda A_27a : \iota.(\lambda V0P \in (2^{A_27a}).(ap\ V0P\ (ap\ (c_2Emin_2E_40\ ($

Definition 11 We define $c_2Eprim_rec_2E_3C$ to be $\lambda V0m \in ty_2Enum_2Enum.\lambda V1n \in ty_2Enum_2Enum.$

Definition 12 We define $c_2Ebool_2E_5C_2F$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap\ (c_2Ebool_2E_21\ 2)\ (\lambda V2t \in$

Definition 13 We define $c_2Earithmetic_2E_3C_3D$ to be $\lambda V0m \in ty_2Enum_2Enum.\lambda V1n \in ty_2Enum_2Enum.$

Let $c_2Enum_2EZERO_REP : \iota$ be given. Assume the following.

$$c_2Enum_2EZERO_REP \in \omega \quad (6)$$

Definition 14 We define c_2Enum_2E0 to be $(ap\ c_2Enum_2EABS_num\ c_2Enum_2EZERO_REP).$

Assume the following.

$$\begin{aligned} & (\forall V0m \in ty_2Enum_2Enum.(\forall V1n \in ty_2Enum_2Enum. \\ & p\ (ap\ (ap\ c_2Earithmetic_2E_3C_3D\ V0m)\ (ap\ (ap\ c_2Earithmetic_2E_2B \\ & V0m)\ V1n)))))) \end{aligned} \quad (7)$$

Assume the following.

$$True \quad (8)$$

Assume the following.

$$\begin{aligned} & (\forall V0t \in 2.(((True \Leftrightarrow (p\ V0t)) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \Leftrightarrow True) \Leftrightarrow \\ & (p\ V0t)) \wedge (((False \Leftrightarrow (p\ V0t)) \Leftrightarrow (\neg(p\ V0t))) \wedge (((p\ V0t) \Leftrightarrow False) \Leftrightarrow (\neg(p\ V0t))))))) \end{aligned} \quad (9)$$

Assume the following.

$$\begin{aligned} & (\forall V0P \in (2^{ty_2Enum_2Enum}).(((p\ (ap\ V0P\ c_2Enum_2E0)) \wedge \\ & (\forall V1n \in ty_2Enum_2Enum.((p\ (ap\ V0P\ V1n)) \Rightarrow (p\ (ap\ V0P\ (ap\ c_2Enum_2ESUC \\ & V1n)))))) \Rightarrow (\forall V2n \in ty_2Enum_2Enum.(p\ (ap\ V0P\ V2n)))))) \end{aligned} \quad (10)$$

Theorem 1

$$\begin{aligned} & (\forall V0m \in ty_2Enum_2Enum.(\forall V1n \in ty_2Enum_2Enum. \\ & p\ (ap\ (ap\ c_2Earithmetic_2E_3C_3D\ V0m)\ (ap\ (ap\ c_2Earithmetic_2E_2B \\ & V0m)\ V1n)))))) \end{aligned}$$