

thm_2Ereal_topology_2ELE_ADDR
(TML9REmLrfAJmpRYWZ4VECYXvkpAi6zsNBs)

October 26, 2020

Definition 1 We define $c_2Emin_2E_3D$ to be $\lambda A. \lambda x \in A. \lambda y \in A. inj_o (x = y)$ of type $\iota \rightarrow \iota$.

Let $ty_2Enum_2Enum : \iota$ be given. Assume the following.

nonempty *ty_2Enum_2Enum* (1)

Let c_2 be given. Assume the following.

$$c_2Earithmetic_2E_2B \in ((ty_2Enum_2Enum^{ty_2Enum_2Enum})^{ty_2Enum_2Enum}) \quad (2)$$

Definition 2 We define c_2Ebool_2ET to be $(ap \ (ap \ (c_2Emin_2E_3D \ (2^2)) \ (\lambda V0x \in 2.V0x)) \ (\lambda V1x \in 2.V1x))$

Definition 3 We define $c_2.Ebool_2E_21$ to be $\lambda A.27a : \iota.(\lambda V0P \in (2^A_{27a}).(ap\ (ap\ (c_2.Emin_2E_3D\ (2^A_{27a}\ P)\ V)\ 0)\ P)$

Definition 4 We define c_2Ebool_2EF to be $(ap\ (c_2Ebool_2E_21\ 2)\ (\lambda V0t \in 2.V0t))$.

Definition 5 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p\ P \Rightarrow p\ Q)$ of type ι .

Definition 6 We define $c_2Ebool_2E_7E$ to be $(\lambda V0t \in 2.(ap\ (ap\ c_2Emin_2E_3D_3D_3E\ V0t)\ c_2Ebool_2EF))$

Definition 7 We define $c_2Ebool_2E_2F_5C$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap(c_2Ebool_2E_21 2))(\lambda V2t \in 2.$

Let $c_2Enum_2EREPORT_num : \iota$ be given. Assume the following.

$$c.2Enum_2EBEP_num \in (\omega^{a_{ty_2Enum_2Enum}}) \quad (3)$$

Let $c \in \text{Enum}(\text{ESUC-BEP})$ be given. Assume the following

² Σ_{H_2} = $\Sigma_{\text{H}_2} \text{CH}_2\text{Cl}_2 / \Sigma_{\text{H}_2}\text{D}_2\text{O}$ (cm^{-2} cm^3/mol)

_____ ()

Definition 8 We define c_2Enum_2ESUC to be $\lambda V0m \in ty_2Enum_2Enum.(ap\ c_2Enum_2EABS_num\ ($

Definition 9 We define $c_2Emin_2E_40$ to be $\lambda A.\lambda P \in 2^A.\text{if } (\exists x \in A.p\ (ap\ P\ x)) \text{ then } (\text{the } (\lambda x.x \in A \wedge p\ \text{of type } \iota \Rightarrow \iota)$

Definition 10 We define $c_2Ebool_2E_3F$ to be $\lambda A_27a : \iota.(\lambda V0P \in (2^{A_27a}).(ap\ V0P\ (ap\ (c_2Emin_2E_40\ ($

Definition 11 We define $c_2Eprim_rec_2E_3C$ to be $\lambda V0m \in ty_2Enum_2Enum.\lambda V1n \in ty_2Enum_2Enum.$

Definition 12 We define $c_2Ebool_2E_5C_2F$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap\ (c_2Ebool_2E_21\ 2)\ (\lambda V2t \in$

Definition 13 We define $c_2Earithmetic_2E_3C_3D$ to be $\lambda V0m \in ty_2Enum_2Enum.\lambda V1n \in ty_2Enum_2Enum.$

Assume the following.

$$(\forall V0m \in ty_2Enum_2Enum.(\forall V1n \in ty_2Enum_2Enum.(\text{ap } (ap\ c_2Earithmetic_2E_2B\ V0m) V1n) = (\text{ap } (ap\ c_2Earithmetic_2E_2B\ V1n) V0m))) \quad (6)$$

Assume the following.

$$(\forall V0m \in ty_2Enum_2Enum.(\forall V1n \in ty_2Enum_2Enum.(p\ (ap\ (ap\ c_2Earithmetic_2E_3C_3D\ V0m)\ (ap\ (ap\ c_2Earithmetic_2E_2B\ V0m)\ V1n)))))) \quad (7)$$

Theorem 1

$$(\forall V0m \in ty_2Enum_2Enum.(\forall V1n \in ty_2Enum_2Enum.(p\ (ap\ (ap\ c_2Earithmetic_2E_3C_3D\ V1n)\ (ap\ (ap\ c_2Earithmetic_2E_2B\ V0m)\ V1n))))))$$