

thm\_2Ereal\_topoogy\_2ELE\_EXISTS  
 (TMaMdw3hmKKPFaBwXe7tnrmdXYgw97CghMr)

October 26, 2020

Let  $ty\_2Enum\_2Enum : \iota$  be given. Assume the following.

$$nonempty\ ty\_2Enum\_2Enum \quad (1)$$

Let  $c\_2Earithmetic\_2E\_2B : \iota$  be given. Assume the following.

$$c\_2Earithmetic\_2E\_2B \in ((ty\_2Enum\_2Enum^{ty\_2Enum\_2Enum})^{ty\_2Enum\_2Enum}) \quad (2)$$

**Definition 1** We define  $c\_2Emin\_2E\_3D$  to be  $\lambda A. \lambda x \in A. \lambda y \in A. inj\_o (x = y)$  of type  $\iota \Rightarrow \iota$ .

**Definition 2** We define  $c\_2Ebool\_2ET$  to be  $(ap (ap (c\_2Emin\_2E\_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

**Definition 3** We define  $c\_2Ebool\_2E\_21$  to be  $\lambda A\_27a : \iota. (\lambda V0P \in (2^{A\_27a}).(ap (ap (c\_2Emin\_2E\_3D (2^{A\_27a})) (\lambda V1P \in 2.V1P)) (\lambda V2P \in 2.V2P)))$

**Definition 4** We define  $c\_2Ebool\_2EF$  to be  $(ap (c\_2Ebool\_2E\_21 2) (\lambda V0t \in 2.V0t))$ .

**Definition 5** We define  $c\_2Emin\_2E\_3D\_3D\_3E$  to be  $\lambda P \in 2. \lambda Q \in 2. inj\_o (p \Rightarrow p \ Q)$  of type  $\iota$ .

**Definition 6** We define  $c\_2Ebool\_2E\_7E$  to be  $(\lambda V0t \in 2.(ap (ap c\_2Emin\_2E\_3D\_3D\_3E V0t) c\_2Ebool\_2EF))$

**Definition 7** We define  $c\_2Ebool\_2E\_2F\_5C$  to be  $(\lambda V0t1 \in 2. (\lambda V1t2 \in 2. (ap (c\_2Ebool\_2E\_21 2) (\lambda V2t \in 2. inj\_o (V0t1 = V2t))))$

Let  $c\_2Enum\_2EREP\_num : \iota$  be given. Assume the following.

$$c\_2Enum\_2EREP\_num \in (\omega^{ty\_2Enum\_2Enum}) \quad (3)$$

Let  $c\_2Enum\_2ESUC\_REP : \iota$  be given. Assume the following.

$$c\_2Enum\_2ESUC\_REP \in (\omega^{\omega}) \quad (4)$$

Let  $c\_2Enum\_2EABS\_num : \iota$  be given. Assume the following.

$$c\_2Enum\_2EABS\_num \in (ty\_2Enum\_2Enum^{\omega}) \quad (5)$$

**Definition 8** We define  $c\_2Enum\_2ESUC$  to be  $\lambda V0m \in ty\_2Enum\_2Enum.(ap\ c\_2Enum\_2EABS\_num\ ($

**Definition 9** We define  $c_2Emin_{\lambda A.\lambda P \in 2^A}$  to be  $\lambda A.\lambda P \in 2^A$ . if  $(\exists x \in A.p(ap P x))$  then  $(the (\lambda x.x \in A \wedge p$  of type  $\iota \Rightarrow \iota$ .

**Definition 10** We define  $c\_2Ebool\_2E\_3F$  to be  $\lambda A\_27a : \iota.(\lambda V0P \in (2^{A\_27a}).(ap\;V0P\;(ap\;(c\_2Emin\_2E\_40$

**Definition 11** We define  $c\_2Eprim\_rec\_2E\_3C$  to be  $\lambda V0m \in ty\_2Enum\_2Enum. \lambda V1n \in ty\_2Enum\_2Enum.$

**Definition 12** We define  $c\_2Ebool\_2E\_5C\_2F$  to be  $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap\ (c\_2Ebool\_2E\_21\ 2)\ (\lambda V2t \in$

**Definition 13** We define  $c\_2Earthmetic\_2E\_3C\_3D$  to be  $\lambda V0m \in ty\_2Enum\_2Enum. \lambda V1n \in ty\_2Enum\_2Enum.$

Let  $c_2Enum_2EZERO\_REP : \iota$  be given. Assume the following.

*c 2Enum 2EZERO BEP ∈ omega*

define  $c$  ?Enum ?E0 to be ( $an\ c$  ?Enum ?EABS num  $c$  ?E)

Assume the following

(VII)

Assume the following.

$$(\forall V0m \in ty\_2Enum\_2Enum. (\forall V1n \in ty\_2Enum\_2Enum. ((ap\ (ap\ c\_2Earithmetic\_2E\_2B\ V0m)\ V1n) = c\_2Enum\_2E0) \Leftrightarrow ((V0m = c\_2Enum\_2E0) \wedge (V1n = c\_2Enum\_2E0)))) \quad (8)$$

Assume the following.

$$(\forall V0m \in ty\_2Enum\_2Enum. (\forall V1n \in ty\_2Enum\_2Enum. (\forall V2p \in ty\_2Enum\_2Enum. (((ap (ap c_2Earithmetic_2E_2B V0m) V2p) = (ap (ap c_2Earithmetic_2E_2B V1n) V2p)) \Leftrightarrow (V0m = V1n)))))) \quad (9)$$

Assume the following.

$$((\forall V0n \in ty\_2Enum\_2Enum.((p (ap (ap c\_2Earithmetic\_2E\_3C\_3D V0n) c\_2Enum\_2E0)) \Leftrightarrow (V0n = c\_2Enum\_2E0))) \wedge (\forall V1m \in ty\_2Enum\_2Enum.(\forall V2n \in ty\_2Enum\_2Enum.((p (ap (ap c\_2Earithmetic\_2E\_3C\_3D V1m) (ap c\_2Enum\_2ESUC V2n))) \Leftrightarrow ((V1m = (ap c\_2Enum\_2ESUC V2n)) \vee (p (ap (ap c\_2Earithmetic\_2E\_3C\_3D V1m) V2n))))))) \quad (10)$$

Assume the following.

$$True \quad (11)$$

Assume the following.

$$(\forall V0t1 \in 2. (\forall V1t2 \in 2. (((p V0t1) \Rightarrow (p V1t2)) \Rightarrow (((p V1t2) \Rightarrow (p V0t1)) \Rightarrow ((p V0t1) \Leftrightarrow (p V1t2)))))) \quad (12)$$

Assume the following.

$$(\forall V0t \in 2. (((True \wedge (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \wedge True) \Leftrightarrow (p V0t)) \wedge (((False \wedge (p V0t)) \Leftrightarrow False) \wedge (((p V0t) \wedge False) \Leftrightarrow False) \wedge (((p V0t) \wedge (p V0t)) \Leftrightarrow (p V0t)))))) \quad (13)$$

Assume the following.

$$(\forall V0t \in 2. (((True \vee (p V0t)) \Leftrightarrow True) \wedge (((p V0t) \vee True) \Leftrightarrow True) \wedge (((False \vee (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \vee False) \Leftrightarrow (p V0t)) \wedge (((p V0t) \vee (p V0t)) \Leftrightarrow (p V0t)))))) \quad (14)$$

Assume the following.

$$(\forall V0t \in 2. (((True \Rightarrow (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \Rightarrow True) \Leftrightarrow True) \wedge (((False \Rightarrow (p V0t)) \Leftrightarrow True) \wedge (((p V0t) \Rightarrow (p V0t)) \Leftrightarrow True) \wedge (((p V0t) \Rightarrow False) \Leftrightarrow (\neg(p V0t))))))) \quad (15)$$

Assume the following.

$$\forall A\_27a.\text{nonempty } A\_27a \Rightarrow (\forall V0x \in A\_27a. ((V0x = V0x) \Leftrightarrow True)) \quad (16)$$

Assume the following.

$$\forall A\_27a.\text{nonempty } A\_27a \Rightarrow (\forall V0x \in A\_27a. (\forall V1y \in A\_27a. ((V0x = V1y) \Leftrightarrow (V1y = V0x)))) \quad (17)$$

Assume the following.

$$(\forall V0t \in 2. (((True \Leftrightarrow (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \Leftrightarrow True) \Leftrightarrow (p V0t)) \wedge (((False \Leftrightarrow (p V0t)) \Leftrightarrow (\neg(p V0t))) \wedge (((p V0t) \Leftrightarrow False) \Leftrightarrow (\neg(p V0t))))))) \quad (18)$$

Assume the following.

$$\forall A\_27a.\text{nonempty } A\_27a \Rightarrow (\forall V0P \in 2. (\forall V1Q \in (2^{A\_27a}). ((\exists V2x \in A\_27a. ((p V0P) \wedge (p (ap V1Q V2x)))) \Leftrightarrow ((p V0P) \wedge (\exists V3x \in A\_27a. (p (ap V1Q V3x))))))) \quad (19)$$

Assume the following.

$$(\forall V0t1 \in 2. (\forall V1t2 \in 2. (\forall V2t3 \in 2. (((p V0t1) \Rightarrow ((p V1t2) \Rightarrow (p V2t3))) \Leftrightarrow (((p V0t1) \wedge (p V1t2)) \Rightarrow (p V2t3)))))) \quad (20)$$

Assume the following.

$$\begin{aligned} & (\forall V0x \in 2.(\forall V1x\_27 \in 2.(\forall V2y \in 2.(\forall V3y\_27 \in \\ & 2.(((p V0x) \Leftrightarrow (p V1x\_27)) \wedge ((p V1x\_27) \Rightarrow ((p V2y) \Leftrightarrow (p V3y\_27)))))) \Rightarrow \quad (21) \\ & (((p V0x) \Rightarrow (p V2y)) \Leftrightarrow ((p V1x\_27) \Rightarrow (p V3y\_27))))))) \end{aligned}$$

Assume the following.

$$\begin{aligned} & \forall A\_27a.\text{nonempty } A\_27a \Rightarrow (\forall V0a \in A\_27a.(\exists V1x \in \\ & A\_27a.(V1x = V0a))) \quad (22) \end{aligned}$$

Assume the following.

$$\begin{aligned} & \forall A\_27a.\text{nonempty } A\_27a \Rightarrow (\forall V0P \in (2^{A\_27a}).(\forall V1Q \in \\ & 2.(((\exists V2x \in A\_27a.(p (ap V0P V2x))) \Rightarrow (p V1Q)) \Leftrightarrow (\forall V3x \in \\ & A\_27a.((p (ap V0P V3x)) \Rightarrow (p V1Q))))) \quad (23) \end{aligned}$$

Assume the following.

$$\begin{aligned} & (\forall V0P \in (2^{ty\_2Enum\_2Enum}).(((p (ap V0P c\_2Enum\_2E0)) \wedge \\ & (\forall V1n \in ty\_2Enum\_2Enum.((p (ap V0P V1n)) \Rightarrow (p (ap V0P (ap c\_2Enum\_2ESUC \\ & V1n)))))) \Rightarrow (\forall V2n \in ty\_2Enum\_2Enum.(p (ap V0P V2n)))))) \quad (24) \end{aligned}$$

Assume the following.

$$(\forall V0m \in ty\_2Enum\_2Enum.(\forall V1n \in ty\_2Enum\_2Enum.(\begin{aligned} & ((ap c\_2Enum\_2ESUC V0m) = (ap c\_2Enum\_2ESUC V1n)) \Leftrightarrow (V0m = V1n))) \quad (25) \end{aligned})$$

### Theorem 1

$$\begin{aligned} & (\forall V0m \in ty\_2Enum\_2Enum.(\forall V1n \in ty\_2Enum\_2Enum.(\begin{aligned} & (p (ap (ap c\_2Earithmetic\_2E\_3C\_3D V0m) V1n)) \Leftrightarrow (\exists V2d \in ty\_2Enum\_2Enum. \\ & (V1n = (ap (ap c\_2Earithmetic\_2E\_2B V0m) V2d)))) \end{aligned}) \end{aligned})$$