

thm_2Ereal__topology_2ELIM__AT__ID
(TMa5aSUd77hiSC7wYCTj1YNTtNgjDTeqgzM)

October 26, 2020

Definition 1 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define $c_2Ebool_2E_2ET$ to be $(ap (ap (c_2Emin_2E_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

Definition 3 We define $c_2Ebool_2E_2E21$ to be $\lambda A_27a : \iota.(\lambda V0P \in (2^{A_27a}).(ap (ap (c_2Emin_2E_3D (2^{A_27a}))$

Definition 4 We define $c_2Ebool_2E_2EF$ to be $(ap (c_2Ebool_2E_2E21 2) (\lambda V0t \in 2.V0t))$.

Definition 5 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p P \Rightarrow p Q)$ of type ι .

Definition 6 We define $c_2Ebool_2E_2E7E$ to be $(\lambda V0t \in 2.(ap (ap c_2Emin_2E_3D_3D_3E V0t) c_2Ebool_2E_2EF$

Definition 7 We define $c_2Ebool_2E_2E2F_5C$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_2E21 2) (\lambda V2t \in 2.V2t))$

Definition 8 We define $c_2Epred_set_2EUNIV$ to be $\lambda A_27a : \iota.(\lambda V0x \in A_27a.c_2Ebool_2E_2ET)$.

Let $ty_2Ereal_topology_2Eenet : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A0.nonempty A0 \Rightarrow nonempty (ty_2Ereal_topology_2Eenet A0) \quad (1)$$

Let $ty_2Erealax_2Ereal : \iota$ be given. Assume the following.

$$nonempty ty_2Erealax_2Ereal \quad (2)$$

Let $ty_2Epair_2Eprod : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A0.nonempty A0 \Rightarrow \forall A1.nonempty A1 \Rightarrow nonempty (ty_2Epair_2Eprod A0 A1) \quad (3)$$

Let $c_2Epair_2EABS_prod : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A_27a.nonempty A_27a \Rightarrow \forall A_27b.nonempty A_27b \Rightarrow c_2Epair_2EABS_prod A_27a A_27b \in ((ty_2Epair_2Eprod A_27a A_27b)^{(2^{A_27b})^{A_27a}}) \quad (4)$$

Definition 9 We define $c_2Epair_2E_2C$ to be $\lambda A.27a : \iota.\lambda A.27b : \iota.\lambda V0x \in A.27a.\lambda V1y \in A.27b.(ap (c_2Ereal_topology_2EDist : \iota$ be given. Assume the following.

$$c_2Ereal_topology_2EDist \in (ty_2Erealax_2Ereal^{(ty_2Epair_2Eprod\ ty_2Erealax_2Ereal\ ty_2Erealax_2Ereal)}) \quad (5)$$

Let $ty_2Ehreal_2Ehreal : \iota$ be given. Assume the following.

$$nonempty\ ty_2Ehreal_2Ehreal \quad (6)$$

Let $c_2Erealax_2Ereal_REP_CLASS : \iota$ be given. Assume the following.

$$c_2Erealax_2Ereal_REP_CLASS \in ((2^{(ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal\ ty_2Ehreal_2Ehreal)})^{ty_2Erealax_2Ereal}) \quad (7)$$

Definition 10 We define $c_2Emin_2E_40$ to be $\lambda A.\lambda P \in 2^A$.if $(\exists x \in A.p (ap\ P\ x))$ then (the $(\lambda x.x \in A \wedge P\ x)$ of type $\iota \Rightarrow \iota$).

Definition 11 We define $c_2Erealax_2Ereal_REP$ to be $\lambda V0a \in ty_2Erealax_2Ereal.(ap (c_2Emin_2E_40 ($

Let $c_2Erealax_2Etrealm_lt : \iota$ be given. Assume the following.

$$c_2Erealax_2Etrealm_lt \in ((2^{(ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal\ ty_2Ehreal_2Ehreal)})^{(ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal)}) \quad (8)$$

Definition 12 We define $c_2Erealax_2Ereal_lt$ to be $\lambda V0T1 \in ty_2Erealax_2Ereal.\lambda V1T2 \in ty_2Erealax_2Ereal$

Definition 13 We define $c_2Ereal_2Ereal_lte$ to be $\lambda V0x \in ty_2Erealax_2Ereal.\lambda V1y \in ty_2Erealax_2Ereal$

Let $c_2Enum_2EZERO_REP : \iota$ be given. Assume the following.

$$c_2Enum_2EZERO_REP \in \omega \quad (9)$$

Let $ty_2Enum_2Enum : \iota$ be given. Assume the following.

$$nonempty\ ty_2Enum_2Enum \quad (10)$$

Let $c_2Enum_2EABS_num : \iota$ be given. Assume the following.

$$c_2Enum_2EABS_num \in (ty_2Enum_2Enum^{\omega}) \quad (11)$$

Definition 14 We define c_2Enum_2E0 to be $(ap\ c_2Enum_2EABS_num\ c_2Enum_2EZERO_REP)$.

Let $c_2Ereal_2Ereal_of_num : \iota$ be given. Assume the following.

$$c_2Ereal_2Ereal_of_num \in (ty_2Erealax_2Ereal^{ty_2Enum_2Enum}) \quad (12)$$

Let $c_2Ereal_topology_2Emk_net : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A.27a.nonempty\ A.27a \Rightarrow c_2Ereal_topology_2Emk_net\ A.27a \in ((ty_2Ereal_topology_2Emk_net\ A.27a)^{(2^{A.27a})^{A.27a}}) \quad (13)$$

Definition 15 We define $c_2Ereal_topology_2Eat$ to be $\lambda V0a \in ty_2Erealax_2Ereal.(ap (c_2Ereal_topology_2Eat$

Definition 16 We define c_2Ebool_2EIN to be $\lambda A_27a : \iota.(\lambda V0x \in A_27a.(\lambda V1f \in (2^{A_27a}).(ap V1f V0x)))$

Let $c_2Ereal_topology_2Eenetord : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A_27a.nonempty A_27a \Rightarrow c_2Ereal_topology_2Eenetord A_27a \in ((2^{A_27a})^{A_27a})^{(ty_2Ereal_topology_2Eenetord A_27a)} \quad (14)$$

Definition 17 We define $c_2Ereal_topology_2Ewithin$ to be $\lambda A_27a : \iota.\lambda V0net \in (ty_2Ereal_topology_2Ewithin$

Definition 18 We define c_2Ebool_2E3F to be $\lambda A_27a : \iota.(\lambda V0P \in (2^{A_27a}).(ap V0P (ap (c_2Emin_2E40$

Definition 19 We define $c_2Ebool_2E5C_2F$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E21 2) (\lambda V2t \in$

Definition 20 We define $c_2Ereal_topology_2Etrivial_limit$ to be $\lambda A_27a : \iota.\lambda V0net \in (ty_2Ereal_topology_2Etrivial_limit$

Definition 21 We define $c_2Ereal_topology_2Eeventually$ to be $\lambda A_27a : \iota.\lambda V0p \in (2^{A_27a}).\lambda V1net \in (ty_2Ereal_topology_2Eeventually$

Definition 22 We define $c_2Ereal_topology_2E2D_2D_3E$ to be $\lambda A_27a : \iota.\lambda V0f \in (ty_2Erealax_2Ereal^{A_27a})$

Assume the following.

$$True \quad (15)$$

Assume the following.

$$\forall A_27a.nonempty A_27a \Rightarrow (\forall V0t \in 2.((\forall V1x \in A_27a.(p V0t)) \Leftrightarrow (p V0t))) \quad (16)$$

Assume the following.

$$\forall A_27a.nonempty A_27a \Rightarrow (\forall V0x \in A_27a.(\forall V1y \in A_27a.((V0x = V1y) \Leftrightarrow (V1y = V0x)))) \quad (17)$$

Assume the following.

$$(\forall V0t \in 2.(((True \Leftrightarrow (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \Leftrightarrow True) \Leftrightarrow (p V0t)) \wedge (((False \Leftrightarrow (p V0t)) \Leftrightarrow (\neg (p V0t))) \wedge (((p V0t) \Leftrightarrow False) \Leftrightarrow (\neg (p V0t))))) \quad (18)$$

Assume the following.

$$(\forall V0x \in ty_2Erealax_2Ereal.((ap (ap (c_2Ereal_topology_2Ewithin ty_2Erealax_2Ereal) (ap c_2Ereal_topology_2Eat V0x)) (c_2Epred_set_2EUNIV ty_2Erealax_2Ereal)) = (ap c_2Ereal_topology_2Eat V0x))) \quad (19)$$

Assume the following.

$$(\forall V0a \in ty_2Erealax_2Ereal.(\forall V1s \in (2^{ty_2Erealax_2Ereal}).(p (ap (ap (ap (c_2Ereal_topology_2E2D_2D_3E ty_2Erealax_2Ereal) (\lambda V2x \in ty_2Erealax_2Ereal.V2x)) V0a) (ap (ap (c_2Ereal_topology_2Ewithin ty_2Erealax_2Ereal) (ap c_2Ereal_topology_2Eat V0a)) V1s)))))) \quad (20)$$

Theorem 1

$(\forall V0a \in ty_2Erealax_2Ereal.(p (ap (ap (ap (c_2Ereal_topology_2E_2D_2D_3E$
 $ty_2Erealax_2Ereal) (\lambda V1x \in ty_2Erealax_2Ereal.V1x)) V0a)$
 $(ap c_2Ereal_topology_2Eat V0a))))$