

# thm\_2Ereal\_\_topology\_2ELIM\_\_WITHIN\_\_INTERIOR (TMXzu8nhFff7c1d3excGyK8H7uo1Q8oSh22)

October 26, 2020

**Definition 1** We define  $c\_2Emin\_2E\_3D$  to be  $\lambda A.\lambda x \in A.\lambda y \in A.inj\_o (x = y)$  of type  $\iota \Rightarrow \iota$ .

**Definition 2** We define  $c\_2Ebool\_2ET$  to be  $(ap (ap (c\_2Emin\_2E\_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

**Definition 3** We define  $c\_2Ebool\_2E\_21$  to be  $\lambda A\_27a : \iota.(\lambda V0P \in (2^{A\_27a}).(ap (ap (c\_2Emin\_2E\_3D (2^{A\_27a}))$

**Definition 4** We define  $c\_2Ebool\_2EF$  to be  $(ap (c\_2Ebool\_2E\_21 2) (\lambda V0t \in 2.V0t))$ .

**Definition 5** We define  $c\_2Emin\_2E\_3D\_3D\_3E$  to be  $\lambda P \in 2.\lambda Q \in 2.inj\_o (p P \Rightarrow p Q)$  of type  $\iota$ .

**Definition 6** We define  $c\_2Ebool\_2E\_7E$  to be  $(\lambda V0t \in 2.(ap (ap c\_2Emin\_2E\_3D\_3D\_3E V0t) c\_2Ebool\_2EF$

**Definition 7** We define  $c\_2Ebool\_2E\_2F\_5C$  to be  $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c\_2Ebool\_2E\_21 2) (\lambda V2t \in 2.V2t$

Let  $ty\_2Epair\_2Eprod : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A0.nonempty A0 \Rightarrow \forall A1.nonempty A1 \Rightarrow nonempty (ty\_2Epair\_2Eprod A0 A1) \quad (1)$$

Let  $c\_2Epair\_2EABS\_prod : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A\_27a.nonempty A\_27a \Rightarrow \forall A\_27b.nonempty A\_27b \Rightarrow c\_2Epair\_2EABS\_prod A\_27a A\_27b \in ((ty\_2Epair\_2Eprod A\_27a A\_27b)^{(2^{A\_27b})^{A\_27a}}) \quad (2)$$

**Definition 8** We define  $c\_2Epair\_2E\_2C$  to be  $\lambda A\_27a : \iota.\lambda A\_27b : \iota.\lambda V0x \in A\_27a.\lambda V1y \in A\_27b.(ap (c\_2E$

Let  $ty\_2Erealax\_2Ereal : \iota$  be given. Assume the following.

$$nonempty ty\_2Erealax\_2Ereal \quad (3)$$

Let  $c\_2Ereal\_topology\_2EDist : \iota$  be given. Assume the following.

$$c\_2Ereal\_topology\_2EDist \in (ty\_2Erealax\_2Ereal^{(ty\_2Epair\_2Eprod\ ty\_2Erealax\_2Ereal\ ty\_2Erealax\_2Ereal)}) \quad (4)$$

Let  $c\_2Enum\_2EZERO\_REP : \iota$  be given. Assume the following.

$$c\_2Enum\_2EZERO\_REP \in \omega \quad (5)$$

Let  $ty\_2Enum\_2Enum : \iota$  be given. Assume the following.

$$nonempty\ ty\_2Enum\_2Enum \quad (6)$$

Let  $c\_2Enum\_2EABS\_num : \iota$  be given. Assume the following.

$$c\_2Enum\_2EABS\_num \in (ty\_2Enum\_2Enum^{\omega}) \quad (7)$$

**Definition 9** We define  $c\_2Enum\_2E0$  to be  $(ap\ c\_2Enum\_2EABS\_num\ c\_2Enum\_2EZERO\_REP)$ .

Let  $c\_2Ereal\_2Ereal\_of\_num : \iota$  be given. Assume the following.

$$c\_2Ereal\_2Ereal\_of\_num \in (ty\_2Erealax\_2Ereal^{ty\_2Enum\_2Enum}) \quad (8)$$

Let  $ty\_2Ehreal\_2Ehreal : \iota$  be given. Assume the following.

$$nonempty\ ty\_2Ehreal\_2Ehreal \quad (9)$$

Let  $c\_2Erealax\_2Ereal\_REP\_CLASS : \iota$  be given. Assume the following.

$$c\_2Erealax\_2Ereal\_REP\_CLASS \in ((2^{(ty\_2Epair\_2Eprod\ ty\_2Ehreal\_2Ehreal\ ty\_2Ehreal\_2Ehreal)})^{ty\_2Erealax}) \quad (10)$$

**Definition 10** We define  $c\_2Emin\_2E\_40$  to be  $\lambda A.\lambda P \in 2^A.\text{if } (\exists x \in A.p\ (ap\ P\ x)) \text{ then } (the\ (\lambda x.x \in A \wedge P\ x)) \text{ of type } \iota \Rightarrow \iota$ .

**Definition 11** We define  $c\_2Erealax\_2Ereal\_REP$  to be  $\lambda V0a \in ty\_2Erealax\_2Ereal.(ap\ (c\_2Emin\_2E\_40\ (the\ (\lambda x.x \in A \wedge P\ x))\ of\ type\ \iota \Rightarrow \iota)))$

Let  $c\_2Erealax\_2Etrealt\_lt : \iota$  be given. Assume the following.

$$c\_2Erealax\_2Etrealt\_lt \in ((2^{(ty\_2Epair\_2Eprod\ ty\_2Ehreal\_2Ehreal\ ty\_2Ehreal\_2Ehreal)})^{(ty\_2Epair\_2Eprod\ ty\_2Ehreal\_2Ehreal)}) \quad (11)$$

**Definition 12** We define  $c\_2Erealax\_2Ereal\_lt$  to be  $\lambda V0T1 \in ty\_2Erealax\_2Ereal.\lambda V1T2 \in ty\_2Erealax\_2Ereal.(ap\ (c\_2Emin\_2E\_40\ (the\ (\lambda x.x \in A \wedge P\ x))\ of\ type\ \iota \Rightarrow \iota)))$

Let  $ty\_2Ereal\_topology\_2Enet : \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A0.nonempty\ A0 \Rightarrow nonempty\ (ty\_2Ereal\_topology\_2Enet\ A0) \quad (12)$$

Let  $c\_2Ereal\_topology\_2Enetord : \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A\_27a.nonempty\ A\_27a \Rightarrow c\_2Ereal\_topology\_2Enetord\ A\_27a \in (((2^{A\_27a})^{A\_27a})^{(ty\_2Ereal\_topology\_2Enet\ A\_27a)}) \quad (13)$$

**Definition 13** We define  $c\_Ebool\_2E\_3F$  to be  $\lambda A\_27a : \iota. (\lambda V0P \in (2^{A-27a}). (ap\ V0P\ (ap\ (c\_2Emin\_2E\_40$

**Definition 14** We define  $c\_Ebool\_2E\_5C\_2F$  to be  $(\lambda V0t1 \in 2. (\lambda V1t2 \in 2. (ap\ (c\_2Ebool\_2E\_21\ 2)\ (\lambda V2t \in$

**Definition 15** We define  $c\_2Ereal\_topology\_2Etrivial\_limit$  to be  $\lambda A\_27a : \iota. \lambda V0net \in (ty\_2Ereal\_topology$

**Definition 16** We define  $c\_2Ereal\_topology\_2Eeventually$  to be  $\lambda A\_27a : \iota. \lambda V0p \in (2^{A-27a}). \lambda V1net \in (ty\_2$

**Definition 17** We define  $c\_2Ereal\_topology\_2E\_2D\_2D\_3E$  to be  $\lambda A\_27a : \iota. \lambda V0f \in (ty\_2Erealax\_2Ereal^A$

**Definition 18** We define  $c\_2Ereal\_2Ereal\_lte$  to be  $\lambda V0x \in ty\_2Erealax\_2Ereal. \lambda V1y \in ty\_2Erealax\_2Er$

Let  $c\_2Ereal\_topology\_2Emk\_net : \iota \Rightarrow \iota$  be given. Assume the following.

$$\begin{aligned} & \forall A\_27a. nonempty\ A\_27a \Rightarrow c\_2Ereal\_topology\_2Emk\_net \\ & A\_27a \in ((ty\_2Ereal\_topology\_2Enet\ A\_27a)^{(2^{A-27a})^{A-27a}}) \end{aligned} \quad (14)$$

**Definition 19** We define  $c\_2Ereal\_topology\_2Eat$  to be  $\lambda V0a \in ty\_2Erealax\_2Ereal. (ap\ (c\_2Ereal\_topolo$

**Definition 20** We define  $c\_Ebool\_2EIN$  to be  $\lambda A\_27a : \iota. (\lambda V0x \in A\_27a. (\lambda V1f \in (2^{A-27a}). (ap\ V1f\ V0x))$

**Definition 21** We define  $c\_2Ereal\_topology\_2Ewithin$  to be  $\lambda A\_27a : \iota. \lambda V0net \in (ty\_2Ereal\_topology\_2Em$

**Definition 22** We define  $c\_2Epred\_set\_2ESUBSET$  to be  $\lambda A\_27a : \iota. \lambda V0s \in (2^{A-27a}). \lambda V1t \in (2^{A-27a}). (ap\ ($

**Definition 23** We define  $c\_2Ereal\_topology\_2EOpen$  to be  $\lambda V0s \in (2^{ty\_2Erealax\_2Ereal}). (ap\ (c\_2Ebool\_2E2$

Let  $c\_2Epred\_set\_2EGSPEC : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\begin{aligned} & \forall A\_27a. nonempty\ A\_27a \Rightarrow \forall A\_27b. nonempty\ A\_27b \Rightarrow c\_2Epred\_set\_2EGSPEC \\ & A\_27a\ A\_27b \in ((2^{A-27a})^{((ty\_2Epair\_2Eprod\ A\_27a\ 2)^{A-27b})}) \end{aligned} \quad (15)$$

**Definition 24** We define  $c\_2Ereal\_topology\_2Einterior$  to be  $\lambda V0s \in (2^{ty\_2Erealax\_2Ereal}). (ap\ (c\_2Epred\_s$

Assume the following.

$$True \quad (16)$$

Assume the following.

$$\begin{aligned} & \forall A\_27a. nonempty\ A\_27a \Rightarrow (\forall V0t \in 2. ((\forall V1x \in \\ & A\_27a. (p\ V0t)) \Leftrightarrow (p\ V0t))) \end{aligned} \quad (17)$$

Assume the following.

$$\begin{aligned} & (\forall V0t \in 2. (((True \Rightarrow (p\ V0t)) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \Rightarrow True) \Leftrightarrow \\ & True) \wedge (((False \Rightarrow (p\ V0t)) \Leftrightarrow True) \wedge (((p\ V0t) \Rightarrow (p\ V0t)) \Leftrightarrow True) \wedge (( \\ & (p\ V0t) \Rightarrow False) \Leftrightarrow (\neg (p\ V0t)))))) \end{aligned} \quad (18)$$

Assume the following.

$$\begin{aligned} & \forall A\_27a. nonempty\ A\_27a \Rightarrow (\forall V0x \in A\_27a. ((V0x = V0x) \Leftrightarrow \\ & True)) \end{aligned} \quad (19)$$

Assume the following.

$$\forall A.27a.nonempty A.27a \Rightarrow (\forall V0x \in A.27a. (\forall V1y \in A.27a. ((V0x = V1y) \Leftrightarrow (V1y = V0x)))) \quad (20)$$

Assume the following.

$$\begin{aligned} & (\forall V0t \in 2. (((True \Leftrightarrow (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \Leftrightarrow True) \Leftrightarrow \\ & (p V0t)) \wedge (((False \Leftrightarrow (p V0t)) \Leftrightarrow \neg(p V0t)) \wedge (((p V0t) \Leftrightarrow False) \Leftrightarrow \neg( \\ & p V0t)))))) \end{aligned} \quad (21)$$

Assume the following.

$$\begin{aligned} & (\forall V0t1 \in 2. (\forall V1t2 \in 2. (\forall V2t3 \in 2. (((p V0t1) \Rightarrow \\ & ((p V1t2) \Rightarrow (p V2t3))) \Leftrightarrow (((p V0t1) \wedge (p V1t2)) \Rightarrow (p V2t3)))))) \end{aligned} \quad (22)$$

Assume the following.

$$\begin{aligned} & (\forall V0x \in 2. (\forall V1x.27 \in 2. (\forall V2y \in 2. (\forall V3y.27 \in \\ & 2. (((((p V0x) \Leftrightarrow (p V1x.27)) \wedge ((p V1x.27) \Rightarrow ((p V2y) \Leftrightarrow (p V3y.27)))) \Rightarrow \\ & (((p V0x) \Rightarrow (p V2y)) \Leftrightarrow ((p V1x.27) \Rightarrow (p V3y.27))))))))) \end{aligned} \quad (23)$$

Assume the following.

$$\begin{aligned} & (\forall V0p \in (2^{ty.2Erealax.2Ereal}). (\forall V1s \in (2^{ty.2Erealax.2Ereal}). \\ & (\forall V2x \in ty.2Erealax.2Ereal. ((p (ap (ap (c.2Ebool.2EIN ty.2Erealax.2Ereal) \\ & V2x) (ap c.2Ereal\_topology\_2Einterior V1s))) \Rightarrow ((p (ap (ap (c.2Ereal\_topology\_2Eeventually \\ & ty.2Erealax.2Ereal) V0p) (ap (ap (c.2Ereal\_topology\_2Ewithin \\ & ty.2Erealax.2Ereal) (ap c.2Ereal\_topology\_2Eat V2x)) V1s))) \Leftrightarrow \\ & (p (ap (ap (c.2Ereal\_topology\_2Eeventually ty.2Erealax.2Ereal) \\ & V0p) (ap c.2Ereal\_topology\_2Eat V2x))))))))) \end{aligned} \quad (24)$$

### Theorem 1

$$\begin{aligned} & (\forall V0f \in (ty.2Erealax.2Ereal^{ty.2Erealax.2Ereal}). (\forall V1l \in \\ & ty.2Erealax.2Ereal. (\forall V2s \in (2^{ty.2Erealax.2Ereal}). ( \\ & \forall V3x \in ty.2Erealax.2Ereal. ((p (ap (ap (c.2Ebool.2EIN ty.2Erealax.2Ereal) \\ & V3x) (ap c.2Ereal\_topology\_2Einterior V2s))) \Rightarrow ((p (ap (ap (ap \\ & (c.2Ereal\_topology\_2E\_2D\_2D\_3E ty.2Erealax.2Ereal) V0f) V1l) \\ & (ap (ap (c.2Ereal\_topology\_2Ewithin ty.2Erealax.2Ereal) (ap \\ & c.2Ereal\_topology\_2Eat V3x)) V2s))) \Leftrightarrow (p (ap (ap (ap (c.2Ereal\_topology\_2E\_2D\_2D\_3E \\ & ty.2Erealax.2Ereal) V0f) V1l) (ap c.2Ereal\_topology\_2Eat V3x))))))))) \end{aligned}$$