

# thm\_2Ereal\_\_topology\_2ELINEAR\_\_SUB (TMMxcVqdqnPA4H5iwt54shd13fh5eEz5q4R)

October 26, 2020

**Definition 1** We define  $c\_2Emin\_2E\_3D$  to be  $\lambda A.\lambda x \in A.\lambda y \in A.inj\_o (x = y)$  of type  $\iota \Rightarrow \iota$ .

**Definition 2** We define  $c\_2Ebool\_2E\_7E$  to be  $(ap (ap (c\_2Emin\_2E\_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

**Definition 3** We define  $c\_2Ebool\_2E\_21$  to be  $\lambda A.\lambda a : \iota.(\lambda V0P \in (2^{A-27a}).(ap (ap (c\_2Emin\_2E\_3D (2^{A-27a}))$

**Definition 4** We define  $c\_2Ebool\_2E\_7E$  to be  $(ap (c\_2Ebool\_2E\_21 2) (\lambda V0t \in 2.V0t))$ .

**Definition 5** We define  $c\_2Emin\_2E\_3D\_3D\_3E$  to be  $\lambda P \in 2.\lambda Q \in 2.inj\_o (p P \Rightarrow p Q)$  of type  $\iota$ .

**Definition 6** We define  $c\_2Ebool\_2E\_7E$  to be  $(\lambda V0t \in 2.(ap (ap c\_2Emin\_2E\_3D\_3D\_3E V0t) c\_2Ebool\_2E\_7E$

**Definition 7** We define  $c\_2Ebool\_2E\_2F\_5C$  to be  $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c\_2Ebool\_2E\_21 2) (\lambda V2t \in 2.V2t))$

Let  $ty\_2Ehreal\_2Ehreal : \iota$  be given. Assume the following.

$$nonempty\ ty\_2Ehreal\_2Ehreal \tag{1}$$

Let  $ty\_2Epair\_2Eprod : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A0.nonempty\ A0 \Rightarrow \forall A1.nonempty\ A1 \Rightarrow nonempty\ (ty\_2Epair\_2Eprod\ A0\ A1) \tag{2}$$

Let  $ty\_2Erealax\_2Ereal : \iota$  be given. Assume the following.

$$nonempty\ ty\_2Erealax\_2Ereal \tag{3}$$

Let  $c\_2Erealax\_2Ereal\_REP\_CLASS : \iota$  be given. Assume the following.

$$c\_2Erealax\_2Ereal\_REP\_CLASS \in ((2^{(ty\_2Epair\_2Eprod\ ty\_2Ehreal\_2Ehreal\ ty\_2Ehreal\_2Ehreal)})\ ty\_2Erealax\_2Ereal) \tag{4}$$

**Definition 8** We define  $c\_2Emin\_2E\_40$  to be  $\lambda A.\lambda P \in 2^A.if\ (\exists x \in A.p\ (ap\ P\ x))\ then\ (the\ (\lambda x.x \in A \wedge p$  of type  $\iota \Rightarrow \iota$ .

**Definition 9** We define  $c\_2Erealax\_2Ereal\_REP$  to be  $\lambda V0a \in ty\_2Erealax\_2Ereal.(ap (c\_2Emin\_2E.40 (ty\_2Erealax\_2Ereal\_neg : \iota$  be given. Assume the following.

$$c\_2Erealax\_2Ereal\_neg \in ((ty\_2Epair\_2Eprod ty\_2Ehreal\_2Ehreal ty\_2Ehreal\_2Ehreal)(ty\_2Epair\_2Eprod ty\_2Ehreal\_2Ehreal ty\_2Ehreal\_2Ehreal)) \quad (5)$$

Let  $c\_2Erealax\_2Ereal\_eq : \iota$  be given. Assume the following.

$$c\_2Erealax\_2Ereal\_eq \in ((2^{(ty\_2Epair\_2Eprod ty\_2Ehreal\_2Ehreal ty\_2Ehreal\_2Ehreal)})(ty\_2Epair\_2Eprod ty\_2Ehreal\_2Ehreal ty\_2Ehreal\_2Ehreal)) \quad (6)$$

Let  $c\_2Erealax\_2Ereal\_ABS\_CLASS : \iota$  be given. Assume the following.

$$c\_2Erealax\_2Ereal\_ABS\_CLASS \in (ty\_2Erealax\_2Ereal)^{(2^{(ty\_2Epair\_2Eprod ty\_2Ehreal\_2Ehreal ty\_2Ehreal\_2Ehreal)}} \quad (7)$$

**Definition 10** We define  $c\_2Erealax\_2Ereal\_ABS$  to be  $\lambda V0r \in (ty\_2Epair\_2Eprod ty\_2Ehreal\_2Ehreal ty\_2Ehreal\_2Ehreal)$

**Definition 11** We define  $c\_2Erealax\_2Ereal\_neg$  to be  $\lambda V0T1 \in ty\_2Erealax\_2Ereal.(ap c\_2Erealax\_2Ereal$

Let  $c\_2Erealax\_2Ereal\_add : \iota$  be given. Assume the following.

$$c\_2Erealax\_2Ereal\_add \in (((ty\_2Epair\_2Eprod ty\_2Ehreal\_2Ehreal ty\_2Ehreal\_2Ehreal)(ty\_2Epair\_2Eprod ty\_2Ehreal\_2Ehreal ty\_2Ehreal\_2Ehreal))^{(ty\_2Epair\_2Eprod ty\_2Ehreal\_2Ehreal ty\_2Ehreal\_2Ehreal)}) \quad (8)$$

**Definition 12** We define  $c\_2Erealax\_2Ereal\_add$  to be  $\lambda V0T1 \in ty\_2Erealax\_2Ereal.\lambda V1T2 \in ty\_2Erealax\_2Ereal$

**Definition 13** We define  $c\_2Ereal\_2Ereal\_sub$  to be  $\lambda V0x \in ty\_2Erealax\_2Ereal.\lambda V1y \in ty\_2Erealax\_2Ereal$

Let  $c\_2Erealax\_2Ereal\_mul : \iota$  be given. Assume the following.

$$c\_2Erealax\_2Ereal\_mul \in (((ty\_2Epair\_2Eprod ty\_2Ehreal\_2Ehreal ty\_2Ehreal\_2Ehreal)(ty\_2Epair\_2Eprod ty\_2Ehreal\_2Ehreal ty\_2Ehreal\_2Ehreal))^{(ty\_2Epair\_2Eprod ty\_2Ehreal\_2Ehreal ty\_2Ehreal\_2Ehreal)}) \quad (9)$$

**Definition 14** We define  $c\_2Erealax\_2Ereal\_mul$  to be  $\lambda V0T1 \in ty\_2Erealax\_2Ereal.\lambda V1T2 \in ty\_2Erealax\_2Ereal$

**Definition 15** We define  $c\_2Ereal\_topology\_2Elinear$  to be  $\lambda V0f \in (ty\_2Erealax\_2Ereal)^{ty\_2Erealax\_2Ereal}$

Assume the following.

$$True \quad (10)$$

Assume the following.

$$\forall A\_27a.nonempty A\_27a \Rightarrow (\forall V0t \in 2.((\forall V1x \in A\_27a.(p V0t) \Leftrightarrow (p V0t))) \Leftrightarrow (p V0t))) \quad (11)$$

Assume the following.

$$(\forall V0t \in 2.(((True \Rightarrow (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \Rightarrow True) \Leftrightarrow True) \wedge (((False \Rightarrow (p V0t)) \Leftrightarrow True) \wedge (((p V0t) \Rightarrow (p V0t)) \Leftrightarrow True) \wedge ((p V0t) \Rightarrow False) \Leftrightarrow (\neg (p V0t)))))) \quad (12)$$

Assume the following.

$$\forall A\_27a.nonempty\ A\_27a \Rightarrow (\forall V0x \in A\_27a. ((V0x = V0x) \Leftrightarrow True)) \quad (13)$$

Assume the following.

$$\forall A\_27a.nonempty\ A\_27a \Rightarrow (\forall V0x \in A\_27a. (\forall V1y \in A\_27a. ((V0x = V1y) \Leftrightarrow (V1y = V0x)))) \quad (14)$$

Assume the following.

$$(\forall V0t \in 2. (((True \Leftrightarrow (p\ V0t)) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \Leftrightarrow True) \Leftrightarrow (p\ V0t)) \wedge (((False \Leftrightarrow (p\ V0t)) \Leftrightarrow \neg(p\ V0t)) \wedge (((p\ V0t) \Leftrightarrow False) \Leftrightarrow \neg(p\ V0t)))))) \quad (15)$$

Assume the following.

$$(\forall V0t1 \in 2. (\forall V1t2 \in 2. (\forall V2t3 \in 2. (((p\ V0t1) \Rightarrow ((p\ V1t2) \Rightarrow (p\ V2t3))) \Leftrightarrow (((p\ V0t1) \wedge (p\ V1t2)) \Rightarrow (p\ V2t3)))))) \quad (16)$$

Assume the following.

$$(\forall V0x \in 2. (\forall V1x\_27 \in 2. (\forall V2y \in 2. (\forall V3y\_27 \in 2. (((p\ V0x) \Leftrightarrow (p\ V1x\_27)) \wedge ((p\ V1x\_27) \Rightarrow ((p\ V2y) \Leftrightarrow (p\ V3y\_27)))))) \Rightarrow (((p\ V0x) \Rightarrow (p\ V2y)) \Leftrightarrow ((p\ V1x\_27) \Rightarrow (p\ V3y\_27)))))) \quad (17)$$

Assume the following.

$$(\forall V0f \in (ty\_2Erealax\_2Ereal^{ty\_2Erealax\_2Ereal}). (\forall V1x \in ty\_2Erealax\_2Ereal. ((p\ (ap\ c\_2Ereal\_topology\_2Elinear\ V0f)) \Rightarrow ((ap\ V0f\ (ap\ c\_2Erealax\_2Ereal\_neg\ V1x)) = (ap\ c\_2Erealax\_2Ereal\_neg\ (ap\ V0f\ V1x)))))) \quad (18)$$

Assume the following.

$$(\forall V0f \in (ty\_2Erealax\_2Ereal^{ty\_2Erealax\_2Ereal}). (\forall V1x \in ty\_2Erealax\_2Ereal. (\forall V2y \in ty\_2Erealax\_2Ereal. ((p\ (ap\ c\_2Ereal\_topology\_2Elinear\ V0f)) \Rightarrow ((ap\ V0f\ (ap\ (ap\ c\_2Erealax\_2Ereal\_add\ V1x\ V2y)) = (ap\ (ap\ c\_2Erealax\_2Ereal\_add\ (ap\ V0f\ V1x))\ (ap\ V0f\ V2y)))))) \quad (19)$$

### Theorem 1

$$(\forall V0f \in (ty\_2Erealax\_2Ereal^{ty\_2Erealax\_2Ereal}). (\forall V1x \in ty\_2Erealax\_2Ereal. (\forall V2y \in ty\_2Erealax\_2Ereal. ((p\ (ap\ c\_2Ereal\_topology\_2Elinear\ V0f)) \Rightarrow ((ap\ V0f\ (ap\ (ap\ c\_2Ereal\_2Ereal\_sub\ V1x\ V2y)) = (ap\ (ap\ c\_2Ereal\_2Ereal\_sub\ (ap\ V0f\ V1x))\ (ap\ V0f\ V2y)))))) \quad (19)$$