

thm_2Ereal_topology_2ELINEAR_SUM_MUL (TMa8pydSDCdhN2RniLb6AA8XN4hSHKnqKdd)

October 26, 2020

Definition 1 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define $c_2Ebool_2E_2T$ to be $(ap (ap (c_2Emin_2E_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

Definition 3 We define $c_2Ebool_2E_21$ to be $\lambda A.27a : \iota.(\lambda V0P \in (2^{A-27a}).(ap (ap (c_2Emin_2E_3D (2^{A-27a}))$

Definition 4 We define $c_2Ebool_2E_2F$ to be $(ap (c_2Ebool_2E_21 2) (\lambda V0t \in 2.V0t))$.

Definition 5 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p P \Rightarrow p Q)$ of type ι .

Definition 6 We define $c_2Ebool_2E_27E$ to be $(\lambda V0t \in 2.(ap (ap c_2Emin_2E_3D_3D_3E V0t) c_2Ebool_2E_2F$

Let $ty_2Ehreal_2Ehreal : \iota$ be given. Assume the following.

$$nonempty\ ty_2Ehreal_2Ehreal \tag{1}$$

Let $ty_2Epair_2Eprod : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A0.nonempty\ A0 \Rightarrow \forall A1.nonempty\ A1 \Rightarrow nonempty\ (ty_2Epair_2Eprod\ A0\ A1) \tag{2}$$

Let $ty_2Erealax_2Ereal : \iota$ be given. Assume the following.

$$nonempty\ ty_2Erealax_2Ereal \tag{3}$$

Let $c_2Erealax_2Ereal_REP_CLASS : \iota$ be given. Assume the following.

$$c_2Erealax_2Ereal_REP_CLASS \in ((2^{(ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal\ ty_2Ehreal_2Ehreal)})\ ty_2Erealax_2Ereal) \tag{4}$$

Definition 7 We define $c_2Emin_2E_40$ to be $\lambda A.\lambda P \in 2^A.if (\exists x \in A.p (ap P x)) \mathbf{then} (the (\lambda x.x \in A \wedge p$ of type $\iota \Rightarrow \iota$.

Definition 8 We define $c_2Erealax_2Ereal_REP$ to be $\lambda V0a \in ty_2Erealax_2Ereal.(ap (c_2Emin_2E_40 (ty$

Definition 18 We define c_2Ebool_2ECOND to be $\lambda A_27a : \iota. (\lambda V0t \in 2. (\lambda V1t1 \in A_27a. (\lambda V2t2 \in A_27a. ($

Definition 19 We define $c_2Ebool_2E_5C_2F$ to be $(\lambda V0t1 \in 2. (\lambda V1t2 \in 2. (ap (c_2Ebool_2E_21 2) (\lambda V2t \in$

Definition 20 We define $c_2Epred_set_2EINSERT$ to be $\lambda A_27a : \iota. \lambda V0x \in A_27a. \lambda V1s \in (2^{A_27a}). (ap (c_2$

Definition 21 We define $c_2Epred_set_2EEMPTY$ to be $\lambda A_27a : \iota. (\lambda V0x \in A_27a. c_2Ebool_2EF).$

Definition 22 We define $c_2Epred_set_2EFINITE$ to be $\lambda A_27a : \iota. \lambda V0s \in (2^{A_27a}). (ap (c_2Ebool_2E_21 2)$

Definition 23 We define $c_2Eiterate_2EITSET$ to be $\lambda A_27a : \iota. \lambda A_27b : \iota. \lambda V0f \in ((A_27a^{A_27a})^{A_27b}). \lambda V$

Definition 24 We define $c_2Eiterate_2Eiterate$ to be $\lambda A_27a : \iota. \lambda A_27b : \iota. \lambda V0op \in ((A_27b^{A_27b})^{A_27b}). \lambda V$

Definition 25 We define $c_2Eiterate_2ESum$ to be $\lambda A_27a : \iota. (ap (c_2Eiterate_2Eiterate A_27a ty_2Erealax$

Definition 26 We define $c_2Ereal_topology_2Elinear$ to be $\lambda V0f \in (ty_2Erealax_2Ereal^{ty_2Erealax_2Ereal})$

Assume the following.

$$True \tag{11}$$

Assume the following.

$$\forall A_27a. nonempty A_27a \Rightarrow (\forall V0t \in 2. ((\forall V1x \in A_27a. (p V0t)) \Leftrightarrow (p V0t))) \tag{12}$$

Assume the following.

$$\begin{aligned} & (\forall V0t \in 2. (((True \wedge (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \wedge True) \Leftrightarrow \\ & (p V0t)) \wedge (((False \wedge (p V0t)) \Leftrightarrow False) \wedge (((p V0t) \wedge False) \Leftrightarrow False) \wedge \\ & (((p V0t) \wedge (p V0t)) \Leftrightarrow (p V0t)))))) \end{aligned} \tag{13}$$

Assume the following.

$$\begin{aligned} & (\forall V0t \in 2. (((True \Rightarrow (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \Rightarrow True) \Leftrightarrow \\ & True) \wedge (((False \Rightarrow (p V0t)) \Leftrightarrow True) \wedge (((p V0t) \Rightarrow (p V0t)) \Leftrightarrow True) \wedge ((\\ & (p V0t) \Rightarrow False) \Leftrightarrow \neg (p V0t)))))) \end{aligned} \tag{14}$$

Assume the following.

$$\forall A_27a. nonempty A_27a \Rightarrow (\forall V0x \in A_27a. ((V0x = V0x) \Leftrightarrow True)) \tag{15}$$

Assume the following.

$$\forall A_27a. nonempty A_27a \Rightarrow (\forall V0x \in A_27a. (\forall V1y \in A_27a. ((V0x = V1y) \Leftrightarrow (V1y = V0x)))) \tag{16}$$

Assume the following.

$$\begin{aligned} & (\forall V0t \in 2. (((True \Leftrightarrow (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \Leftrightarrow True) \Leftrightarrow \\ & (p V0t)) \wedge (((False \Leftrightarrow (p V0t)) \Leftrightarrow \neg (p V0t)) \wedge (((p V0t) \Leftrightarrow False) \Leftrightarrow \neg (\\ & p V0t)))))) \end{aligned} \tag{17}$$

Assume the following.

$$(\forall V0t1 \in 2.(\forall V1t2 \in 2.(\forall V2t3 \in 2.(((p V0t1) \Rightarrow ((p V1t2) \Rightarrow (p V2t3))) \Leftrightarrow (((p V0t1) \wedge (p V1t2)) \Rightarrow (p V2t3)))))) \quad (18)$$

Assume the following.

$$(\forall V0x \in 2.(\forall V1x_{.27} \in 2.(\forall V2y \in 2.(\forall V3y_{.27} \in 2.(((p V0x) \Leftrightarrow (p V1x_{.27})) \wedge ((p V1x_{.27}) \Rightarrow ((p V2y) \Leftrightarrow (p V3y_{.27})))))) \Rightarrow ((p V0x) \Rightarrow (p V2y)) \Leftrightarrow ((p V1x_{.27}) \Rightarrow (p V3y_{.27})))))) \quad (19)$$

Assume the following.

$$(\forall V0f \in (ty_2Erealax_2Ereal^{ty_2Erealax_2Ereal}).(\forall V1c \in ty_2Erealax_2Ereal.(\forall V2x \in ty_2Erealax_2Ereal.((p (ap c_2Ereal_topology_2Elinear V0f)) \Rightarrow ((ap V0f (ap (ap c_2Erealax_2Ereal_mul V1c) V2x)) = (ap (ap c_2Erealax_2Ereal_mul V1c) (ap V0f V2x))))))) \quad (20)$$

Assume the following.

$$\forall A_{.27a}.nonempty A_{.27a} \Rightarrow (\forall V0f \in (ty_2Erealax_2Ereal^{ty_2Erealax_2Ereal}).(\forall V1g \in (ty_2Erealax_2Ereal^{A_{.27a}}).(\forall V2s \in (2^{A_{.27a}}).(((p (ap c_2Ereal_topology_2Elinear V0f)) \wedge (p (ap (c_2Epred_set_2EFINITE A_{.27a}) V2s))) \Rightarrow ((ap V0f (ap (ap (c_2Eiterate_2ESum A_{.27a}) V2s) V1g)) = (ap (ap (c_2Eiterate_2ESum A_{.27a}) V2s) (ap (ap (c_2Ecombin_2Eo A_{.27a} ty_2Erealax_2Ereal ty_2Erealax_2Ereal) V0f) V1g))))))) \quad (21)$$

Theorem 1

$$\forall A_{.27a}.nonempty A_{.27a} \Rightarrow (\forall V0f \in (ty_2Erealax_2Ereal^{ty_2Erealax_2Ereal}).(\forall V1s \in (2^{A_{.27a}}).(\forall V2c \in (ty_2Erealax_2Ereal^{A_{.27a}}).(\forall V3v \in (ty_2Erealax_2Ereal^{A_{.27a}}).(((p (ap c_2Ereal_topology_2Elinear V0f)) \wedge (p (ap (c_2Epred_set_2EFINITE A_{.27a}) V1s))) \Rightarrow ((ap V0f (ap (ap (c_2Eiterate_2ESum A_{.27a}) V1s) (\lambda V4i \in A_{.27a}.(ap (ap c_2Erealax_2Ereal_mul (ap V2c V4i)) (ap V3v V4i)))))) = (ap (ap (c_2Eiterate_2ESum A_{.27a}) V1s) (\lambda V5i \in A_{.27a}.(ap (ap c_2Erealax_2Ereal_mul (ap V2c V5i)) (ap V0f (ap V3v V5i)))))))))))))$$