

# thm\_2Ereal\_topology\_2ENOT\_EVENTUALLY (TMP4UWRHd7BwJvqPMb5BLt5SBo26RYVG2rA)

October 26, 2020

**Definition 1** We define  $c\_2Emin\_2E\_3D$  to be  $\lambda A.\lambda x \in A.\lambda y \in A.inj\_o (x = y)$  of type  $\iota \Rightarrow \iota$ .

**Definition 2** We define  $c\_2Ebool\_2E\_2T$  to be  $(ap (ap (c\_2Emin\_2E\_3D (2^2))) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x)$

Let  $ty\_2Ereal\_topology\_2E\_2net : \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A0.nonempty A0 \Rightarrow nonempty (ty\_2Ereal\_topology\_2E\_2net A0) \quad (1)$$

Let  $c\_2Ereal\_topology\_2E\_2netord : \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A\_27a.nonempty A\_27a \Rightarrow c\_2Ereal\_topology\_2E\_2netord A\_27a \in (((2^{A\_27a})^{A\_27a})^{(ty\_2Ereal\_topology\_2E\_2net A\_27a)}) \quad (2)$$

**Definition 3** We define  $c\_2Emin\_2E\_40$  to be  $\lambda A.\lambda P \in 2^A.if (\exists x \in A.p (ap P x))$  then (the  $(\lambda x.x \in A \wedge p x)$ ) of type  $\iota \Rightarrow \iota$ .

**Definition 4** We define  $c\_2Ebool\_2E\_3F$  to be  $\lambda A\_27a : \iota.(\lambda V0P \in (2^{A\_27a}).(ap V0P (ap (c\_2Emin\_2E\_40 A\_27a) P)))$

**Definition 5** We define  $c\_2Ebool\_2E\_21$  to be  $\lambda A\_27a : \iota.(\lambda V0P \in (2^{A\_27a}).(ap (ap (c\_2Emin\_2E\_3D (2^{A\_27a}) P) P)))$

**Definition 6** We define  $c\_2Ebool\_2E\_2F$  to be  $(ap (c\_2Ebool\_2E\_21 2) (\lambda V0t \in 2.V0t))$ .

**Definition 7** We define  $c\_2Emin\_2E\_3D\_3D\_3E$  to be  $\lambda P \in 2.\lambda Q \in 2.inj\_o (p P \Rightarrow p Q)$  of type  $\iota$ .

**Definition 8** We define  $c\_2Ebool\_2E\_7E$  to be  $(\lambda V0t \in 2.(ap (ap c\_2Emin\_2E\_3D\_3D\_3E V0t) c\_2Ebool\_2E\_2F))$

**Definition 9** We define  $c\_2Ebool\_2E\_2F\_5C$  to be  $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c\_2Ebool\_2E\_21 2) (\lambda V2t \in 2.V2t))))$

**Definition 10** We define  $c\_2Ebool\_2E\_5C\_2F$  to be  $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c\_2Ebool\_2E\_21 2) (\lambda V2t \in 2.V2t))))$

**Definition 11** We define  $c\_2Ereal\_topology\_2E\_2trivial\_limit$  to be  $\lambda A\_27a : \iota.\lambda V0net \in (ty\_2Ereal\_topology\_2E\_2net A\_27a)$

**Definition 12** We define  $c\_2Ereal\_topology\_2E\_2eventually$  to be  $\lambda A\_27a : \iota.\lambda V0p \in (2^{A\_27a}).\lambda V1net \in (ty\_2Ereal\_topology\_2E\_2net A\_27a)$

Assume the following.

$$True \quad (3)$$

Assume the following.

$$(\forall V0t1 \in 2. (\forall V1t2 \in 2. (((p V0t1) \Rightarrow (p V1t2)) \Rightarrow (((p V1t2) \Rightarrow (p V0t1)) \Rightarrow ((p V0t1) \Leftrightarrow (p V1t2)))))) \quad (4)$$

Assume the following.

$$(\forall V0t \in 2. (((p V0t) \Rightarrow False) \Rightarrow (\neg(p V0t)))) \quad (5)$$

Assume the following.

$$(\forall V0t \in 2. ((\neg(p V0t)) \Rightarrow ((p V0t) \Rightarrow False))) \quad (6)$$

Assume the following.

$$(\forall V0t \in 2. (((True) \Rightarrow (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \Rightarrow True) \Leftrightarrow True) \wedge (((False) \Rightarrow (p V0t)) \Leftrightarrow True) \wedge (((p V0t) \Rightarrow (p V0t)) \Leftrightarrow True) \wedge (((p V0t) \Rightarrow False) \Leftrightarrow (\neg(p V0t)))) \quad (7)$$

Assume the following.

$$((\forall V0t \in 2. ((\neg(\neg(p V0t))) \Leftrightarrow (p V0t)) \wedge ((\neg True) \Leftrightarrow False) \wedge ((\neg False) \Leftrightarrow True))) \quad (8)$$

Assume the following.

$$\forall A.27a.nonempty A.27a \Rightarrow (\forall V0x \in A.27a. (\forall V1y \in A.27a. ((V0x = V1y) \Leftrightarrow (V1y = V0x)))) \quad (9)$$

Assume the following.

$$(\forall V0t \in 2. (((True) \Leftrightarrow (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \Leftrightarrow True) \Leftrightarrow (p V0t)) \wedge (((False) \Leftrightarrow (p V0t)) \Leftrightarrow (\neg(p V0t))) \wedge (((p V0t) \Leftrightarrow False) \Leftrightarrow (\neg(p V0t)))) \quad (10)$$

Assume the following.

$$\forall A.27a.nonempty A.27a \Rightarrow (\forall V0P \in (2^{A.27a}). ((\neg(\forall V1x \in A.27a. (p (ap V0P V1x)))) \Leftrightarrow (\exists V2x \in A.27a. (\neg(p (ap V0P V2x)))))) \quad (11)$$

Assume the following.

$$\forall A.27a.nonempty A.27a \Rightarrow (\forall V0P \in 2. (\forall V1Q \in (2^{A.27a}). (((p V0P) \vee (\exists V2x \in A.27a. (p (ap V1Q V2x)))) \Leftrightarrow (\exists V3x \in A.27a. ((p V0P) \vee (p (ap V1Q V3x)))))) \quad (12)$$

Assume the following.

$$\forall A.27a.nonempty A.27a \Rightarrow (\forall V0P \in (2^{A.27a}). (\forall V1Q \in 2. ((\exists V2x \in A.27a. ((p (ap V0P V2x)) \wedge (p V1Q))) \Leftrightarrow ((\exists V3x \in A.27a. (p (ap V0P V3x))) \wedge (p V1Q)))) \quad (13)$$

Assume the following.

$$\forall A.27a.nonempty A.27a \Rightarrow (\forall V0P \in 2. (\forall V1Q \in (2^{A.27a}). ((\exists V2x \in A.27a. ((p V0P) \wedge (p (ap V1Q V2x)))) \Leftrightarrow ((p V0P) \wedge (\exists V3x \in A.27a. (p (ap V1Q V3x)))))))) \quad (14)$$

Assume the following.

$$(\forall V0A \in 2. (\forall V1B \in 2. (((p V0A) \vee (p V1B)) \Leftrightarrow ((p V1B) \vee (p V0A)))) \quad (15)$$

Assume the following.

$$(\forall V0t \in 2. ((\neg(\neg(p V0t))) \Leftrightarrow (p V0t))) \quad (16)$$

Assume the following.

$$(\forall V0A \in 2. ((p V0A) \Rightarrow ((\neg(p V0A)) \Rightarrow False))) \quad (17)$$

Assume the following.

$$(\forall V0A \in 2. (\forall V1B \in 2. (((\neg((p V0A) \vee (p V1B))) \Rightarrow False) \Leftrightarrow (((p V0A) \Rightarrow False) \Rightarrow ((\neg(p V1B)) \Rightarrow False)))) \quad (18)$$

Assume the following.

$$(\forall V0A \in 2. (\forall V1B \in 2. (((\neg(\neg(p V0A) \vee (p V1B))) \Rightarrow False) \Leftrightarrow ((p V0A) \Rightarrow ((\neg(p V1B)) \Rightarrow False)))) \quad (19)$$

Assume the following.

$$(\forall V0A \in 2. (((\neg(p V0A)) \Rightarrow False) \Rightarrow (((p V0A) \Rightarrow False) \Rightarrow False))) \quad (20)$$

Assume the following.

$$(\forall V0p \in 2. (\forall V1q \in 2. (\forall V2r \in 2. (((p V0p) \Leftrightarrow ((p V1q) \Leftrightarrow (p V2r))) \Leftrightarrow (((p V0p) \vee ((p V1q) \vee (p V2r))) \wedge (((p V0p) \vee (\neg(p V2r))) \vee (\neg(p V1q)))) \wedge (((p V1q) \vee ((\neg(p V2r)) \vee (\neg(p V0p)))) \wedge ((p V2r) \vee ((\neg(p V1q)) \vee (\neg(p V0p)))))))))) \quad (21)$$

Assume the following.

$$(\forall V0p \in 2. (\forall V1q \in 2. (\forall V2r \in 2. (((p V0p) \Leftrightarrow ((p V1q) \wedge (p V2r))) \Leftrightarrow (((p V0p) \vee ((\neg(p V1q)) \vee (\neg(p V2r)))) \wedge (((p V1q) \vee (\neg(p V0p))) \wedge ((p V2r) \vee (\neg(p V0p)))))))) \quad (22)$$

Assume the following.

$$(\forall V0p \in 2. (\forall V1q \in 2. (\forall V2r \in 2. (((p V0p) \Leftrightarrow ((p V1q) \vee (p V2r))) \Leftrightarrow (((p V0p) \vee (\neg(p V1q))) \wedge (((p V0p) \vee (\neg(p V2r)))) \wedge ((p V1q) \vee ((p V2r) \vee (\neg(p V0p)))))))) \quad (23)$$

Assume the following.

$$\begin{aligned}
& (\forall V0p \in 2. (\forall V1q \in 2. (\forall V2r \in 2. (((p \ V0p) \Leftrightarrow ( \\
& (p \ V1q) \Rightarrow (p \ V2r))) \Leftrightarrow (((p \ V0p) \vee (p \ V1q)) \wedge (((p \ V0p) \vee \neg(p \ V2r))) \wedge ( \\
& \neg(p \ V1q)) \vee ((p \ V2r) \vee \neg(p \ V0p)))))))))) \quad (24)
\end{aligned}$$

Assume the following.

$$\begin{aligned}
& (\forall V0p \in 2. (\forall V1q \in 2. (((p \ V0p) \Leftrightarrow \neg(p \ V1q))) \Leftrightarrow (((p \ V0p) \vee \\
& (p \ V1q)) \wedge (\neg(p \ V1q)) \vee \neg(p \ V0p)))))) \quad (25)
\end{aligned}$$

**Theorem 1**

$$\begin{aligned}
& \forall A.27a.nonempty \ A.27a \Rightarrow (\forall V0net \in (ty.2Ereal\_topology.2Enet \\
& A.27a). (\forall V1p \in (2^{A.27a}). (((\forall V2x \in A.27a. (\neg(p \ (ap \\
& V1p \ V2x)))) \wedge (\neg(p \ (ap \ (c.2Ereal\_topology.2Etrivial\_limit \ A.27a) \\
& V0net)))) \Rightarrow (\neg(p \ (ap \ (ap \ (c.2Ereal\_topology.2Eventually \ A.27a) \\
& V1p) \ V0net))))))
\end{aligned}$$