

thm_2Ereal__topology_2EOPEN__CONTAINS__INTERVAL (TMFZ1Rh8LJNJcsRDsta7qgCZ9b4Emx451ek)

October 26, 2020

Definition 1 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o(x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define c_2Ebool_2ET to be $(ap (ap (c_2Emin_2E_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

Let $ty_2Elist_2Elist : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A0.nonempty A0 \Rightarrow nonempty (ty_2Elist_2Elist A0) \quad (1)$$

Let $c_2Elist_2ENIL : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A_27a.nonempty A_27a \Rightarrow c_2Elist_2ENIL A_27a \in (ty_2Elist_2Elist A_27a) \quad (2)$$

Let $ty_2Epair_2Eprod : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A0.nonempty A0 \Rightarrow \forall A1.nonempty A1 \Rightarrow nonempty (ty_2Epair_2Eprod A0 A1) \quad (3)$$

Let $c_2Elist_2ECONS : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A_27a.nonempty A_27a \Rightarrow c_2Elist_2ECONS A_27a \in (((ty_2Elist_2Elist A_27a)^{(ty_2Elist_2Elist A_27a)})^{A_27a}) \quad (4)$$

Let $ty_2Erealax_2Ereal : \iota$ be given. Assume the following.

$$nonempty ty_2Erealax_2Ereal \quad (5)$$

Let $c_2Elist_2EHD : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A_27a.nonempty A_27a \Rightarrow c_2Elist_2EHD A_27a \in (A_27a^{(ty_2Elist_2Elist A_27a)}) \quad (6)$$

Let $c_2Epair_2ESND : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned} \forall A_27a.nonempty\ A_27a \Rightarrow \forall A_27b.nonempty\ A_27b \Rightarrow c_2Epair_2ESND \\ A_27a\ A_27b \in (A_27b)^{(ty_2Epair_2Eprod\ A_27a\ A_27b)} \end{aligned} \quad (7)$$

Let $ty_2Ehreal_2Ehreal : \iota$ be given. Assume the following.

$$nonempty\ ty_2Ehreal_2Ehreal \quad (8)$$

Let $c_2Erealax_2Ereal_REP_CLASS : \iota$ be given. Assume the following.

$$c_2Erealax_2Ereal_REP_CLASS \in ((2^{(ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal\ ty_2Ehreal_2Ehreal)})^{ty_2Erealax_2Ereal_REP_CLASS}) \quad (9)$$

Definition 3 We define $c_2Emin_2E_40$ to be $\lambda A.\lambda P \in 2^A.$ if $(\exists x \in A.p\ (ap\ P\ x))$ then (the $(\lambda x.x \in A \wedge p\ x)$ of type $\iota \Rightarrow \iota$).

Definition 4 We define $c_2Ebool_2E_21$ to be $\lambda A_27a : \iota.(\lambda V0P \in (2^{A_27a}).(ap\ (ap\ (c_2Emin_2E_3D\ (2^{A_27a}))\ P)\ V0t))$.

Definition 5 We define $c_2Erealax_2Ereal_REP$ to be $\lambda V0a \in ty_2Erealax_2Ereal.(ap\ (c_2Emin_2E_40\ (2^{V0a})))$.

Let $c_2Erealax_2Etrealm_lt : \iota$ be given. Assume the following.

$$c_2Erealax_2Etrealm_lt \in ((2^{(ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal\ ty_2Ehreal_2Ehreal)})^{(ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal)}) \quad (10)$$

Definition 6 We define $c_2Erealax_2Ereal_lt$ to be $\lambda V0T1 \in ty_2Erealax_2Ereal.\lambda V1T2 \in ty_2Erealax_2Ereal.$

Definition 7 We define $c_2Ebool_2E_2F$ to be $(ap\ (c_2Ebool_2E_21\ 2)\ (\lambda V0t \in 2.V0t))$.

Definition 8 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o\ (p\ P \Rightarrow p\ Q)$ of type ι .

Definition 9 We define $c_2Ebool_2E_7E$ to be $(\lambda V0t \in 2.(ap\ (ap\ c_2Emin_2E_3D_3D_3E\ V0t)\ c_2Ebool_2E_2F))$.

Definition 10 We define $c_2Ereal_2Ereal_lte$ to be $\lambda V0x \in ty_2Erealax_2Ereal.\lambda V1y \in ty_2Erealax_2Ereal.$

Let $c_2Epair_2EFST : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned} \forall A_27a.nonempty\ A_27a \Rightarrow \forall A_27b.nonempty\ A_27b \Rightarrow c_2Epair_2EFST \\ A_27a\ A_27b \in (A_27a)^{(ty_2Epair_2Eprod\ A_27a\ A_27b)} \end{aligned} \quad (11)$$

Definition 11 We define $c_2Ebool_2E_2F_5C$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap\ (c_2Ebool_2E_21\ 2)\ (\lambda V2t \in 2.V2t))))$.

Let $c_2Epair_2EABS_prod : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned} \forall A_27a.nonempty\ A_27a \Rightarrow \forall A_27b.nonempty\ A_27b \Rightarrow c_2Epair_2EABS_prod \\ A_27a\ A_27b \in ((ty_2Epair_2Eprod\ A_27a\ A_27b)^{(2^{A_27b})^{A_27a}}) \end{aligned} \quad (12)$$

Definition 12 We define $c_2Epair_2E_2C$ to be $\lambda A_27a : \iota. \lambda A_27b : \iota. \lambda V0x \in A_27a. \lambda V1y \in A_27b. (ap (c_2Epred_set_2EGSPEC : \iota \Rightarrow \iota \Rightarrow \iota)$ be given. Assume the following.

$$\begin{aligned} \forall A_27a.nonempty\ A_27a \Rightarrow \forall A_27b.nonempty\ A_27b \Rightarrow c_2Epred_set_2EGSPEC \\ A_27a\ A_27b \in ((2^{A_27a})^{(ty_2Epair_2Eprod\ A_27a\ 2)^{A_27b}}) \end{aligned} \quad (13)$$

Definition 13 We define $c_2Ereal_topology_2ECLOSED_interval$ to be $\lambda V0l \in (ty_2Elist_2Elist\ (ty_2Epa$

Definition 14 We define c_2Ebool_2EIN to be $\lambda A_27a : \iota. (\lambda V0x \in A_27a. (\lambda V1f \in (2^{A_27a}). (ap\ V1f\ V0x))$

Definition 15 We define $c_2Epred_set_2ESUBSET$ to be $\lambda A_27a : \iota. \lambda V0s \in (2^{A_27a}). \lambda V1t \in (2^{A_27a}). (ap$

Let $c_2Ereal_topology_2EOPEN_interval : \iota$ be given. Assume the following.

$$c_2Ereal_topology_2EOPEN_interval \in ((2^{ty_2Erealax_2Ereal})^{(ty_2Epair_2Eprod\ ty_2Erealax_2Ereal\ ty_2Ereal)} \quad (14)$$

Definition 16 We define $c_2Ebool_2E_3F$ to be $\lambda A_27a : \iota. (\lambda V0P \in (2^{A_27a}). (ap\ V0P\ (ap\ (c_2Emin_2E_40$

Let $c_2Ereal_topology_2EDist : \iota$ be given. Assume the following.

$$c_2Ereal_topology_2EDist \in (ty_2Erealax_2Ereal^{(ty_2Epair_2Eprod\ ty_2Erealax_2Ereal\ ty_2Erealax_2Ereal)}) \quad (15)$$

Let $c_2Enum_2EZERO_REP : \iota$ be given. Assume the following.

$$c_2Enum_2EZERO_REP \in \omega \quad (16)$$

Let $ty_2Enum_2Enum : \iota$ be given. Assume the following.

$$nonempty\ ty_2Enum_2Enum \quad (17)$$

Let $c_2Enum_2EABS_num : \iota$ be given. Assume the following.

$$c_2Enum_2EABS_num \in (ty_2Enum_2Enum^{\omega}) \quad (18)$$

Definition 17 We define c_2Enum_2E0 to be $(ap\ c_2Enum_2EABS_num\ c_2Enum_2EZERO_REP)$.

Let $c_2Ereal_2Ereal_of_num : \iota$ be given. Assume the following.

$$c_2Ereal_2Ereal_of_num \in (ty_2Erealax_2Ereal^{ty_2Enum_2Enum}) \quad (19)$$

Definition 18 We define $c_2Ereal_topology_2EOpen$ to be $\lambda V0s \in (2^{ty_2Erealax_2Ereal}). (ap\ (c_2Ebool_2E_2$

Assume the following.

$$True \quad (20)$$

Assume the following.

$$\begin{aligned} \forall A_27a.nonempty\ A_27a \Rightarrow (\forall V0t \in 2. ((\forall V1x \in \\ A_27a. (p\ V0t)) \Leftrightarrow (p\ V0t))) \end{aligned} \quad (21)$$

Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow (\forall V0x \in A_27a. ((V0x = V0x) \Leftrightarrow True)) \quad (22)$$

Assume the following.

$$\begin{aligned} & ((\forall V0s \in (2^{ty_2Erealax_2Ereal}). ((p (ap\ c_2Ereal_topology_2EOpen \\ & V0s)) \Leftrightarrow (\forall V1x \in ty_2Erealax_2Ereal. ((p (ap (ap (c_2Ebool_2EIN \\ & ty_2Erealax_2Ereal) V1x) V0s)) \Rightarrow (\exists V2a \in ty_2Erealax_2Ereal. \\ & (\exists V3b \in ty_2Erealax_2Ereal. ((p (ap (ap (c_2Ebool_2EIN\ ty_2Erealax_2Ereal) \\ & V1x) (ap\ c_2Ereal_topology_2EOPEN_interval (ap (ap (c_2Epair_2E_2C \\ & ty_2Erealax_2Ereal\ ty_2Erealax_2Ereal) V2a) V3b)))) \wedge (p (ap (\\ & ap (c_2Epred_set_2ESUBSET\ ty_2Erealax_2Ereal) (ap\ c_2Ereal_topology_2ECLOSED_interval \\ & (ap (ap (c_2Elist_2ECONS (ty_2Epair_2Eprod\ ty_2Erealax_2Ereal \\ & ty_2Erealax_2Ereal)) (ap (ap (c_2Epair_2E_2C\ ty_2Erealax_2Ereal \\ & ty_2Erealax_2Ereal) V2a) V3b)) (c_2Elist_2ENIL (ty_2Epair_2Eprod \\ & ty_2Erealax_2Ereal\ ty_2Erealax_2Ereal)))))) V0s)))))) \wedge (\\ & \forall V4s \in (2^{ty_2Erealax_2Ereal}). ((p (ap\ c_2Ereal_topology_2EOpen \\ & V4s)) \Leftrightarrow (\forall V5x \in ty_2Erealax_2Ereal. ((p (ap (ap (c_2Ebool_2EIN \\ & ty_2Erealax_2Ereal) V5x) V4s)) \Rightarrow (\exists V6a \in ty_2Erealax_2Ereal. \\ & (\exists V7b \in ty_2Erealax_2Ereal. ((p (ap (ap (c_2Ebool_2EIN\ ty_2Erealax_2Ereal) \\ & V5x) (ap\ c_2Ereal_topology_2EOPEN_interval (ap (ap (c_2Epair_2E_2C \\ & ty_2Erealax_2Ereal\ ty_2Erealax_2Ereal) V6a) V7b)))) \wedge (p (ap (\\ & ap (c_2Epred_set_2ESUBSET\ ty_2Erealax_2Ereal) (ap\ c_2Ereal_topology_2EOPEN_interval \\ & (ap (ap (c_2Epair_2E_2C\ ty_2Erealax_2Ereal\ ty_2Erealax_2Ereal) \\ & V6a) V7b)) V4s))))))))) \end{aligned} \quad (23)$$

Theorem 1

$$\begin{aligned} & ((\forall V0s \in (2^{ty_2Erealax_2Ereal}). ((p (ap\ c_2Ereal_topology_2EOpen \\ & V0s)) \Leftrightarrow (\forall V1x \in ty_2Erealax_2Ereal. ((p (ap (ap (c_2Ebool_2EIN \\ & ty_2Erealax_2Ereal) V1x) V0s)) \Rightarrow (\exists V2a \in ty_2Erealax_2Ereal. \\ & (\exists V3b \in ty_2Erealax_2Ereal. ((p (ap (ap (c_2Ebool_2EIN\ ty_2Erealax_2Ereal) \\ & V1x) (ap\ c_2Ereal_topology_2EOPEN_interval (ap (ap (c_2Epair_2E_2C \\ & ty_2Erealax_2Ereal\ ty_2Erealax_2Ereal) V2a) V3b)))) \wedge (p (ap (\\ & ap (c_2Epred_set_2ESUBSET\ ty_2Erealax_2Ereal) (ap\ c_2Ereal_topology_2ECLOSED_interval \\ & (ap (ap (c_2Elist_2ECONS (ty_2Epair_2Eprod\ ty_2Erealax_2Ereal \\ & ty_2Erealax_2Ereal)) (ap (ap (c_2Epair_2E_2C\ ty_2Erealax_2Ereal \\ & ty_2Erealax_2Ereal) V2a) V3b)) (c_2Elist_2ENIL (ty_2Epair_2Eprod \\ & ty_2Erealax_2Ereal\ ty_2Erealax_2Ereal)))))) V0s)))))) \end{aligned}$$