

thm_2Ereal__topology_2EOPEN__INTERVAL__LEFT
(TMPDTUySCwKT-
gUcdj5PcMRnn8DvJEAHPVEq)

October 26, 2020

Definition 1 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define $c_2Ebool_2E_2T$ to be $(ap (ap (c_2Emin_2E_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

Definition 3 We define $c_2Ebool_2E_21$ to be $\lambda A_27a : \iota.(\lambda V0P \in (2^{A_27a}).(ap (ap (c_2Emin_2E_3D (2^{A_27a}))$

Definition 4 We define $c_2Ebool_2E_2F$ to be $(ap (c_2Ebool_2E_21 2) (\lambda V0t \in 2.V0t))$.

Definition 5 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p P \Rightarrow p Q)$ of type ι .

Definition 6 We define $c_2Ebool_2E_27E$ to be $(\lambda V0t \in 2.(ap (ap c_2Emin_2E_3D_3D_3E V0t) c_2Ebool_2E_2F))$

Definition 7 We define $c_2Ebool_2E_2F_5C$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21 2) (\lambda V2t \in 2.V2t))$

Let $ty_2Ehreal_2Ehreal : \iota$ be given. Assume the following.

$$nonempty\ ty_2Ehreal_2Ehreal \tag{1}$$

Let $ty_2Epair_2Eprod : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A0.nonempty\ A0 \Rightarrow \forall A1.nonempty\ A1 \Rightarrow nonempty\ (ty_2Epair_2Eprod\ A0\ A1) \tag{2}$$

Let $ty_2Erealax_2Ereal : \iota$ be given. Assume the following.

$$nonempty\ ty_2Erealax_2Ereal \tag{3}$$

Let $c_2Erealax_2Ereal_REP_CLASS : \iota$ be given. Assume the following.

$$c_2Erealax_2Ereal_REP_CLASS \in ((2^{(ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal\ ty_2Ehreal_2Ehreal)})\ ty_2Erealax_2Ereal) \tag{4}$$

Definition 8 We define $c_2Emin_2E_40$ to be $\lambda A.\lambda P \in 2^A.$ **if** $(\exists x \in A.p (ap P x))$ **then** $(the (\lambda x.x \in A \wedge p$
of type $\iota \Rightarrow \iota$.

Definition 9 We define $c_2Erealax_2Ereal_REP$ to be $\lambda V0a \in ty_2Erealax_2Ereal.(ap (c_2Emin_2E_40 (ty$
Let $c_2Erealax_2Ereal_lt : \iota$ be given. Assume the following.

$$c_2Erealax_2Ereal_lt \in ((2^{(ty_2Epair_2Eprod ty_2Ehreal_2Ehreal ty_2Ehreal_2Ehreal)})(ty_2Epair_2Eprod ty_2Ehreal_2Ehreal)) \quad (5)$$

Definition 10 We define $c_2Erealax_2Ereal_lt$ to be $\lambda V0T1 \in ty_2Erealax_2Ereal.\lambda V1T2 \in ty_2Erealax_2Ereal$
Let $c_2Epair_2EABS_prod : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A_27a.nonempty A_27a \Rightarrow \forall A_27b.nonempty A_27b \Rightarrow c_2Epair_2EABS_prod A_27a A_27b \in ((ty_2Epair_2Eprod A_27a A_27b)^{(2^{A_27b})^{A_27a}}) \quad (6)$$

Definition 11 We define $c_2Epair_2E_2C$ to be $\lambda A_27a : \iota.\lambda A_27b : \iota.\lambda V0x \in A_27a.\lambda V1y \in A_27b.(ap (c_2E$
Let $c_2Epred_set_2EGSPEC : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A_27a.nonempty A_27a \Rightarrow \forall A_27b.nonempty A_27b \Rightarrow c_2Epred_set_2EGSPEC A_27a A_27b \in ((2^{A_27a})^{(ty_2Epair_2Eprod A_27a 2)^{A_27b}}) \quad (7)$$

Definition 12 We define c_2Ebool_2EIN to be $\lambda A_27a : \iota.(\lambda V0x \in A_27a.(\lambda V1f \in (2^{A_27a}).(ap V1f V0x))$
Let $c_2Ereal_topology_2EDist : \iota$ be given. Assume the following.

$$c_2Ereal_topology_2EDist \in (ty_2Erealax_2Ereal^{(ty_2Epair_2Eprod ty_2Erealax_2Ereal ty_2Erealax_2Ereal)}) \quad (8)$$

Let $c_2Enum_2EZERO_REP : \iota$ be given. Assume the following.

$$c_2Enum_2EZERO_REP \in \omega \quad (9)$$

Let $ty_2Enum_2Enum : \iota$ be given. Assume the following.

$$nonempty ty_2Enum_2Enum \quad (10)$$

Let $c_2Enum_2EABS_num : \iota$ be given. Assume the following.

$$c_2Enum_2EABS_num \in (ty_2Enum_2Enum^{\omega}) \quad (11)$$

Definition 13 We define c_2Enum_2E0 to be $(ap c_2Enum_2EABS_num c_2Enum_2EZERO_REP)$.

Let $c_2Ereal_2Ereal_of_num : \iota$ be given. Assume the following.

$$c_2Ereal_2Ereal_of_num \in (ty_2Erealax_2Ereal^{ty_2Enum_2Enum}) \quad (12)$$

Definition 14 We define $c_2Ebool_2E_3F$ to be $\lambda A_27a : \iota.(\lambda V0P \in (2^{A_27a}).(ap V0P (ap (c_2Emin_2E_40$

Definition 15 We define $c_Ereal_topology_2EOpen$ to be $\lambda V0s \in (2^{ty_2Erealax_2Ereal}).(ap (c_Ebool_2E_2$

Assume the following.

$$True \quad (13)$$

Assume the following.

$$\forall A_27a.nonempty A_27a \Rightarrow (\forall V0t \in 2.((\forall V1x \in A_27a.(p V0t)) \Leftrightarrow (p V0t))) \quad (14)$$

Assume the following.

$$\begin{aligned} & (\forall V0t \in 2.(((True \Leftrightarrow (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \Leftrightarrow True) \Leftrightarrow \\ & (p V0t)) \wedge (((False \Leftrightarrow (p V0t)) \Leftrightarrow \neg(p V0t)) \wedge (((p V0t) \Leftrightarrow False) \Leftrightarrow \neg(\\ & p V0t)))))) \quad (15) \end{aligned}$$

Assume the following.

$$\begin{aligned} & (\forall V0a \in ty_2Erealax_2Ereal.(p (ap c_Ereal_topology_2EOpen \\ & (ap (c_2Epred_set_2EGSPEC ty_2Erealax_2Ereal ty_2Erealax_2Ereal) \\ & (\lambda V1x \in ty_2Erealax_2Ereal.(ap (ap (c_2Epair_2E_2C ty_2Erealax_2Ereal \\ & 2) V1x) (ap (ap c_2Erealax_2Ereal_lt V1x) V0a)))))) \quad (16) \end{aligned}$$

Theorem 1

$$\begin{aligned} & (\forall V0b \in ty_2Erealax_2Ereal.(p (ap c_Ereal_topology_2EOpen \\ & (ap (c_2Epred_set_2EGSPEC ty_2Erealax_2Ereal ty_2Erealax_2Ereal) \\ & (\lambda V1x \in ty_2Erealax_2Ereal.(ap (ap (c_2Epair_2E_2C ty_2Erealax_2Ereal \\ & 2) V1x) (ap (ap c_2Erealax_2Ereal_lt V1x) V0b)))))) \end{aligned}$$