

# thm\_2Ereal\_\_topology\_2EOPEN\_\_INTERVAL\_\_RIGHT (TMY1RELhCNdsSDKUVq6Q9z5a6BNnmzx7VZV)

October 26, 2020

**Definition 1** We define  $c\_2Emin\_2E\_3D$  to be  $\lambda A.\lambda x \in A.\lambda y \in A.inj\_o (x = y)$  of type  $\iota \Rightarrow \iota$ .

**Definition 2** We define  $c\_2Ebool\_2E\_2T$  to be  $(ap (ap (c\_2Emin\_2E\_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

**Definition 3** We define  $c\_2Ebool\_2E\_21$  to be  $\lambda A.\lambda a : \iota.(\lambda V0P \in (2^{A-27a}).(ap (ap (c\_2Emin\_2E\_3D (2^{A-27a}))$

**Definition 4** We define  $c\_2Ebool\_2E\_2F$  to be  $(ap (c\_2Ebool\_2E\_21 2) (\lambda V0t \in 2.V0t))$ .

**Definition 5** We define  $c\_2Emin\_2E\_3D\_3D\_3E$  to be  $\lambda P \in 2.\lambda Q \in 2.inj\_o (p P \Rightarrow p Q)$  of type  $\iota$ .

**Definition 6** We define  $c\_2Ebool\_2E\_7E$  to be  $(\lambda V0t \in 2.(ap (ap c\_2Emin\_2E\_3D\_3D\_3E V0t) c\_2Ebool\_2E\_2F$

**Definition 7** We define  $c\_2Ebool\_2E\_2F\_5C$  to be  $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c\_2Ebool\_2E\_21 2) (\lambda V2t \in 2.V2t))$

Let  $ty\_2Ehreal\_2Ehreal : \iota$  be given. Assume the following.

$$nonempty\ ty\_2Ehreal\_2Ehreal \tag{1}$$

Let  $ty\_2Epair\_2Eprod : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A0.nonempty\ A0 \Rightarrow \forall A1.nonempty\ A1 \Rightarrow nonempty\ (ty\_2Epair\_2Eprod\ A0\ A1) \tag{2}$$

Let  $ty\_2Erealax\_2Ereal : \iota$  be given. Assume the following.

$$nonempty\ ty\_2Erealax\_2Ereal \tag{3}$$

Let  $c\_2Erealax\_2Ereal\_REP\_CLASS : \iota$  be given. Assume the following.

$$c\_2Erealax\_2Ereal\_REP\_CLASS \in ((2^{(ty\_2Epair\_2Eprod\ ty\_2Ehreal\_2Ehreal\ ty\_2Ehreal\_2Ehreal)})ty\_2Erealax\_2Ereal) \tag{4}$$

**Definition 8** We define  $c\_2Emin\_2E40$  to be  $\lambda A.\lambda P \in 2^A.$ **if**  $(\exists x \in A.p (ap P x))$  **then**  $(the (\lambda x.x \in A \wedge p$   
of type  $\iota \Rightarrow \iota$ .

**Definition 9** We define  $c\_2Erealax\_2Ereal\_REP$  to be  $\lambda V0a \in ty\_2Erealax\_2Ereal.(ap (c\_2Emin\_2E40 (ty$   
Let  $c\_2Erealax\_2Ereal\_lt : \iota$  be given. Assume the following.

$$c\_2Erealax\_2Ereal\_lt \in ((2^{(ty\_2Epair\_2Eprod ty\_2Ehreal\_2Ehreal ty\_2Ehreal\_2Ehreal)})(ty\_2Epair\_2Eprod ty\_2Ehreal\_2Ehreal)) \quad (5)$$

**Definition 10** We define  $c\_2Erealax\_2Ereal\_lt$  to be  $\lambda V0T1 \in ty\_2Erealax\_2Ereal.\lambda V1T2 \in ty\_2Erealax\_2Ereal$

**Definition 11** We define  $c\_2Ereal\_2Ereal\_gt$  to be  $\lambda V0x \in ty\_2Erealax\_2Ereal.\lambda V1y \in ty\_2Erealax\_2Ereal$   
Let  $c\_2Epair\_2EABS\_prod : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A\_27a.nonempty A\_27a \Rightarrow \forall A\_27b.nonempty A\_27b \Rightarrow c\_2Epair\_2EABS\_prod A\_27a A\_27b \in ((ty\_2Epair\_2Eprod A\_27a A\_27b)^{(2^{A\_27b})^{A\_27a}}) \quad (6)$$

**Definition 12** We define  $c\_2Epair\_2E2C$  to be  $\lambda A\_27a : \iota.\lambda A\_27b : \iota.\lambda V0x \in A\_27a.\lambda V1y \in A\_27b.(ap (c\_2E$   
Let  $c\_2Epred\_set\_2EGSPEC : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A\_27a.nonempty A\_27a \Rightarrow \forall A\_27b.nonempty A\_27b \Rightarrow c\_2Epred\_set\_2EGSPEC A\_27a A\_27b \in ((2^{A\_27a})^{(ty\_2Epair\_2Eprod A\_27a 2)^{A\_27b}}) \quad (7)$$

**Definition 13** We define  $c\_2Ebool\_2EIN$  to be  $\lambda A\_27a : \iota.(\lambda V0x \in A\_27a.(\lambda V1f \in (2^{A\_27a}).(ap V1f V0x)))$   
Let  $c\_2Ereal\_topology\_2EDist : \iota$  be given. Assume the following.

$$c\_2Ereal\_topology\_2EDist \in (ty\_2Erealax\_2Ereal^{(ty\_2Epair\_2Eprod ty\_2Erealax\_2Ereal ty\_2Erealax\_2Ereal)}) \quad (8)$$

Let  $c\_2Enum\_2EZERO\_REP : \iota$  be given. Assume the following.

$$c\_2Enum\_2EZERO\_REP \in \omega \quad (9)$$

Let  $ty\_2Enum\_2Enum : \iota$  be given. Assume the following.

$$nonempty ty\_2Enum\_2Enum \quad (10)$$

Let  $c\_2Enum\_2EABS\_num : \iota$  be given. Assume the following.

$$c\_2Enum\_2EABS\_num \in (ty\_2Enum\_2Enum^{\omega}) \quad (11)$$

**Definition 14** We define  $c\_2Enum\_2E0$  to be  $(ap c\_2Enum\_2EABS\_num c\_2Enum\_2EZERO\_REP)$ .

Let  $c\_2Ereal\_2Ereal\_of\_num : \iota$  be given. Assume the following.

$$c\_2Ereal\_2Ereal\_of\_num \in (ty\_2Erealax\_2Ereal^{ty\_2Enum\_2Enum}) \quad (12)$$

**Definition 15** We define  $c\_2Ebool\_2E\_3F$  to be  $\lambda A\_27a : \iota. (\lambda V0P \in (2^{A\_27a}). (ap\ V0P\ (ap\ (c\_2Emin\_2E\_40$

**Definition 16** We define  $c\_2Ereal\_topology\_2EOpen$  to be  $\lambda V0s \in (2^{ty\_2Erealax\_2Ereal}). (ap\ (c\_2Ebool\_2E\_2$

Assume the following.

$$True \tag{13}$$

Assume the following.

$$\forall A\_27a. nonempty\ A\_27a \Rightarrow (\forall V0t \in 2. ((\forall V1x \in A\_27a. (p\ V0t) \Leftrightarrow (p\ V1x))) \Leftrightarrow (p\ V0t))) \tag{14}$$

Assume the following.

$$\forall A\_27a. nonempty\ A\_27a \Rightarrow (\forall V0x \in A\_27a. (\forall V1y \in A\_27a. ((V0x = V1y) \Leftrightarrow (V1y = V0x)))) \tag{15}$$

Assume the following.

$$(\forall V0t \in 2. (((True \Leftrightarrow (p\ V0t)) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \Leftrightarrow True) \Leftrightarrow (p\ V0t)) \wedge (((False \Leftrightarrow (p\ V0t)) \Leftrightarrow \neg(p\ V0t)) \wedge (((p\ V0t) \Leftrightarrow False) \Leftrightarrow \neg(p\ V0t)))))) \tag{16}$$

Assume the following.

$$(\forall V0a \in ty\_2Erealax\_2Ereal. (p\ (ap\ c\_2Ereal\_topology\_2EOpen\ (ap\ (c\_2Epred\_set\_2EGSPEC\ ty\_2Erealax\_2Ereal\ ty\_2Erealax\_2Ereal)\ (\lambda V1x \in ty\_2Erealax\_2Ereal. (ap\ (ap\ (c\_2Epair\_2E\_2C\ ty\_2Erealax\_2Ereal\ 2)\ V1x)\ (ap\ (ap\ c\_2Ereal\_gt\ V1x\ V0a)))))))))) \tag{17}$$

**Theorem 1**

$$(\forall V0a \in ty\_2Erealax\_2Ereal. (p\ (ap\ c\_2Ereal\_topology\_2EOpen\ (ap\ (c\_2Epred\_set\_2EGSPEC\ ty\_2Erealax\_2Ereal\ ty\_2Erealax\_2Ereal)\ (\lambda V1x \in ty\_2Erealax\_2Ereal. (ap\ (ap\ (c\_2Epair\_2E\_2C\ ty\_2Erealax\_2Ereal\ 2)\ V1x)\ (ap\ (ap\ c\_2Ereal\_lt\ V0a\ V1x))))))))))$$