

thm_2Ereal_topology_2EOPEN_IN_OPEN (TMarKZ31MdPGbMsoUah1WaNqic7Rs8b5SYP)

October 26, 2020

Definition 1 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define $c_2Ebool_2E_ET$ to be $(ap (ap (c_2Emin_2E_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

Definition 3 We define $c_2Ebool_2E_EIN$ to be $\lambda A_27a : \iota.(\lambda V0x \in A_27a.(\lambda V1f \in (2^{A_27a}).(ap V1f V0x)))$

Definition 4 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p P \Rightarrow p Q)$ of type ι .

Definition 5 We define $c_2Ebool_2E_21$ to be $\lambda A_27a : \iota.(\lambda V0P \in (2^{A_27a}).(ap (ap (c_2Emin_2E_3D (2^{A_27a}))$

Definition 6 We define $c_2Ebool_2E_2F_5C$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21 2) (\lambda V2t \in 2$

Let $ty_2Epair_2Eprod : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A0.nonempty A0 \Rightarrow \forall A1.nonempty A1 \Rightarrow nonempty (ty_2Epair_2Eprod A0 A1) \tag{1}$$

Let $c_2Epair_2EABS_prod : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A_27a.nonempty A_27a \Rightarrow \forall A_27b.nonempty A_27b \Rightarrow c_2Epair_2EABS_prod A_27a A_27b \in ((ty_2Epair_2Eprod A_27a A_27b)^{(2^{A_27b})^{A_27a}}) \tag{2}$$

Definition 7 We define $c_2Epair_2E_2C$ to be $\lambda A_27a : \iota.\lambda A_27b : \iota.\lambda V0x \in A_27a.\lambda V1y \in A_27b.(ap (c_2E$

Let $c_2Epred_set_2EGSPEC : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A_27a.nonempty A_27a \Rightarrow \forall A_27b.nonempty A_27b \Rightarrow c_2Epred_set_2EGSPEC A_27a A_27b \in ((2^{A_27a})^{((ty_2Epair_2Eprod A_27a 2)^{A_27b})}) \tag{3}$$

Definition 8 We define $c_2Epred_set_2EINTER$ to be $\lambda A_27a : \iota.\lambda V0s \in (2^{A_27a}).\lambda V1t \in (2^{A_27a}).(ap (c_2$

Definition 9 We define $c_2Emin_2E_40$ to be $\lambda A.\lambda P \in 2^A$. **if** $(\exists x \in A.p (ap P x))$ **then** (the $(\lambda x.x \in A \wedge p$ of type $\iota \Rightarrow \iota$).

Definition 10 We define $c_2Ebool_2E_3F$ to be $\lambda A_27a : \iota. (\lambda V0P \in (2^{A_27a}). (ap V0P (ap (c_2Emin_2E_40$

Let $ty_2Etopology_2Etopology : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A0.nonempty A0 \Rightarrow nonempty (ty_2Etopology_2Etopology A0) \quad (4)$$

Let $c_2Etopology_2Eopen_in : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A_27a.nonempty A_27a \Rightarrow c_2Etopology_2Eopen_in A_27a \in ((2^{(2^{A_27a})})^{(ty_2Etopology_2Etopology A_27a)}) \quad (5)$$

Let $c_2Etopology_2Etopology : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A_27a.nonempty A_27a \Rightarrow c_2Etopology_2Etopology A_27a \in ((ty_2Etopology_2Etopology A_27a)^{(2^{(2^{A_27a})})}) \quad (6)$$

Definition 11 We define $c_2Ereal_topology_2Esubtopology$ to be $\lambda A_27a : \iota. \lambda V0top \in (ty_2Etopology_2Etopology$

Let $ty_2Erealax_2Ereal : \iota$ be given. Assume the following.

$$nonempty ty_2Erealax_2Ereal \quad (7)$$

Let $c_2Ereal_topology_2EDist : \iota$ be given. Assume the following.

$$c_2Ereal_topology_2EDist \in (ty_2Erealax_2Ereal^{(ty_2Epair_2Eprod ty_2Erealax_2Ereal ty_2Erealax_2Ereal)}) \quad (8)$$

Let $ty_2Ehreal_2Ehreal : \iota$ be given. Assume the following.

$$nonempty ty_2Ehreal_2Ehreal \quad (9)$$

Let $c_2Erealax_2Ereal_REP_CLASS : \iota$ be given. Assume the following.

$$c_2Erealax_2Ereal_REP_CLASS \in ((2^{(ty_2Epair_2Eprod ty_2Ehreal_2Ehreal ty_2Ehreal_2Ehreal)})^{ty_2Erealax_2Ereal}) \quad (10)$$

Definition 12 We define $c_2Erealax_2Ereal_REP$ to be $\lambda V0a \in ty_2Erealax_2Ereal. (ap (c_2Emin_2E_40 (t$

Let $c_2Erealax_2Etreallt : \iota$ be given. Assume the following.

$$c_2Erealax_2Etreallt \in ((2^{(ty_2Epair_2Eprod ty_2Ehreal_2Ehreal ty_2Ehreal_2Ehreal)})^{(ty_2Epair_2Eprod ty_2Ehreal_2Ehreal)}) \quad (11)$$

Definition 13 We define $c_2Erealax_2Ereal_lt$ to be $\lambda V0T1 \in ty_2Erealax_2Ereal. \lambda V1T2 \in ty_2Erealax_2Ereal.$

Let $c_2Enum_2EZERO_REP : \iota$ be given. Assume the following.

$$c_2Enum_2EZERO_REP \in \omega \quad (12)$$

Let $ty_2Enum_2Enum : \iota$ be given. Assume the following.

$$nonempty ty_2Enum_2Enum \quad (13)$$

Let $c_2Enum_2EABS_num : \iota$ be given. Assume the following.

$$c_2Enum_2EABS_num \in (ty_2Enum_2Enum^{\omega}) \quad (14)$$

Definition 14 We define c_Enum_2E0 to be $(ap\ c_Enum_2EABS_num\ c_Enum_2EZERO_REP)$.

Let $c_Ereal_2Ereal_of_num : \iota$ be given. Assume the following.

$$c_Ereal_2Ereal_of_num \in (ty_2Erealax_2Ereal^{ty_2Enum_2Enum}) \quad (15)$$

Definition 15 We define $c_Ereal_topology_2EOpen$ to be $\lambda V0s \in (2^{ty_2Erealax_2Ereal}).(ap\ (c_Ebool_2E2$

Definition 16 We define $c_Ereal_topology_2Eeuclidean$ to be $(ap\ (c_2Etopology_2Etopology\ ty_2Erealax$

Assume the following.

$$True \quad (16)$$

Assume the following.

$$\forall A.27a.nonempty\ A.27a \Rightarrow (\forall V0x \in A.27a.((V0x = V0x) \Leftrightarrow True)) \quad (17)$$

Assume the following.

$$\forall A.27a.nonempty\ A.27a \Rightarrow (\forall V0x \in A.27a.(\forall V1y \in A.27a.((V0x = V1y) \Leftrightarrow (V1y = V0x)))) \quad (18)$$

Assume the following.

$$\begin{aligned} & \forall A.27a.nonempty\ A.27a \Rightarrow (\forall V0r \in (2^{A.27a}).(\forall V1p \in (2^{A.27a}).(\forall V2q \in (2^{A.27a}).(((ap\ (ap\ (c_2Epred_set_2EINTER\ A.27a)\ V1p)\ V2q) = (ap\ (ap\ (c_2Epred_set_2EINTER\ A.27a)\ V2q)\ V1p)) \wedge \\ & (((ap\ (ap\ (c_2Epred_set_2EINTER\ A.27a)\ V1p)\ V2q))\ V0r) = (ap\ (ap\ (c_2Epred_set_2EINTER\ A.27a)\ (ap\ (ap\ (c_2Epred_set_2EINTER\ A.27a)\ V1p)\ V2q))\ V0r)) \wedge (((ap\ (ap\ (c_2Epred_set_2EINTER\ A.27a)\ V1p)\ V2q))\ V0r) = (ap\ (ap\ (c_2Epred_set_2EINTER\ A.27a)\ (ap\ (ap\ (c_2Epred_set_2EINTER\ A.27a)\ V2q)\ V1p))\ V0r)) \wedge (((ap\ (ap\ (c_2Epred_set_2EINTER\ A.27a)\ V1p)\ V1p) = V1p) \wedge ((ap\ (ap\ (c_2Epred_set_2EINTER\ A.27a)\ (ap\ (ap\ (c_2Epred_set_2EINTER\ A.27a)\ V1p)\ V1p))\ V2q) = (ap\ (ap\ (c_2Epred_set_2EINTER\ A.27a)\ V1p)\ V2q))))))))) \quad (19) \end{aligned}$$

Assume the following.

$$\begin{aligned} & \forall A.27a.nonempty\ A.27a \Rightarrow (\forall V0top \in (ty_2Etopology_2Etopology\ A.27a).(\forall V1u \in (2^{A.27a}).(\forall V2s \in (2^{A.27a}).((p\ (ap\ (ap\ (c_2Etopology_2Eopen_in\ A.27a)\ (ap\ (ap\ (c_2Ereal_topology_2Esubtopology\ A.27a)\ V0top)\ V1u))\ V2s)) \Leftrightarrow (\exists V3t \in (2^{A.27a}).((p\ (ap\ (ap\ (c_2Etopology_2Eopen_in\ A.27a)\ V0top)\ V3t)) \wedge (V2s = (ap\ (ap\ (c_2Epred_set_2EINTER\ A.27a)\ V3t)\ V1u))))))))) \quad (20) \end{aligned}$$

Assume the following.

$$(\forall V0s \in (2^{ty_2Erealax_2Ereal}).((p\ (ap\ c_Ereal_topology_2EOpen\ V0s)) \Leftrightarrow (p\ (ap\ (ap\ (c_2Etopology_2Eopen_in\ ty_2Erealax_2Ereal)\ c_2Ereal_topology_2Eeuclidean)\ V0s)))) \quad (21)$$

Theorem 1

$$\begin{aligned} & (\forall V0s \in (2^{ty_2Erealax_2Ereal}).(\forall V1u \in (2^{ty_2Erealax_2Ereal}). \\ & ((p (ap (ap (c_2Etopology_2Eopen_in ty_2Erealax_2Ereal) (ap \\ & (ap (c_2Ereal_topology_2Esubtopology ty_2Erealax_2Ereal) \\ & c_2Ereal_topology_2Euclidean) V1u)) V0s)) \Leftrightarrow (\exists V2t \in (\\ & 2^{ty_2Erealax_2Ereal}).((p (ap c_2Ereal_topology_2EOpen V2t)) \wedge \\ & (V0s = (ap (ap (c_2Epred_set_2EINTER ty_2Erealax_2Ereal) V1u) \\ & V2t)))))) \end{aligned}$$