

# thm\_2Ereal\_\_topology\_2EOPEN\_\_POSITIVE\_\_ORTHANT (TMSY4YcTipaYo9mdv5E1R4DMuDvVXwd7RnX)

October 26, 2020

Let  $c\_2Enum\_2EZERO\_REP : \iota$  be given. Assume the following.

$$c\_2Enum\_2EZERO\_REP \in \omega \tag{1}$$

Let  $ty\_2Enum\_2Enum : \iota$  be given. Assume the following.

$$nonempty\ ty\_2Enum\_2Enum \tag{2}$$

Let  $c\_2Enum\_2EABS\_num : \iota$  be given. Assume the following.

$$c\_2Enum\_2EABS\_num \in (ty\_2Enum\_2Enum^{\omega}) \tag{3}$$

**Definition 1** We define  $c\_2Emin\_2E\_3D$  to be  $\lambda A.\lambda x \in A.\lambda y \in A.inj\_o (x = y)$  of type  $\iota \Rightarrow \iota$ .

**Definition 2** We define  $c\_2Enum\_2E0$  to be  $(ap\ c\_2Enum\_2EABS\_num\ c\_2Enum\_2EZERO\_REP)$ .

Let  $ty\_2Erealx\_2Ereal : \iota$  be given. Assume the following.

$$nonempty\ ty\_2Erealx\_2Ereal \tag{4}$$

Let  $c\_2Ereal\_2Ereal\_of\_num : \iota$  be given. Assume the following.

$$c\_2Ereal\_2Ereal\_of\_num \in (ty\_2Erealx\_2Ereal^{ty\_2Enum\_2Enum}) \tag{5}$$

**Definition 3** We define  $c\_2Ebool\_2ET$  to be  $(ap\ (ap\ (c\_2Emin\_2E\_3D\ (2^2))\ (\lambda V0x \in 2.V0x))\ (\lambda V1x \in 2.V1x))$

**Definition 4** We define  $c\_2Ebool\_2E\_21$  to be  $\lambda A.\lambda a : \iota.(\lambda V0P \in (2^{A-27a}).(ap\ (ap\ (c\_2Emin\_2E\_3D\ (2^{A-27a})))$

**Definition 5** We define  $c\_2Ebool\_2EF$  to be  $(ap\ (c\_2Ebool\_2E\_21\ 2)\ (\lambda V0t \in 2.V0t))$ .

**Definition 6** We define  $c\_2Emin\_2E\_3D\_3D\_3E$  to be  $\lambda P \in 2.\lambda Q \in 2.inj\_o (p\ P \Rightarrow p\ Q)$  of type  $\iota$ .



**Definition 14** We define `c_2Ebool_2E_3F` to be  $\lambda A_27a : \iota. (\lambda V0P \in (2^{A_27a}). (ap\ V0P\ (ap\ (c_2Emin_2E_40$

**Definition 15** We define `c_2Ereal_topology_2EOpen` to be  $\lambda V0s \in (2^{ty_2Erealax_2Ereal}). (ap\ (c_2Ebool_2E_2$

Assume the following.

$$True \tag{13}$$

Assume the following.

$$\begin{aligned} & (\forall V0t \in 2. (((True \Rightarrow (p\ V0t)) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \Rightarrow True) \Leftrightarrow \\ & True) \wedge (((False \Rightarrow (p\ V0t)) \Leftrightarrow True) \wedge (((p\ V0t) \Rightarrow (p\ V0t)) \Leftrightarrow True) \wedge (( \\ & (p\ V0t) \Rightarrow False) \Leftrightarrow (\neg (p\ V0t)))))) \end{aligned} \tag{14}$$

Assume the following.

$$\begin{aligned} & (\forall V0a \in ty_2Erealax_2Ereal. (p\ (ap\ c_2Ereal\_topology_2EOpen \\ & (ap\ (c_2Epred\_set_2EGSPEC\ ty_2Erealax_2Ereal\ ty_2Erealax_2Ereal) \\ & (\lambda V1x \in ty_2Erealax_2Ereal. (ap\ (ap\ (c_2Epair_2E_2C\ ty_2Erealax_2Ereal \\ & 2)\ V1x)\ (ap\ (ap\ c_2Erealax_2Ereal\_lt\ V0a)\ V1x)))))) \end{aligned} \tag{15}$$

**Theorem 1**

$$\begin{aligned} & (p\ (ap\ c_2Ereal\_topology_2EOpen\ (ap\ (c_2Epred\_set_2EGSPEC \\ & ty_2Erealax_2Ereal\ ty_2Erealax_2Ereal)\ (\lambda V0x \in ty_2Erealax_2Ereal. \\ & (ap\ (ap\ (c_2Epair_2E_2C\ ty_2Erealax_2Ereal\ 2)\ V0x)\ (ap\ (ap\ c_2Erealax_2Ereal\_lt \\ & (ap\ c_2Ereal_2Ereal\_of\_num\ c_2Enum_2E0)\ V0x)))))) \end{aligned}$$