

thm_2Ereal__topology_2EREAL__EQ__SQUARE__ABS
 (TMKiGgQd-
 pQF6iFSnM5WBC6ecAE1xjHBVGeF)

October 26, 2020

Definition 1 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define c_2Ebool_2ET to be $(ap (ap (c_2Emin_2E_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

Definition 3 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p P \Rightarrow p Q)$ of type ι .

Definition 4 We define $c_2Ebool_2E_21$ to be $\lambda A.\lambda a : \iota.(\lambda V0P \in (2^{A-27a}).(ap (ap (c_2Emin_2E_3D (2^{A-27a}))$

Definition 5 We define $c_2Ebool_2E_2F_5C$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21 2) (\lambda V2t \in 2.V2t)))$

Let $c_2Enum_2EZERO_REP : \iota$ be given. Assume the following.

$$c_2Enum_2EZERO_REP \in \omega \tag{1}$$

Let $ty_2Enum_2Enum : \iota$ be given. Assume the following.

$$nonempty\ ty_2Enum_2Enum \tag{2}$$

Let $c_2Enum_2EABS_num : \iota$ be given. Assume the following.

$$c_2Enum_2EABS_num \in (ty_2Enum_2Enum^{\omega}) \tag{3}$$

Definition 6 We define c_2Enum_2E0 to be $(ap\ c_2Enum_2EABS_num\ c_2Enum_2EZERO_REP)$.

Definition 7 We define $c_2Earithmic_2EZERO$ to be c_2Enum_2E0 .

Let $c_2Enum_2EREP_num : \iota$ be given. Assume the following.

$$c_2Enum_2EREP_num \in (\omega^{ty_2Enum_2Enum}) \tag{4}$$

Let $c_2Enum_2ESUC_REP : \iota$ be given. Assume the following.

$$c_2Enum_2ESUC_REP \in (\omega^{\omega}) \tag{5}$$

Definition 8 We define c_2Enum_2ESUC to be $\lambda V0m \in ty_2Enum_2Enum.(ap\ c_2Enum_2EABS_num$

Let $c_2Earithmetic_2E_2B : \iota$ be given. Assume the following.

$$c_2Earithmetic_2E_2B \in ((ty_2Enum_2Enum^{ty_2Enum_2Enum})^{ty_2Enum_2Enum}) \quad (6)$$

Definition 9 We define $c_2Earithmetic_2EBIT2$ to be $\lambda V0n \in ty_2Enum_2Enum.(ap\ (ap\ c_2Earithmetic_2E_2B$

Definition 10 We define $c_2Earithmetic_2ENUMERAL$ to be $\lambda V0x \in ty_2Enum_2Enum.V0x$.

Let $ty_2Erealax_2Ereal : \iota$ be given. Assume the following.

$$nonempty\ ty_2Erealax_2Ereal \quad (7)$$

Let $c_2Ereal_2Epow : \iota$ be given. Assume the following.

$$c_2Ereal_2Epow \in ((ty_2Erealax_2Ereal^{ty_2Enum_2Enum})^{ty_2Erealax_2Ereal}) \quad (8)$$

Let $ty_2Ehreal_2Ehreal : \iota$ be given. Assume the following.

$$nonempty\ ty_2Ehreal_2Ehreal \quad (9)$$

Let $ty_2Epair_2Eprod : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A0.nonempty\ A0 \Rightarrow \forall A1.nonempty\ A1 \Rightarrow nonempty\ (ty_2Epair_2Eprod\ A0\ A1) \quad (10)$$

Let $c_2Erealax_2Ereal_2REP_2CLASS : \iota$ be given. Assume the following.

$$c_2Erealax_2Ereal_2REP_2CLASS \in ((2^{(ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal\ ty_2Ehreal_2Ehreal)})^{ty_2Erealax_2Ereal}) \quad (11)$$

Definition 11 We define $c_2Emin_2E_40$ to be $\lambda A.\lambda P \in 2^A$.if $(\exists x \in A.p\ (ap\ P\ x))$ then $(the\ (\lambda x.x \in A \wedge P\ x))$ of type $\iota \Rightarrow \iota$.

Definition 12 We define $c_2Erealax_2Ereal_2REP$ to be $\lambda V0a \in ty_2Erealax_2Ereal.(ap\ (c_2Emin_2E_40\ (the\ (\lambda x.x \in A \wedge P\ x))))$

Let $c_2Erealax_2Etrealm_neg : \iota$ be given. Assume the following.

$$c_2Erealax_2Etrealm_neg \in ((ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal\ ty_2Ehreal_2Ehreal)^{ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal\ ty_2Ehreal_2Ehreal}) \quad (12)$$

Let $c_2Erealax_2Etrealm_eq : \iota$ be given. Assume the following.

$$c_2Erealax_2Etrealm_eq \in ((2^{(ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal\ ty_2Ehreal_2Ehreal)})^{ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal\ ty_2Ehreal_2Ehreal}) \quad (13)$$

Let $c_2Erealax_2Ereal_2ABS_2CLASS : \iota$ be given. Assume the following.

$$c_2Erealax_2Ereal_2ABS_2CLASS \in (ty_2Erealax_2Ereal^{(2^{(ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal\ ty_2Ehreal_2Ehreal)})}) \quad (14)$$

Definition 13 We define $c_2Erealax_2Ereal_ABS$ to be $\lambda V0r \in (ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal\ ty_2Ehreal\ ty_2Ehreal)$

Definition 14 We define $c_2Erealax_2Ereal_neg$ to be $\lambda V0T1 \in ty_2Erealax_2Ereal.(ap\ c_2Erealax_2Ereal)$

Let $c_2Ereal_2Ereal_of_num : \iota$ be given. Assume the following.

$$c_2Ereal_2Ereal_of_num \in (ty_2Erealax_2Ereal^{ty_2Eenum_2Eenum}) \quad (15)$$

Let $c_2Erealax_2Etrealt_lt : \iota$ be given. Assume the following.

$$c_2Erealax_2Etrealt_lt \in ((2^{(ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal\ ty_2Ehreal_2Ehreal)})(ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal)) \quad (16)$$

Definition 15 We define $c_2Erealax_2Ereal_lt$ to be $\lambda V0T1 \in ty_2Erealax_2Ereal.\lambda V1T2 \in ty_2Erealax_2Ereal$

Definition 16 We define c_2Ebool_2Eef to be $(ap\ (c_2Ebool_2E21\ 2)\ (\lambda V0t \in 2.V0t))$.

Definition 17 We define c_2Ebool_2E7E to be $(\lambda V0t \in 2.(ap\ (ap\ c_2Emin_2E3D_3D_3E\ V0t)\ c_2Ebool_2E7E))$

Definition 18 We define $c_2Ereal_2Ereal_lte$ to be $\lambda V0x \in ty_2Erealax_2Ereal.\lambda V1y \in ty_2Erealax_2Ereal$

Definition 19 We define c_2Ebool_2ECOND to be $\lambda A_27a : \iota.(\lambda V0t \in 2.(\lambda V1t1 \in A_27a.(\lambda V2t2 \in A_27a.(ap\ (ap\ (ap\ c_2Ebool_2ECOND\ V0t)\ V1t1)\ V2t2)\ V0t)))$

Definition 20 We define c_2Ereal_2Eabs to be $\lambda V0x \in ty_2Erealax_2Ereal.(ap\ (ap\ (ap\ (c_2Ebool_2ECOND\ V0x)\ V0x)\ V0x)\ V0x)$

Assume the following.

$$True \quad (17)$$

Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow (\forall V0t \in 2.((\forall V1x \in A_27a.(p\ V0t) \Leftrightarrow (p\ V1x))) \Leftrightarrow (p\ V0t))) \quad (18)$$

Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow (\forall V0x \in A_27a.((V0x = V0x) \Leftrightarrow True)) \quad (19)$$

Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow (\forall V0x \in A_27a.(\forall V1y \in A_27a.((V0x = V1y) \Leftrightarrow (V1y = V0x)))) \quad (20)$$

Assume the following.

$$\begin{aligned} & (\forall V0x \in ty_2Erealax_2Ereal.(\forall V1y \in ty_2Erealax_2Ereal. \\ & (((p\ (ap\ (ap\ c_2Ereal_2Ereal_lte\ V0x)\ V1y)) \wedge (p\ (ap\ (ap\ c_2Ereal_2Ereal_lte\ V1y)\ V0x))) \Leftrightarrow (V0x = V1y)))) \end{aligned} \quad (21)$$

Assume the following.

$$\begin{aligned}
& (\forall V0x \in ty_2Erealax_2Ereal. (\forall V1y \in ty_2Erealax_2Ereal. \\
& ((p (ap (ap c_2Ereal_2Ereal_lte (ap c_2Ereal_2Eabs V0x)) (ap c_2Ereal_2Eabs \\
& V1y))) \Leftrightarrow (p (ap (ap c_2Ereal_2Ereal_lte (ap (ap c_2Ereal_2Epow \\
& V0x) (ap c_2Earithmetic_2ENUMERAL (ap c_2Earithmetic_2EBIT2 \\
& c_2Earithmetic_2EZERO)))) (ap (ap c_2Ereal_2Epow V1y) (ap c_2Earithmetic_2ENUMERAL \\
& (ap c_2Earithmetic_2EBIT2 c_2Earithmetic_2EZERO))))))))) \\
& \hspace{15em} (22)
\end{aligned}$$

Theorem 1

$$\begin{aligned}
& (\forall V0x \in ty_2Erealax_2Ereal. (\forall V1y \in ty_2Erealax_2Ereal. \\
& (((ap c_2Ereal_2Eabs V0x) = (ap c_2Ereal_2Eabs V1y)) \Leftrightarrow ((ap (ap c_2Ereal_2Epow \\
& V0x) (ap c_2Earithmetic_2ENUMERAL (ap c_2Earithmetic_2EBIT2 \\
& c_2Earithmetic_2EZERO))) = (ap (ap c_2Ereal_2Epow V1y) (ap c_2Earithmetic_2ENUMERAL \\
& (ap c_2Earithmetic_2EBIT2 c_2Earithmetic_2EZERO)))))))))
\end{aligned}$$