

# thm\_2Ereal\_topology\_2ESERIES\_NEG (TMHtRADs2stizAbmNRjZSkeFj2L8q9wi2dd)

October 26, 2020

**Definition 1** We define  $c\_2Emin\_2E\_3D$  to be  $\lambda A.\lambda x \in A.\lambda y \in A.inj\_o (x = y)$  of type  $\iota \Rightarrow \iota$ .

**Definition 2** We define  $c\_2Ebool\_2E\_7E$  to be  $(ap (ap (c\_2Emin\_2E\_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

**Definition 3** We define  $c\_2Ebool\_2E\_21$  to be  $\lambda A.\lambda P \in 2^A.(ap (ap (c\_2Emin\_2E\_3D (2^{A-27a})) (\lambda V0P \in 2.V0P)) (\lambda V1P \in 2.V1P))$

**Definition 4** We define  $c\_2Ebool\_2E\_2F$  to be  $(ap (c\_2Ebool\_2E\_21 2) (\lambda V0t \in 2.V0t))$ .

**Definition 5** We define  $c\_2Emin\_2E\_3D\_3D\_3E$  to be  $\lambda P \in 2.\lambda Q \in 2.inj\_o (p P \Rightarrow p Q)$  of type  $\iota$ .

**Definition 6** We define  $c\_2Ebool\_2E\_7E$  to be  $(\lambda V0t \in 2.(ap (ap c\_2Emin\_2E\_3D\_3D\_3E V0t) c\_2Ebool\_2E\_2F))$

**Definition 7** We define  $c\_2Ebool\_2E\_2F\_5C$  to be  $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c\_2Ebool\_2E\_21 2) (\lambda V2t \in 2.V2t)) (\lambda V3t \in 2.V3t)))$

Let  $ty\_2Ehreal\_2Ehreal : \iota$  be given. Assume the following.

$$nonempty\ ty\_2Ehreal\_2Ehreal \tag{1}$$

Let  $ty\_2Epair\_2Eprod : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A0.nonempty\ A0 \Rightarrow \forall A1.nonempty\ A1 \Rightarrow nonempty\ (ty\_2Epair\_2Eprod\ A0\ A1) \tag{2}$$

Let  $ty\_2Erealax\_2Ereal : \iota$  be given. Assume the following.

$$nonempty\ ty\_2Erealax\_2Ereal \tag{3}$$

Let  $c\_2Erealax\_2Ereal\_REP\_CLASS : \iota$  be given. Assume the following.

$$c\_2Erealax\_2Ereal\_REP\_CLASS \in ((2^{(ty\_2Epair\_2Eprod\ ty\_2Ehreal\_2Ehreal\ ty\_2Ehreal\_2Ehreal)})\ ty\_2Erealax\_2Ereal) \tag{4}$$

**Definition 8** We define  $c\_2Emin\_2E\_40$  to be  $\lambda A.\lambda P \in 2^A.if (\exists x \in A.p (ap P x)) \mathbf{then} (the (\lambda x.x \in A \wedge p x))$  of type  $\iota \Rightarrow \iota$ .

**Definition 9** We define  $c\_2Erealax\_2Ereal\_REP$  to be  $\lambda V0a \in ty\_2Erealax\_2Ereal.(ap (c\_2Emin\_2E40 (ty$

Let  $c\_2Erealax\_2Etrealm\_neg : \iota$  be given. Assume the following.

$$c\_2Erealax\_2Etrealm\_neg \in ((ty\_2Epair\_2Eprod ty\_2Ehreal\_2Ehreal ty\_2Ehreal\_2Ehreal)^{(ty\_2Epair\_2Eprod ty\_2Ehreal\_2Ehreal ty\_2Ehreal\_2Ehreal)}) \quad (5)$$

Let  $c\_2Erealax\_2Etrealm\_eq : \iota$  be given. Assume the following.

$$c\_2Erealax\_2Etrealm\_eq \in ((2^{(ty\_2Epair\_2Eprod ty\_2Ehreal\_2Ehreal ty\_2Ehreal\_2Ehreal)})^{(ty\_2Epair\_2Eprod ty\_2Ehreal\_2Ehreal)}) \quad (6)$$

Let  $c\_2Erealax\_2Ereal\_ABS\_CLASS : \iota$  be given. Assume the following.

$$c\_2Erealax\_2Ereal\_ABS\_CLASS \in (ty\_2Erealax\_2Ereal)^{(2^{(ty\_2Epair\_2Eprod ty\_2Ehreal\_2Ehreal ty\_2Ehreal\_2Ehreal)})} \quad (7)$$

**Definition 10** We define  $c\_2Erealax\_2Ereal\_ABS$  to be  $\lambda V0r \in (ty\_2Epair\_2Eprod ty\_2Ehreal\_2Ehreal ty$

**Definition 11** We define  $c\_2Erealax\_2Ereal\_neg$  to be  $\lambda V0T1 \in ty\_2Erealax\_2Ereal.(ap c\_2Erealax\_2Ereal$

Let  $ty\_2Ereal\_topology\_2Enet : \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A0.nonempty A0 \Rightarrow nonempty (ty\_2Ereal\_topology\_2Enet A0) \quad (8)$$

Let  $ty\_2Eenum\_2Eenum : \iota$  be given. Assume the following.

$$nonempty ty\_2Eenum\_2Eenum \quad (9)$$

Let  $c\_2Eenum\_2EREP\_num : \iota$  be given. Assume the following.

$$c\_2Eenum\_2EREP\_num \in (\omega^{ty\_2Eenum\_2Eenum}) \quad (10)$$

Let  $c\_2Eenum\_2ESUC\_REP : \iota$  be given. Assume the following.

$$c\_2Eenum\_2ESUC\_REP \in (\omega^{\omega}) \quad (11)$$

Let  $c\_2Eenum\_2EABS\_num : \iota$  be given. Assume the following.

$$c\_2Eenum\_2EABS\_num \in (ty\_2Eenum\_2Eenum^{\omega}) \quad (12)$$

**Definition 12** We define  $c\_2Eenum\_2ESUC$  to be  $\lambda V0m \in ty\_2Eenum\_2Eenum.(ap c\_2Eenum\_2EABS\_num$

**Definition 13** We define  $c\_2Ebool\_2E3F$  to be  $\lambda A\_27a : \iota.(\lambda V0P \in (2^{A\_27a}).(ap V0P (ap (c\_2Emin\_2E40$

**Definition 14** We define  $c\_2Eprim\_rec\_2E3C$  to be  $\lambda V0m \in ty\_2Eenum\_2Eenum.\lambda V1n \in ty\_2Eenum\_2Eenum$

**Definition 15** We define  $c\_2Earithmetic\_2E3E$  to be  $\lambda V0m \in ty\_2Eenum\_2Eenum.\lambda V1n \in ty\_2Eenum\_2Eenum$

**Definition 16** We define  $c\_2Ebool\_2E5C\_2F$  to be  $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c\_2Ebool\_2E21 2) (\lambda V2t \in$

**Definition 17** We define  $c\_2Earithmetic\_2E3E\_3D$  to be  $\lambda V0m \in ty\_2Eenum\_2Eenum.\lambda V1n \in ty\_2Eenum\_2Eenum$



**Definition 30** We define  $c\_2Epred\_set\_2EEMPTY$  to be  $\lambda A\_27a : \iota. (\lambda V0x \in A\_27a. c\_2Ebool\_2EF)$ .

**Definition 31** We define  $c\_2Epred\_set\_2EFINITE$  to be  $\lambda A\_27a : \iota. \lambda V0s \in (2^{A\_27a}). (ap (c\_2Ebool\_2E21$

**Definition 32** We define  $c\_2Eiterate\_2EITSET$  to be  $\lambda A\_27a : \iota. \lambda A\_27b : \iota. \lambda V0f \in ((A\_27a^{A\_27a})^{A\_27b}). \lambda V$

**Definition 33** We define  $c\_2Eiterate\_2Eiterate$  to be  $\lambda A\_27a : \iota. \lambda A\_27b : \iota. \lambda V0op \in ((A\_27b^{A\_27b})^{A\_27b}). \lambda V$

**Definition 34** We define  $c\_2Eiterate\_2ESum$  to be  $\lambda A\_27a : \iota. (ap (c\_2Eiterate\_2Eiterate A\_27a ty\_2Erealax$

Let  $c\_2Ereal\_topology\_2EDist : \iota$  be given. Assume the following.

$$c\_2Ereal\_topology\_2EDist \in (ty\_2Erealax\_2Ereal^{(ty\_2Epair\_2Eprod ty\_2Erealax\_2Ereal ty\_2Erealax\_2Ereal)}) \quad (18)$$

Let  $c\_2Erealax\_2Etrealt\_lt : \iota$  be given. Assume the following.

$$c\_2Erealax\_2Etrealt\_lt \in ((2^{(ty\_2Epair\_2Eprod ty\_2Ehreal\_2Ehreal ty\_2Ehreal\_2Ehreal)})^{(ty\_2Epair\_2Eprod ty\_2Ehreal\_2Ehreal)}) \quad (19)$$

**Definition 35** We define  $c\_2Erealax\_2Ereal\_lt$  to be  $\lambda V0T1 \in ty\_2Erealax\_2Ereal. \lambda V1T2 \in ty\_2Erealax$

Let  $c\_2Ereal\_topology\_2Eenetord : \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A\_27a. nonempty A\_27a \Rightarrow c\_2Ereal\_topology\_2Eenetord A\_27a \in (((2^{A\_27a})^{A\_27a})^{(ty\_2Ereal\_topology\_2Eenet A\_27a)}) \quad (20)$$

**Definition 36** We define  $c\_2Ereal\_topology\_2Etrivial\_limit$  to be  $\lambda A\_27a : \iota. \lambda V0net \in (ty\_2Ereal\_topology$

**Definition 37** We define  $c\_2Ereal\_topology\_2Eeventually$  to be  $\lambda A\_27a : \iota. \lambda V0p \in (2^{A\_27a}). \lambda V1net \in (ty\_2$

Let  $c\_2Ereal\_2Ereal\_of\_num : \iota$  be given. Assume the following.

$$c\_2Ereal\_2Ereal\_of\_num \in (ty\_2Erealax\_2Ereal^{ty\_2Eenum\_2Eenum}) \quad (21)$$

**Definition 38** We define  $c\_2Ereal\_topology\_2E2D\_2D\_3E$  to be  $\lambda A\_27a : \iota. \lambda V0f \in (ty\_2Erealax\_2Ereal^{A$

**Definition 39** We define  $c\_2Ereal\_topology\_2ESums$  to be  $\lambda V0f \in (ty\_2Erealax\_2Ereal^{ty\_2Eenum\_2Eenum}). \lambda$

Assume the following.

$$True \quad (22)$$

Assume the following.

$$\forall A\_27a. nonempty A\_27a \Rightarrow (\forall V0t \in 2. ((\forall V1x \in A\_27a. (p V0t)) \Leftrightarrow (p V0t))) \quad (23)$$

Assume the following.

$$\begin{aligned} & (\forall V0t \in 2. (((True \Rightarrow (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \Rightarrow True) \Leftrightarrow \\ & True) \wedge (((False \Rightarrow (p V0t)) \Leftrightarrow True) \wedge (((p V0t) \Rightarrow (p V0t)) \Leftrightarrow True) \wedge (( \\ & (p V0t) \Rightarrow False) \Leftrightarrow (\neg (p V0t)))))) \end{aligned} \quad (24)$$

Assume the following.

$$\forall A\_27a.nonempty\ A\_27a \Rightarrow (\forall V0x \in A\_27a. (\forall V1y \in A\_27a. ((V0x = V1y) \Leftrightarrow (V1y = V0x)))) \quad (25)$$

Assume the following.

$$(\forall V0t \in 2. (((True \Leftrightarrow (p\ V0t)) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \Leftrightarrow True) \Leftrightarrow (p\ V0t)) \wedge (((False \Leftrightarrow (p\ V0t)) \Leftrightarrow \neg(p\ V0t)) \wedge (((p\ V0t) \Leftrightarrow False) \Leftrightarrow \neg(p\ V0t)))))) \quad (26)$$

Assume the following.

$$(\forall V0t1 \in 2. (\forall V1t2 \in 2. (\forall V2t3 \in 2. (((p\ V0t1) \Rightarrow ((p\ V1t2) \Rightarrow (p\ V2t3))) \Leftrightarrow (((p\ V0t1) \wedge (p\ V1t2)) \Rightarrow (p\ V2t3)))))) \quad (27)$$

Assume the following.

$$(\forall V0x \in 2. (\forall V1x\_27 \in 2. (\forall V2y \in 2. (\forall V3y\_27 \in 2. (((p\ V0x) \Leftrightarrow (p\ V1x\_27)) \wedge ((p\ V1x\_27) \Rightarrow ((p\ V2y) \Leftrightarrow (p\ V3y\_27)))))) \Rightarrow (((p\ V0x) \Rightarrow (p\ V2y)) \Leftrightarrow ((p\ V1x\_27) \Rightarrow (p\ V3y\_27)))))) \quad (28)$$

Assume the following.

$$\forall A\_27a.nonempty\ A\_27a \Rightarrow (\forall V0f \in (ty\_2Erealax\_2Ereal^{A\_27a}). (\forall V1s \in (2^{A\_27a}). ((ap\ (ap\ (c\_2Eiterate\_2ESum\ A\_27a)\ V1s)\ (\lambda V2x \in A\_27a. (ap\ c\_2Erealax\_2Ereal\_neg\ (ap\ V0f\ V2x)))))) = (ap\ c\_2Erealax\_2Ereal\_neg\ (ap\ (ap\ (c\_2Eiterate\_2ESum\ A\_27a)\ V1s)\ V0f)))))) \quad (29)$$

Assume the following.

$$\forall A\_27a.nonempty\ A\_27a \Rightarrow (\forall V0net \in (ty\_2Ereal\_topology\_2Enet\ A\_27a). (\forall V1f \in (ty\_2Erealax\_2Ereal^{A\_27a}). (\forall V2l \in ty\_2Erealax\_2Ereal. ((p\ (ap\ (ap\ (ap\ (c\_2Ereal\_topology\_2E\_2D\_2D\_3E\ A\_27a)\ V1f)\ V2l)\ V0net)) \Rightarrow (p\ (ap\ (ap\ (ap\ (c\_2Ereal\_topology\_2E\_2D\_2D\_3E\ A\_27a)\ (\lambda V3x \in A\_27a. (ap\ c\_2Erealax\_2Ereal\_neg\ (ap\ V1f\ V3x))))\ (ap\ c\_2Erealax\_2Ereal\_neg\ V2l))\ V0net)))))) \quad (30)$$

### Theorem 1

$$(\forall V0x \in (ty\_2Erealax\_2Ereal^{ty\_2Enum\_2Enum}). (\forall V1x0 \in ty\_2Erealax\_2Ereal. (\forall V2s \in (2^{ty\_2Enum\_2Enum}). ((p\ (ap\ (ap\ (ap\ c\_2Ereal\_topology\_2Esums\ V0x)\ V1x0)\ V2s)) \Rightarrow (p\ (ap\ (ap\ (ap\ c\_2Ereal\_topology\_2Esums\ (\lambda V3n \in ty\_2Enum\_2Enum. (ap\ c\_2Erealax\_2Ereal\_neg\ (ap\ V0x\ V3n))))\ (ap\ c\_2Erealax\_2Ereal\_neg\ V1x0))\ V2s))))))$$