

thm_2Ereal_topology_2ESETDIST_EMPTY
(TMLQwZTFEFLECWnji-
iZRLu42Q3txQ8nxiGA)

October 26, 2020

Definition 1 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define c_2Ebool_2ET to be $(ap (ap (c_2Emin_2E_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

Definition 3 We define $c_2Ebool_2E_21$ to be $\lambda A_27a : \iota.(\lambda V0P \in (2^{A_27a}).(ap (ap (c_2Emin_2E_3D (2^{A_27a}))$

Definition 4 We define c_2Ebool_2EF to be $(ap (c_2Ebool_2E_21 2) (\lambda V0t \in 2.V0t))$.

Definition 5 We define c_2Ebool_2EIN to be $\lambda A_27a : \iota.(\lambda V0x \in A_27a.(\lambda V1f \in (2^{A_27a}).(ap V1f V0x)))$

Definition 6 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p P \Rightarrow p Q)$ of type ι .

Definition 7 We define $c_2Ebool_2E_2F_5C$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21 2) (\lambda V2t \in 2.V2t)))$

Let $ty_2Erealax_2Ereal : \iota$ be given. Assume the following.

$$nonempty\ ty_2Erealax_2Ereal \tag{1}$$

Let $ty_2Epair_2Eprod : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A0.nonempty\ A0 \Rightarrow \forall A1.nonempty\ A1 \Rightarrow nonempty\ (ty_2Epair_2Eprod\ A0\ A1) \tag{2}$$

Let $c_2Ereal_topology_2EDist : \iota$ be given. Assume the following.

$$c_2Ereal_topology_2EDist \in (ty_2Erealax_2Ereal^{(ty_2Epair_2Eprod\ ty_2Erealax_2Ereal\ ty_2Erealax_2Ereal)}) \tag{3}$$

Let $c_2Epair_2ESND : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow \forall A_27b.nonempty\ A_27b \Rightarrow c_2Epair_2ESND\ A_27a\ A_27b \in (A_27b^{(ty_2Epair_2Eprod\ A_27a\ A_27b)}) \tag{4}$$

Let $c_2Epair_2EFST : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned} \forall A_27a.nonempty\ A_27a \Rightarrow \forall A_27b.nonempty\ A_27b \Rightarrow c_2Epair_2EFST \\ A_27a\ A_27b \in (A_27a^{(ty_2Epair_2Eprod\ A_27a\ A_27b)}) \end{aligned} \quad (5)$$

Definition 8 We define $c_2Epair_2EUNCURRY$ to be $\lambda A_27a : \iota.\lambda A_27b : \iota.\lambda A_27c : \iota.\lambda V0f \in ((A_27c^{A_27b})$

Let $c_2Epred_set_2EGSPEC : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned} \forall A_27a.nonempty\ A_27a \Rightarrow \forall A_27b.nonempty\ A_27b \Rightarrow c_2Epred_set_2EGSPEC \\ A_27a\ A_27b \in ((2^{A_27a})^{(ty_2Epair_2Eprod\ A_27a\ 2)^{A_27b}}) \end{aligned} \quad (6)$$

Let $ty_2Ehreal_2Ehreal : \iota$ be given. Assume the following.

$$nonempty\ ty_2Ehreal_2Ehreal \quad (7)$$

Let $c_2Erealax_2Ereal_REP_CLASS : \iota$ be given. Assume the following.

$$c_2Erealax_2Ereal_REP_CLASS \in ((2^{(ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal\ ty_2Ehreal_2Ehreal)})^{ty_2Erealax_2Ereal_REP_CLASS}) \quad (8)$$

Definition 9 We define $c_2Emin_2E.40$ to be $\lambda A.\lambda P \in 2^A$. **if** $(\exists x \in A.p\ (ap\ P\ x))$ **then** (the $(\lambda x.x \in A \wedge p\ x)$ of type $\iota \Rightarrow \iota$).

Definition 10 We define $c_2Erealax_2Ereal_REP$ to be $\lambda V0a \in ty_2Erealax_2Ereal.(ap\ (c_2Emin_2E.40\ (ap\ P\ x)))$

Let $c_2Erealax_2Etreall_lt : \iota$ be given. Assume the following.

$$c_2Erealax_2Etreall_lt \in ((2^{(ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal\ ty_2Ehreal_2Ehreal)})^{(ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal)}) \quad (9)$$

Definition 11 We define $c_2Erealax_2Ereal_lt$ to be $\lambda V0T1 \in ty_2Erealax_2Ereal.\lambda V1T2 \in ty_2Erealax_2Ereal$

Definition 12 We define $c_2Ebool_2E.7E$ to be $(\lambda V0t \in 2.(ap\ (ap\ c_2Emin_2E.3D_3D_3E\ V0t)\ c_2Ebool_2E.7E))$

Definition 13 We define $c_2Ereal_2Ereal_lte$ to be $\lambda V0x \in ty_2Erealax_2Ereal.\lambda V1y \in ty_2Erealax_2Ereal$

Definition 14 We define $c_2Eiterate_2Einf$ to be $\lambda V0s \in (2^{ty_2Erealax_2Ereal}).(ap\ (c_2Emin_2E.40\ ty_2Erealax_2Ereal))$

Let $c_2Enum_2EZERO_REP : \iota$ be given. Assume the following.

$$c_2Enum_2EZERO_REP \in \omega \quad (10)$$

Let $ty_2Enum_2Enum : \iota$ be given. Assume the following.

$$nonempty\ ty_2Enum_2Enum \quad (11)$$

Let $c_2Enum_2EABS_num : \iota$ be given. Assume the following.

$$c_2Enum_2EABS_num \in (ty_2Enum_2Enum^{\omega}) \quad (12)$$

Assume the following.

$$\begin{aligned} & \forall A.27a.nonempty\ A.27a \Rightarrow (\forall V0t1 \in A.27a. (\forall V1t2 \in \\ & A.27a. (((ap\ (ap\ (ap\ (c.2Ebool.2ECOND\ A.27a)\ c.2Ebool.2ET)\ V0t1) \\ & V1t2) = V0t1) \wedge ((ap\ (ap\ (ap\ (c.2Ebool.2ECOND\ A.27a)\ c.2Ebool.2EF) \\ & V0t1)\ V1t2) = V1t2)))))) \end{aligned} \quad (21)$$

Assume the following.

$$\begin{aligned} & (\forall V0s \in (2^{ty.2Erealx.2Ereal}). (\forall V1t \in (2^{ty.2Erealx.2Ereal}). \\ & ((ap\ c.2Ereal_topology.2Esetdist\ (ap\ (ap\ (c.2Epair.2E.2C\ (2^{ty.2Erealx.2Ereal}) \\ & (2^{ty.2Erealx.2Ereal}))\ V0s)\ V1t)) = (ap\ (ap\ (ap\ (c.2Ebool.2ECOND \\ & ty.2Erealx.2Ereal)\ (ap\ (ap\ c.2Ebool.2E.5C.2F\ (ap\ (ap\ (c.2Emin.2E.3D \\ & (2^{ty.2Erealx.2Ereal}))\ V0s)\ (c.2Epred_set.2EEMPTY\ ty.2Erealx.2Ereal)))) \\ & (ap\ (ap\ (c.2Emin.2E.3D\ (2^{ty.2Erealx.2Ereal}))\ V1t)\ (c.2Epred_set.2EEMPTY \\ & ty.2Erealx.2Ereal))))\ (ap\ c.2Ereal.2Ereal_of_num\ c.2Enum.2E0)) \\ & (ap\ c.2Eiterate.2Einf\ (ap\ (c.2Epred_set.2EGSPEC\ ty.2Erealx.2Ereal \\ & (ty.2Epair.2Eprod\ ty.2Erealx.2Ereal\ ty.2Erealx.2Ereal)) \\ & (ap\ (c.2Epair.2EUNCURRY\ ty.2Erealx.2Ereal\ ty.2Erealx.2Ereal \\ & (ty.2Epair.2Eprod\ ty.2Erealx.2Ereal\ 2))\ (\lambda V2x \in ty.2Erealx.2Ereal. \\ & (\lambda V3y \in ty.2Erealx.2Ereal. (ap\ (ap\ (c.2Epair.2E.2C\ ty.2Erealx.2Ereal \\ & 2)\ (ap\ c.2Ereal_topology.2EDist\ (ap\ (ap\ (c.2Epair.2E.2C\ ty.2Erealx.2Ereal \\ & ty.2Erealx.2Ereal)\ V2x)\ V3y))))\ (ap\ (ap\ c.2Ebool.2E.2F.5C\ (ap \\ & (ap\ (c.2Ebool.2EIN\ ty.2Erealx.2Ereal)\ V2x)\ V0s))\ (ap\ (ap\ (c.2Ebool.2EIN \\ & ty.2Erealx.2Ereal)\ V3y)\ V1t)))))))))) \end{aligned} \quad (22)$$

Theorem 1

$$\begin{aligned} & ((\forall V0t \in (2^{ty.2Erealx.2Ereal}). ((ap\ c.2Ereal_topology.2Esetdist \\ & (ap\ (ap\ (c.2Epair.2E.2C\ (2^{ty.2Erealx.2Ereal})\ (2^{ty.2Erealx.2Ereal})) \\ & (c.2Epred_set.2EEMPTY\ ty.2Erealx.2Ereal))\ V0t)) = (ap\ c.2Ereal.2Ereal_of_num \\ & c.2Enum.2E0))) \wedge (\forall V1s \in (2^{ty.2Erealx.2Ereal}). ((ap\ c.2Ereal_topology.2Esetdist \\ & (ap\ (ap\ (c.2Epair.2E.2C\ (2^{ty.2Erealx.2Ereal})\ (2^{ty.2Erealx.2Ereal})) \\ & V1s)\ (c.2Epred_set.2EEMPTY\ ty.2Erealx.2Ereal)))) = (ap\ c.2Ereal.2Ereal_of_num \\ & c.2Enum.2E0)))) \end{aligned}$$