

thm_2Ereal__topology_2ESPHERE__EMPTY (TMRe4KiQubrpDdJ7FeK4JJYW74LkXi7nG45)

October 26, 2020

Definition 1 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define $c_2Ebool_2E_2T$ to be $(ap (ap (c_2Emin_2E_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

Definition 3 We define $c_2Ebool_2E_21$ to be $\lambda A_27a : \iota.(\lambda V0P \in (2^{A_27a}).(ap (ap (c_2Emin_2E_3D (2^{A_27a}))$

Definition 4 We define $c_2Ebool_2E_2F$ to be $(ap (c_2Ebool_2E_21 2) (\lambda V0t \in 2.V0t))$.

Definition 5 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p P \Rightarrow p Q)$ of type ι .

Definition 6 We define $c_2Ebool_2E_27E$ to be $(\lambda V0t \in 2.(ap (ap c_2Emin_2E_3D_3D_3E V0t) c_2Ebool_2E_2F$

Definition 7 We define $c_2Ebool_2E_2F_5C$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21 2) (\lambda V2t \in 2.V2t))$

Let $c_2Enum_2EZERO_REP : \iota$ be given. Assume the following.

$$c_2Enum_2EZERO_REP \in \omega \tag{1}$$

Let $ty_2Enum_2Enum : \iota$ be given. Assume the following.

$$nonempty\ ty_2Enum_2Enum \tag{2}$$

Let $c_2Enum_2EABS_num : \iota$ be given. Assume the following.

$$c_2Enum_2EABS_num \in (ty_2Enum_2Enum^{\omega}) \tag{3}$$

Definition 8 We define c_2Enum_2E0 to be $(ap c_2Enum_2EABS_num c_2Enum_2EZERO_REP)$.

Let $ty_2Erealx_2Ereal : \iota$ be given. Assume the following.

$$nonempty\ ty_2Erealx_2Ereal \tag{4}$$

Let $c_2Ereal_2Ereal_of_num : \iota$ be given. Assume the following.

$$c_2Ereal_2Ereal_of_num \in (ty_2Erealx_2Ereal^{ty_2Enum_2Enum}) \tag{5}$$

Let $ty_2Ehreal_2Ehreal : \iota$ be given. Assume the following.

$$nonempty\ ty_2Ehreal_2Ehreal \quad (6)$$

Let $ty_2Epair_2Eprod : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A0.nonempty\ A0 \Rightarrow \forall A1.nonempty\ A1 \Rightarrow nonempty\ (ty_2Epair_2Eprod\ A0\ A1) \quad (7)$$

Let $c_2Erealax_2Ereal_REP_CLASS : \iota$ be given. Assume the following.

$$c_2Erealax_2Ereal_REP_CLASS \in ((2^{(ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal\ ty_2Ehreal_2Ehreal)})^{ty_2Erealax}) \quad (8)$$

Definition 9 We define c_2Emin_2E40 to be $\lambda A.\lambda P \in 2^A$. **if** $(\exists x \in A.p\ (ap\ P\ x))$ **then** $(the\ (\lambda x.x \in A \wedge p\ x))$ of type $\iota \Rightarrow \iota$.

Definition 10 We define $c_2Erealax_2Ereal_REP$ to be $\lambda V0a \in ty_2Erealax_2Ereal.(ap\ (c_2Emin_2E40\ (c_2Erealax_2Ereal_REP_CLASS\ V0a)))$

Let $c_2Erealax_2Etreallt : \iota$ be given. Assume the following.

$$c_2Erealax_2Etreallt \in ((2^{(ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal\ ty_2Ehreal_2Ehreal)})^{(ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal)}) \quad (9)$$

Definition 11 We define $c_2Erealax_2Ereal_lt$ to be $\lambda V0T1 \in ty_2Erealax_2Ereal.\lambda V1T2 \in ty_2Erealax_2Ereal.(c_2Erealax_2Etreallt\ V0T1\ V1T2)$

Definition 12 We define $c_2Epred_set_2EEMPTY$ to be $\lambda A_27a : \iota.(\lambda V0x \in A_27a.c_2Ebool_2EF)$.

Let $c_2Epair_2EABS_prod : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow \forall A_27b.nonempty\ A_27b \Rightarrow c_2Epair_2EABS_prod\ A_27a\ A_27b \in ((ty_2Epair_2Eprod\ A_27a\ A_27b)^{(2^{A_27b})^{A_27a}}) \quad (10)$$

Definition 13 We define c_2Epair_2E2C to be $\lambda A_27a : \iota.\lambda A_27b : \iota.\lambda V0x \in A_27a.\lambda V1y \in A_27b.(ap\ (c_2Epair_2EABS_prod\ A_27a\ A_27b)\ (V0x\ V1y))$

Let $c_2Ereal_topology_2Esphere : \iota$ be given. Assume the following.

$$c_2Ereal_topology_2Esphere \in ((2^{ty_2Erealax_2Ereal})^{(ty_2Epair_2Eprod\ ty_2Erealax_2Ereal\ ty_2Erealax_2Ereal)}) \quad (11)$$

Assume the following.

$$True \quad (12)$$

Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow (\forall V0t \in 2.((\forall V1x \in A_27a.(p\ V0t) \Leftrightarrow (p\ V1x)))) \quad (13)$$

Assume the following.

$$\begin{aligned} & (\forall V0t \in 2.(((True \Rightarrow (p\ V0t)) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \Rightarrow True) \Leftrightarrow \\ & True) \wedge (((False \Rightarrow (p\ V0t)) \Leftrightarrow True) \wedge (((p\ V0t) \Rightarrow (p\ V0t)) \Leftrightarrow True) \wedge ((\\ & (p\ V0t) \Rightarrow False) \Leftrightarrow (\neg (p\ V0t)))))) \end{aligned} \quad (14)$$

Assume the following.

$$\begin{aligned}
& (\forall V0a \in ty_2Erealax_2Ereal. (\forall V1r \in ty_2Erealax_2Ereal. \\
& (((ap\ c_2Ereal_topology_2Esphere\ (ap\ (ap\ (c_2Epair_2E_2C\ ty_2Erealax_2Ereal \\
& ty_2Erealax_2Ereal)\ V0a)\ V1r)) = (c_2Epred_set_2EEMPTY\ ty_2Erealax_2Ereal)) \Leftrightarrow \\
& (p\ (ap\ (ap\ c_2Erealax_2Ereal_lt\ V1r)\ (ap\ c_2Ereal_2Ereal_of_num \\
& c_2Enum_2E0))))))
\end{aligned} \tag{15}$$

Theorem 1

$$\begin{aligned}
& (\forall V0a \in ty_2Erealax_2Ereal. (\forall V1r \in ty_2Erealax_2Ereal. \\
& ((p\ (ap\ (ap\ c_2Erealax_2Ereal_lt\ V1r)\ (ap\ c_2Ereal_2Ereal_of_num \\
& c_2Enum_2E0))) \Rightarrow ((ap\ c_2Ereal_topology_2Esphere\ (ap\ (ap\ (c_2Epair_2E_2C \\
& ty_2Erealax_2Ereal\ ty_2Erealax_2Ereal)\ V0a)\ V1r)) = (c_2Epred_set_2EEMPTY \\
& ty_2Erealax_2Ereal))))))
\end{aligned}$$