

# thm\_2Ereal\_\_topology\_2ETRIVIAL\_\_LIMIT\_\_SEQUENTIALLY (TMKBBuTMntMHsdeptbjFa2izU73SidCZgzt)

October 26, 2020

Let  $ty\_2Enum\_2Enum : \iota$  be given. Assume the following.

$$nonempty\ ty\_2Enum\_2Enum \tag{1}$$

Let  $c\_2Enum\_2EREP\_num : \iota$  be given. Assume the following.

$$c\_2Enum\_2EREP\_num \in (\omega^{ty\_2Enum\_2Enum}) \tag{2}$$

Let  $c\_2Enum\_2ESUC\_REP : \iota$  be given. Assume the following.

$$c\_2Enum\_2ESUC\_REP \in (\omega^{\omega}) \tag{3}$$

Let  $c\_2Enum\_2EABS\_num : \iota$  be given. Assume the following.

$$c\_2Enum\_2EABS\_num \in (ty\_2Enum\_2Enum^{\omega}) \tag{4}$$

**Definition 1** We define  $c\_2Emin\_2E\_3D$  to be  $\lambda A.\lambda x \in A.\lambda y \in A.inj\_o (x = y)$  of type  $\iota \Rightarrow \iota$ .

**Definition 2** We define  $c\_2Ebool\_2ET$  to be  $(\lambda p (ap (c\_2Emin\_2E\_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

**Definition 3** We define  $c\_2Ebool\_2E\_21$  to be  $\lambda A.\lambda a : \iota.(\lambda V0P \in (2^{A-27a}).(ap (ap (c\_2Emin\_2E\_3D (2^{A-27a}))$

**Definition 4** We define  $c\_2Enum\_2ESUC$  to be  $\lambda V0m \in ty\_2Enum\_2Enum.(ap c\_2Enum\_2EABS\_num$

Let  $c\_2Enum\_2EZERO\_REP : \iota$  be given. Assume the following.

$$c\_2Enum\_2EZERO\_REP \in \omega \tag{5}$$

**Definition 5** We define  $c\_2Enum\_2E0$  to be  $(ap c\_2Enum\_2EABS\_num c\_2Enum\_2EZERO\_REP)$ .

**Definition 6** We define  $c\_2Ebool\_2EF$  to be  $(\lambda p (c\_2Ebool\_2E\_21 2) (\lambda V0t \in 2.V0t))$ .

**Definition 7** We define  $c\_2Emin\_2E\_3D\_3D\_3E$  to be  $\lambda P \in 2.\lambda Q \in 2.inj\_o (p P \Rightarrow p Q)$  of type  $\iota$ .

**Definition 8** We define  $c\_Ebool\_2E\_7E$  to be  $(\lambda V0t \in 2.(ap (ap c\_Emin\_2E\_3D\_3D\_3E V0t) c\_Ebool\_2E\_7E))$

**Definition 9** We define  $c\_Ebool\_2E\_2F\_5C$  to be  $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c\_Ebool\_2E\_21 2) (\lambda V2t \in 2)))$

**Definition 10** We define  $c\_Emin\_2E\_40$  to be  $\lambda A.\lambda P \in 2^A.$ if  $(\exists x \in A.p (ap P x))$  then (the  $(\lambda x.x \in A \wedge P x)$  of type  $\iota \Rightarrow \iota$ ).

**Definition 11** We define  $c\_Ebool\_2E\_3F$  to be  $\lambda A\_27a : \iota.(\lambda V0P \in (2^{A\_27a}).(ap V0P (ap (c\_Emin\_2E\_40 A\_27a) P)))$

**Definition 12** We define  $c\_Eprim\_rec\_2E\_3C$  to be  $\lambda V0m \in ty\_2Enum\_2Enum.\lambda V1n \in ty\_2Enum\_2Enum.$

**Definition 13** We define  $c\_Ebool\_2E\_5C\_2F$  to be  $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c\_Ebool\_2E\_21 2) (\lambda V2t \in 2)))$

**Definition 14** We define  $c\_Earithmetic\_2E\_3C\_3D$  to be  $\lambda V0m \in ty\_2Enum\_2Enum.\lambda V1n \in ty\_2Enum\_2Enum.$

**Definition 15** We define  $c\_Earithmetic\_2E\_3E$  to be  $\lambda V0m \in ty\_2Enum\_2Enum.\lambda V1n \in ty\_2Enum\_2Enum.$

**Definition 16** We define  $c\_Earithmetic\_2E\_3E\_3D$  to be  $\lambda V0m \in ty\_2Enum\_2Enum.\lambda V1n \in ty\_2Enum\_2Enum.$

Let  $ty\_2Ereal\_topology\_2Enet : \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A0.nonempty A0 \Rightarrow nonempty (ty\_2Ereal\_topology\_2Enet A0) \quad (6)$$

Let  $c\_2Ereal\_topology\_2Emk\_net : \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A\_27a.nonempty A\_27a \Rightarrow c\_2Ereal\_topology\_2Emk\_net A\_27a \in ((ty\_2Ereal\_topology\_2Enet A\_27a)^{(2^{A\_27a})^{A\_27a}}) \quad (7)$$

**Definition 17** We define  $c\_2Ereal\_topology\_2Esequentially$  to be  $(ap (c\_2Ereal\_topology\_2Emk\_net ty\_2Ereal\_topology\_2Enet))$

Let  $c\_2Ereal\_topology\_2Enetord : \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A\_27a.nonempty A\_27a \Rightarrow c\_2Ereal\_topology\_2Enetord A\_27a \in (((2^{A\_27a})^{A\_27a})^{(ty\_2Ereal\_topology\_2Enet A\_27a)}) \quad (8)$$

**Definition 18** We define  $c\_2Ereal\_topology\_2Etrivial\_limit$  to be  $\lambda A\_27a : \iota.\lambda V0net \in (ty\_2Ereal\_topology\_2Enetord A\_27a)$

Assume the following.

$$(\forall V0n \in ty\_2Enum\_2Enum.(\neg(c\_2Enum\_2E0 = (ap c\_2Enum\_2ESUC V0n)))) \quad (9)$$

Assume the following.

$$(\forall V0m \in ty\_2Enum\_2Enum.(p (ap (ap c\_2Earithmetic\_2E\_3C\_3D V0m) V0m))) \quad (10)$$

Assume the following.

$$(\forall V0n \in ty\_2Enum\_2Enum.(\forall V1m \in ty\_2Enum\_2Enum.(p (ap (ap c\_2Earithmetic\_2E\_3E\_3D V0n) V1m)) \Leftrightarrow (p (ap (ap c\_2Earithmetic\_2E\_3C\_3D V1m) V0n)))) \quad (11)$$

Assume the following.

$$True \quad (12)$$

Assume the following.

$$(\forall V0t1 \in 2.(\forall V1t2 \in 2.(((p V0t1) \Rightarrow (p V1t2)) \Rightarrow (((p V1t2) \Rightarrow (p V0t1)) \Rightarrow ((p V0t1) \Leftrightarrow (p V1t2)))))) \quad (13)$$

Assume the following.

$$(\forall V0t \in 2.(((p V0t) \Rightarrow False) \Rightarrow (\neg(p V0t)))) \quad (14)$$

Assume the following.

$$(\forall V0t \in 2.((\neg(p V0t)) \Rightarrow ((p V0t) \Rightarrow False))) \quad (15)$$

Assume the following.

$$(\forall V0t \in 2.(((True \Rightarrow (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \Rightarrow True) \Leftrightarrow True) \wedge (((False \Rightarrow (p V0t)) \Leftrightarrow True) \wedge (((p V0t) \Rightarrow (p V0t)) \Leftrightarrow True) \wedge ((p V0t) \Rightarrow False) \Leftrightarrow (\neg(p V0t)))))) \quad (16)$$

Assume the following.

$$((\forall V0t \in 2.((\neg(\neg(p V0t))) \Leftrightarrow (p V0t))) \wedge (((\neg True) \Leftrightarrow False) \wedge ((\neg False) \Leftrightarrow True))) \quad (17)$$

Assume the following.

$$(\forall V0t \in 2.(((True \Leftrightarrow (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \Leftrightarrow True) \Leftrightarrow (p V0t)) \wedge (((False \Leftrightarrow (p V0t)) \Leftrightarrow (\neg(p V0t))) \wedge (((p V0t) \Leftrightarrow False) \Leftrightarrow (\neg(p V0t)))))) \quad (18)$$

Assume the following.

$$\forall A\_27a.nonempty A\_27a \Rightarrow (\forall V0P \in (2^{A\_27a}).(\forall V1Q \in (2^{A\_27a}).((\forall V2x \in A\_27a.((p (ap V0P V2x)) \wedge (p (ap V1Q V2x)))) \Leftrightarrow ((\forall V3x \in A\_27a.(p (ap V0P V3x))) \wedge (\forall V4x \in A\_27a.(p (ap V1Q V4x))))))) \quad (19)$$

Assume the following.

$$\forall A\_27a.nonempty A\_27a \Rightarrow (\forall V0P \in 2.(\forall V1Q \in (2^{A\_27a}).(((p V0P) \vee (\exists V2x \in A\_27a.(p (ap V1Q V2x)))) \Leftrightarrow (\exists V3x \in A\_27a.((p V0P) \vee (p (ap V1Q V3x))))))) \quad (20)$$

Assume the following.

$$(\forall V0m \in ty\_2Enum\_2Enum.(\forall V1n \in ty\_2Enum\_2Enum.(p (ap (ap (ap (c\_2Ereal\_topology\_2Enetord ty\_2Enum\_2Enum) c\_2Ereal\_topology\_2Esequentially) V0m) V1n)) \Leftrightarrow (p (ap (ap c\_2Earithmic\_2E\_3E\_3D V0m) V1n)))) \quad (21)$$

Assume the following.

$$(\forall V0t \in 2.((\neg(\neg(p V0t))) \Leftrightarrow (p V0t))) \quad (22)$$

Assume the following.

$$(\forall V0A \in 2.((p V0A) \Rightarrow ((\neg(p V0A)) \Rightarrow False))) \quad (23)$$

Assume the following.

$$(\forall V0A \in 2.(\forall V1B \in 2.(((\neg((p V0A) \vee (p V1B))) \Rightarrow False) \Leftrightarrow ((p V0A) \Rightarrow False) \Rightarrow ((\neg(p V1B)) \Rightarrow False)))) \quad (24)$$

Assume the following.

$$(\forall V0A \in 2.(\forall V1B \in 2.(((\neg((\neg(p V0A)) \vee (p V1B))) \Rightarrow False) \Leftrightarrow ((p V0A) \Rightarrow ((\neg(p V1B)) \Rightarrow False)))) \quad (25)$$

Assume the following.

$$(\forall V0A \in 2.(((\neg(p V0A)) \Rightarrow False) \Rightarrow (((p V0A) \Rightarrow False) \Rightarrow False))) \quad (26)$$

Assume the following.

$$(\forall V0p \in 2.(\forall V1q \in 2.(\forall V2r \in 2.(((p V0p) \Leftrightarrow (p V1q) \Leftrightarrow (p V2r)) \Leftrightarrow (((p V0p) \vee ((p V1q) \vee (p V2r))) \wedge (((p V0p) \vee ((\neg(p V2r)) \vee (\neg(p V1q)))) \wedge (((p V1q) \vee ((\neg(p V2r)) \vee (\neg(p V0p)))) \wedge ((p V2r) \vee ((\neg(p V1q)) \vee (\neg(p V0p)))))))))) \quad (27)$$

Assume the following.

$$(\forall V0p \in 2.(\forall V1q \in 2.(\forall V2r \in 2.(((p V0p) \Leftrightarrow (p V1q) \wedge (p V2r)) \Leftrightarrow (((p V0p) \vee ((\neg(p V1q)) \vee (\neg(p V2r)))) \wedge (((p V1q) \vee (\neg(p V0p))) \wedge ((p V2r) \vee (\neg(p V0p)))))))) \quad (28)$$

Assume the following.

$$(\forall V0p \in 2.(\forall V1q \in 2.(\forall V2r \in 2.(((p V0p) \Leftrightarrow (p V1q) \vee (p V2r)) \Leftrightarrow (((p V0p) \vee (\neg(p V1q))) \wedge (((p V0p) \vee (\neg(p V2r))) \wedge ((p V1q) \vee ((p V2r) \vee (\neg(p V0p)))))))) \quad (29)$$

Assume the following.

$$(\forall V0p \in 2.(\forall V1q \in 2.(\forall V2r \in 2.(((p V0p) \Leftrightarrow (p V1q) \Rightarrow (p V2r)) \Leftrightarrow (((p V0p) \vee (p V1q)) \wedge (((p V0p) \vee (\neg(p V2r))) \wedge ((\neg(p V1q)) \vee ((p V2r) \vee (\neg(p V0p)))))))) \quad (30)$$

Assume the following.

$$(\forall V0p \in 2.(\forall V1q \in 2.(((p V0p) \Leftrightarrow (\neg(p V1q))) \Leftrightarrow (((p V0p) \vee (p V1q)) \wedge ((\neg(p V1q)) \vee (\neg(p V0p)))))) \quad (31)$$

**Theorem 1**

$$(\neg(p (ap (c\_2Ereal\_topology\_2Etrivial\_limit ty\_2Enum\_2Enum) c\_2Ereal\_topology\_2Esequentially))))$$