

# thm\_2Ereal\_\_topology\_2EUNCOUNTABLE\_\_OPEN (TMG3j7nZ8Jer6Q7Rfi3gQhHWju8bjmbwL9L)

October 26, 2020

**Definition 1** We define  $c\_2Emin\_2E\_3D$  to be  $\lambda A.\lambda x \in A.\lambda y \in A.inj\_o (x = y)$  of type  $\iota \Rightarrow \iota$ .

**Definition 2** We define  $c\_2Ebool\_2E\_2ET$  to be  $(ap (ap (c\_2Emin\_2E\_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

**Definition 3** We define  $c\_2Ebool\_2E\_2E\_21$  to be  $\lambda A\_27a : \iota.(\lambda V0P \in (2^{A-27a}).(ap (ap (c\_2Emin\_2E\_3D (2^{A-27a}))$

**Definition 4** We define  $c\_2Ebool\_2E\_2EF$  to be  $(ap (c\_2Ebool\_2E\_2E\_21 2) (\lambda V0t \in 2.V0t))$ .

Let  $ty\_2Eenum\_2Eenum : \iota$  be given. Assume the following.

$$nonempty\ ty\_2Eenum\_2Eenum \quad (1)$$

**Definition 5** We define  $c\_2Epred\_set\_2EUNIV$  to be  $\lambda A\_27a : \iota.(\lambda V0x \in A\_27a.c\_2Ebool\_2E\_2ET)$ .

**Definition 6** We define  $c\_2Emin\_2E\_3D\_3D\_3E$  to be  $\lambda P \in 2.\lambda Q \in 2.inj\_o (p \Rightarrow q)$  of type  $\iota$ .

**Definition 7** We define  $c\_2Ebool\_2E\_2EIN$  to be  $\lambda A\_27a : \iota.(\lambda V0x \in A\_27a.(\lambda V1f \in (2^{A-27a}).(ap V1f V0x)))$

**Definition 8** We define  $c\_2Ebool\_2E\_2E\_2F\_5C$  to be  $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c\_2Ebool\_2E\_2E\_21 2) (\lambda V2t \in 2.V2t)))$

**Definition 9** We define  $c\_2Epred\_set\_2EINJ$  to be  $\lambda A\_27a : \iota.\lambda A\_27b : \iota.\lambda V0f \in (A\_27b^{A-27a}).\lambda V1s \in (2^{A-27a})$

**Definition 10** We define  $c\_2Emin\_2E\_40$  to be  $\lambda A.\lambda P \in 2^A.if (\exists x \in A.p (ap P x)) \text{ then } (the (\lambda x.x \in A \wedge P x))$  of type  $\iota \Rightarrow \iota$ .

**Definition 11** We define  $c\_2Ebool\_2E\_3F$  to be  $\lambda A\_27a : \iota.(\lambda V0P \in (2^{A-27a}).(ap V0P (ap (c\_2Emin\_2E\_40$

**Definition 12** We define  $c\_2Epred\_set\_2Ecountable$  to be  $\lambda A\_27a : \iota.\lambda V0s \in (2^{A-27a}).(ap (c\_2Ebool\_2E\_3F$

**Definition 13** We define  $c\_2Epred\_set\_2ESURJ$  to be  $\lambda A\_27a : \iota.\lambda A\_27b : \iota.\lambda V0f \in (A\_27b^{A-27a}).\lambda V1s \in (2^{A-27a})$

**Definition 14** We define  $c\_2Epred\_set\_2EBIJ$  to be  $\lambda A\_27a : \iota.\lambda A\_27b : \iota.\lambda V0f \in (A\_27b^{A-27a}).\lambda V1s \in (2^{A-27a})$

**Definition 15** We define  $c\_2Ecardinal\_2Ecardeq$  to be  $\lambda A\_27a : \iota. \lambda A\_27b : \iota. \lambda V0s1 \in (2^{A\_27a}). \lambda V1s2 \in (2^{A\_27b}).$

**Definition 16** We define  $c\_2Epred\_set\_2EEMPTY$  to be  $\lambda A\_27a : \iota. (\lambda V0x \in A\_27a. c\_2Ebool\_2EF).$

**Definition 17** We define  $c\_2Ebool\_2E7E$  to be  $(\lambda V0t \in 2. (ap (ap c\_2Emin\_2E\_3D\_3D\_3E V0t) c\_2Ebool\_2EF)).$

Let  $ty\_2Erealax\_2Ereal : \iota$  be given. Assume the following.

$$nonempty\ ty\_2Erealax\_2Ereal \quad (2)$$

Let  $ty\_2Epair\_2Eprod : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A0. nonempty\ A0 \Rightarrow \forall A1. nonempty\ A1 \Rightarrow nonempty\ (ty\_2Epair\_2Eprod\ A0\ A1) \quad (3)$$

Let  $c\_2Epair\_2EABS\_prod : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A\_27a. nonempty\ A\_27a \Rightarrow \forall A\_27b. nonempty\ A\_27b \Rightarrow c\_2Epair\_2EABS\_prod\ A\_27a\ A\_27b \in ((ty\_2Epair\_2Eprod\ A\_27a\ A\_27b)^{(2^{A\_27b})^{A\_27a}}) \quad (4)$$

**Definition 18** We define  $c\_2Epair\_2E2C$  to be  $\lambda A\_27a : \iota. \lambda A\_27b : \iota. \lambda V0x \in A\_27a. \lambda V1y \in A\_27b. (ap (c\_2Emin\_2E\_3D\_3D\_3E V0t) c\_2Ebool\_2EF)).$

Let  $c\_2Ereal\_topology\_2EDist : \iota$  be given. Assume the following.

$$c\_2Ereal\_topology\_2EDist \in (ty\_2Erealax\_2Ereal^{(ty\_2Epair\_2Eprod\ ty\_2Erealax\_2Ereal\ ty\_2Erealax\_2Ereal)}) \quad (5)$$

Let  $ty\_2Ehreal\_2Ehreal : \iota$  be given. Assume the following.

$$nonempty\ ty\_2Ehreal\_2Ehreal \quad (6)$$

Let  $c\_2Erealax\_2Ereal\_REP\_CLASS : \iota$  be given. Assume the following.

$$c\_2Erealax\_2Ereal\_REP\_CLASS \in ((2^{(ty\_2Epair\_2Eprod\ ty\_2Ehreal\_2Ehreal\ ty\_2Ehreal\_2Ehreal)})^{ty\_2Erealax\_2Ereal}) \quad (7)$$

**Definition 19** We define  $c\_2Erealax\_2Ereal\_REP$  to be  $\lambda V0a \in ty\_2Erealax\_2Ereal. (ap (c\_2Emin\_2E\_40 V0t) c\_2Ebool\_2EF)).$

Let  $c\_2Erealax\_2Etreallt : \iota$  be given. Assume the following.

$$c\_2Erealax\_2Etreallt \in ((2^{(ty\_2Epair\_2Eprod\ ty\_2Ehreal\_2Ehreal\ ty\_2Ehreal\_2Ehreal)})^{(ty\_2Epair\_2Eprod\ ty\_2Ehreal\_2Ehreal)}) \quad (8)$$

**Definition 20** We define  $c\_2Erealax\_2Ereal\_lt$  to be  $\lambda V0T1 \in ty\_2Erealax\_2Ereal. \lambda V1T2 \in ty\_2Erealax\_2Ereal. (ap (c\_2Emin\_2E\_40 V0t) c\_2Ebool\_2EF)).$

Let  $c\_2Enum\_2EZERO\_REP : \iota$  be given. Assume the following.

$$c\_2Enum\_2EZERO\_REP \in \omega \quad (9)$$

Let  $c\_2Enum\_2EABS\_num : \iota$  be given. Assume the following.

$$c\_2Enum\_2EABS\_num \in (ty\_2Enum\_2Enum^{\omega}) \quad (10)$$

**Definition 21** We define  $c\_2Enum\_2E0$  to be  $(ap\ c\_2Enum\_2EABS\_num\ c\_2Enum\_2EZERO\_REP)$ .

Let  $c\_2Ereal\_2Ereal\_of\_num : \iota$  be given. Assume the following.

$$c\_2Ereal\_2Ereal\_of\_num \in (ty\_2Erealax\_2Ereal^{ty\_2Enum\_2Enum}) \quad (11)$$

**Definition 22** We define  $c\_2Ereal\_topology\_2EOpen$  to be  $\lambda V0s \in (2^{ty\_2Erealax\_2Ereal}).(ap\ (c\_2Ebool\_2E2$

Assume the following.

$$True \quad (12)$$

Assume the following.

$$(\forall V0t1 \in 2.(\forall V1t2 \in 2.(((p\ V0t1) \Rightarrow (p\ V1t2)) \Rightarrow (((p\ V1t2) \Rightarrow (p\ V0t1)) \Rightarrow ((p\ V0t1) \Leftrightarrow (p\ V1t2)))))) \quad (13)$$

Assume the following.

$$(\forall V0t \in 2.(False \Rightarrow (p\ V0t))) \quad (14)$$

Assume the following.

$$\forall A\_27a.nonempty\ A\_27a \Rightarrow (\forall V0t \in 2.((\forall V1x \in A\_27a.(p\ V0t)) \Leftrightarrow (p\ V0t))) \quad (15)$$

Assume the following.

$$(\forall V0t \in 2.(((True \wedge (p\ V0t)) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \wedge True) \Leftrightarrow (p\ V0t)) \wedge (((False \wedge (p\ V0t)) \Leftrightarrow False) \wedge (((p\ V0t) \wedge False) \Leftrightarrow False) \wedge (((p\ V0t) \wedge (p\ V0t)) \Leftrightarrow (p\ V0t)))))) \quad (16)$$

Assume the following.

$$(\forall V0t \in 2.(((True \Rightarrow (p\ V0t)) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \Rightarrow True) \Leftrightarrow True) \wedge (((False \Rightarrow (p\ V0t)) \Leftrightarrow True) \wedge (((p\ V0t) \Rightarrow (p\ V0t)) \Leftrightarrow True) \wedge ((p\ V0t) \Rightarrow False) \Leftrightarrow \neg(p\ V0t)))))) \quad (17)$$

Assume the following.

$$((\forall V0t \in 2.((\neg(\neg(p\ V0t))) \Leftrightarrow (p\ V0t))) \wedge (((\neg True) \Leftrightarrow False) \wedge ((\neg False) \Leftrightarrow True))) \quad (18)$$

Assume the following.

$$\forall A\_27a.nonempty\ A\_27a \Rightarrow (\forall V0x \in A\_27a.(\forall V1y \in A\_27a.((V0x = V1y) \Leftrightarrow (V1y = V0x)))) \quad (19)$$

Assume the following.

$$(\forall V0t \in 2.(((True \Leftrightarrow (p\ V0t)) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \Leftrightarrow True) \Leftrightarrow (p\ V0t)) \wedge (((False \Leftrightarrow (p\ V0t)) \Leftrightarrow \neg(p\ V0t)) \wedge (((p\ V0t) \Leftrightarrow False) \Leftrightarrow \neg(p\ V0t)))))) \quad (20)$$

Assume the following.

$$(\forall V0t1 \in 2.(\forall V1t2 \in 2.(\forall V2t3 \in 2.(((p V0t1) \Rightarrow ((p V1t2) \Rightarrow (p V2t3))) \Leftrightarrow (((p V0t1) \wedge (p V1t2)) \Rightarrow (p V2t3)))))) \quad (21)$$

Assume the following.

$$(\forall V0x \in 2.(\forall V1x_{27} \in 2.(\forall V2y \in 2.(\forall V3y_{27} \in 2.(((p V0x) \Leftrightarrow (p V1x_{27})) \wedge ((p V1x_{27}) \Rightarrow ((p V2y) \Leftrightarrow (p V3y_{27})))))) \Rightarrow ((p V0x) \Rightarrow (p V2y)) \Leftrightarrow ((p V1x_{27}) \Rightarrow (p V3y_{27})))))) \quad (22)$$

Assume the following.

$$(\forall V0s \in (2^{ty\_2Erealax\_2Ereal}).((p (ap (ap (c\_2Ecardinal\_2Ecardeq ty\_2Erealax\_2Ereal ty\_2Erealax\_2Ereal) V0s) (c\_2Epred\_set\_2EUNIV ty\_2Erealax\_2Ereal))) \Rightarrow (\neg (p (ap (c\_2Epred\_set\_2Ecountable ty\_2Erealax\_2Ereal) V0s)))))) \quad (23)$$

Assume the following.

$$(\forall V0s \in (2^{ty\_2Erealax\_2Ereal}).(((p (ap c\_2Ereal\_topology\_2EOpen V0s) \wedge (\neg (V0s = (c\_2Epred\_set\_2EEMPTY ty\_2Erealax\_2Ereal)))) \Rightarrow (p (ap (ap (c\_2Ecardinal\_2Ecardeq ty\_2Erealax\_2Ereal ty\_2Erealax\_2Ereal) V0s) (c\_2Epred\_set\_2EUNIV ty\_2Erealax\_2Ereal)))))) \quad (24)$$

**Theorem 1**

$$(\forall V0s \in (2^{ty\_2Erealax\_2Ereal}).(((p (ap c\_2Ereal\_topology\_2EOpen V0s) \wedge (\neg (V0s = (c\_2Epred\_set\_2EEMPTY ty\_2Erealax\_2Ereal)))) \Rightarrow (\neg (p (ap (c\_2Epred\_set\_2Ecountable ty\_2Erealax\_2Ereal) V0s))))))$$