

# thm\_2Ereal\_\_topology\_2EUNIFORMLY\_\_CONTINUOUS\_\_ON\_\_EQ (TMSPg7fbKnmyPfQrPiZ8s5MTDdUigobZimp)

October 26, 2020

**Definition 1** We define  $c\_2Emin\_2E\_3D$  to be  $\lambda A.\lambda x \in A.\lambda y \in A.inj\_o (x = y)$   
of type  $\iota \Rightarrow \iota$ .

**Definition 2** We define  $c\_2Ebool\_2E\_2E$  to be  $(ap (ap (c\_2Emin\_2E\_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

**Definition 3** We define  $c\_2Ebool\_2E\_21$  to be  $\lambda A\_27a : \iota.(\lambda V0P \in (2^{A\_27a}).(ap (ap (c\_2Emin\_2E\_3D (2^{A\_27a}))$

**Definition 4** We define  $c\_2Ebool\_2E\_2E$  to be  $(ap (c\_2Ebool\_2E\_21 2) (\lambda V0t \in 2.V0t))$ .

**Definition 5** We define  $c\_2Emin\_2E\_3D\_3D\_3E$  to be  $\lambda P \in 2.\lambda Q \in 2.inj\_o (p P \Rightarrow p Q)$   
of type  $\iota$ .

**Definition 6** We define  $c\_2Ebool\_2E\_7E$  to be  $(\lambda V0t \in 2.(ap (ap c\_2Emin\_2E\_3D\_3D\_3E V0t) c\_2Ebool\_2E\_2E$

**Definition 7** We define  $c\_2Ebool\_2E\_2F\_5C$  to be  $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c\_2Ebool\_2E\_21 2) (\lambda V2t \in 2.V2t))$

Let  $ty\_2Epair\_2Eprod : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A0.nonempty A0 \Rightarrow \forall A1.nonempty A1 \Rightarrow nonempty (ty\_2Epair\_2Eprod A0 A1) \quad (1)$$

Let  $c\_2Epair\_2EABS\_prod : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A\_27a.nonempty A\_27a \Rightarrow \forall A\_27b.nonempty A\_27b \Rightarrow c\_2Epair\_2EABS\_prod A\_27a A\_27b \in ((ty\_2Epair\_2Eprod A\_27a A\_27b)^{(2^{A\_27b})^{A\_27a}}) \quad (2)$$

**Definition 8** We define  $c\_2Epair\_2E\_2C$  to be  $\lambda A\_27a : \iota.\lambda A\_27b : \iota.\lambda V0x \in A\_27a.\lambda V1y \in A\_27b.(ap (c\_2E$

Let  $ty\_2Erealx\_2Ereal : \iota$  be given. Assume the following.

$$nonempty ty\_2Erealx\_2Ereal \quad (3)$$

Let  $c\_2Ereal\_topology\_2EDist : \iota$  be given. Assume the following.

$$c\_2Ereal\_topology\_2EDist \in (ty\_2Erealx\_2Ereal^{(ty\_2Epair\_2Eprod ty\_2Erealx\_2Ereal ty\_2Erealx\_2Ereal)}) \quad (4)$$

**Definition 9** We define  $c\_2Ebool\_2EIN$  to be  $\lambda A\_27a : \iota. (\lambda V0x \in A\_27a. (\lambda V1f \in (2^{A\_27a}). (ap\ V1f\ V0x)))$

**Definition 10** We define  $c\_2Emin\_2E40$  to be  $\lambda A. \lambda P \in 2^A. \text{if } (\exists x \in A. p\ (ap\ P\ x)) \text{ then } (the\ (\lambda x. x \in A) \wedge p)$  of type  $\iota \Rightarrow \iota$ .

**Definition 11** We define  $c\_2Ebool\_2E3F$  to be  $\lambda A\_27a : \iota. (\lambda V0P \in (2^{A\_27a}). (ap\ V0P\ (ap\ (c\_2Emin\_2E40\ P))))$

Let  $c\_2Enum\_2EZERO\_REP : \iota$  be given. Assume the following.

$$c\_2Enum\_2EZERO\_REP \in \omega \tag{5}$$

Let  $ty\_2Enum\_2Enum : \iota$  be given. Assume the following.

$$nonempty\ ty\_2Enum\_2Enum \tag{6}$$

Let  $c\_2Enum\_2EABS\_num : \iota$  be given. Assume the following.

$$c\_2Enum\_2EABS\_num \in (ty\_2Enum\_2Enum^{\omega}) \tag{7}$$

**Definition 12** We define  $c\_2Enum\_2E0$  to be  $(ap\ c\_2Enum\_2EABS\_num\ c\_2Enum\_2EZERO\_REP)$ .

Let  $c\_2Ereal\_2Ereal\_of\_num : \iota$  be given. Assume the following.

$$c\_2Ereal\_2Ereal\_of\_num \in (ty\_2Erealax\_2Ereal^{ty\_2Enum\_2Enum}) \tag{8}$$

Let  $ty\_2Ehreal\_2Ehreal : \iota$  be given. Assume the following.

$$nonempty\ ty\_2Ehreal\_2Ehreal \tag{9}$$

Let  $c\_2Erealax\_2Ereal\_REP\_CLASS : \iota$  be given. Assume the following.

$$c\_2Erealax\_2Ereal\_REP\_CLASS \in ((2^{(ty\_2Epair\_2Eprod\ ty\_2Ehreal\_2Ehreal\ ty\_2Ehreal\_2Ehreal)})^{ty\_2Erealax}) \tag{10}$$

**Definition 13** We define  $c\_2Erealax\_2Ereal\_REP$  to be  $\lambda V0a \in ty\_2Erealax\_2Ereal. (ap\ (c\_2Emin\_2E40\ a))$

Let  $c\_2Erealax\_2Etrealt\_lt : \iota$  be given. Assume the following.

$$c\_2Erealax\_2Etrealt\_lt \in ((2^{(ty\_2Epair\_2Eprod\ ty\_2Ehreal\_2Ehreal\ ty\_2Ehreal\_2Ehreal)})^{(ty\_2Epair\_2Eprod\ ty\_2Ehreal\_2Ehreal)}) \tag{11}$$

**Definition 14** We define  $c\_2Erealax\_2Ereal\_lt$  to be  $\lambda V0T1 \in ty\_2Erealax\_2Ereal. \lambda V1T2 \in ty\_2Erealax\_2Ereal.$

**Definition 15** We define  $c\_2Ereal\_topology\_2Euniformly\_continuous\_on$  to be  $\lambda V0f \in (ty\_2Erealax\_2Ereal^{ty\_2Erealax\_2Ereal}). \lambda V1s \in (2^{ty\_2Erealax\_2Ereal}). (ap\ (c\_2Ebool\_2E21\ ty\_2Erealax\_2Ereal\ f\ s))$

Assume the following.

$$True \tag{12}$$

Assume the following.

$$\forall A\_27a. nonempty\ A\_27a \Rightarrow (\forall V0t \in 2. ((\forall V1x \in A\_27a. (p\ V0t)) \Leftrightarrow (p\ V0t))) \tag{13}$$

Assume the following.

$$(\forall V0t \in 2.(((True \Rightarrow (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \Rightarrow True) \Leftrightarrow True) \wedge (((False \Rightarrow (p V0t)) \Leftrightarrow True) \wedge (((p V0t) \Rightarrow (p V0t)) \Leftrightarrow True) \wedge ((p V0t) \Rightarrow False) \Leftrightarrow (\neg(p V0t)))))) \quad (14)$$

Assume the following.

$$\forall A\_27a.nonempty\ A\_27a \Rightarrow (\forall V0x \in A\_27a. (\forall V1y \in A\_27a. ((V0x = V1y) \Leftrightarrow (V1y = V0x)))) \quad (15)$$

Assume the following.

$$(\forall V0t \in 2.(((True \Leftrightarrow (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \Leftrightarrow True) \Leftrightarrow (p V0t)) \wedge (((False \Leftrightarrow (p V0t)) \Leftrightarrow (\neg(p V0t))) \wedge (((p V0t) \Leftrightarrow False) \Leftrightarrow (\neg(p V0t)))))) \quad (16)$$

Assume the following.

$$(\forall V0t1 \in 2. (\forall V1t2 \in 2. (\forall V2t3 \in 2. (((p V0t1) \Rightarrow (p V1t2) \Rightarrow (p V2t3)) \Leftrightarrow (((p V0t1) \wedge (p V1t2)) \Rightarrow (p V2t3)))))) \quad (17)$$

Assume the following.

$$(\forall V0x \in 2. (\forall V1x\_27 \in 2. (\forall V2y \in 2. (\forall V3y\_27 \in 2. (((p V0x) \Leftrightarrow (p V1x\_27)) \wedge ((p V1x\_27) \Rightarrow ((p V2y) \Leftrightarrow (p V3y\_27)))))) \Rightarrow (((p V0x) \Rightarrow (p V2y)) \Leftrightarrow ((p V1x\_27) \Rightarrow (p V3y\_27)))))) \quad (18)$$

Assume the following.

$$(\forall V0r \in 2. (\forall V1p \in 2. (\forall V2q \in 2. (((p V1p) \wedge (p V2q) \Rightarrow (p V0r)) \Leftrightarrow ((p V1p) \Rightarrow ((p V2q) \Rightarrow (p V0r)))))) \quad (19)$$

### Theorem 1

$$(\forall V0f \in (ty\_2Erealax\_2Ereal^{ty\_2Erealax\_2Ereal}). (\forall V1g \in (ty\_2Erealax\_2Ereal^{ty\_2Erealax\_2Ereal}). (\forall V2s \in (2^{ty\_2Erealax\_2Ereal}). (((\forall V3x \in ty\_2Erealax\_2Ereal. ((p (ap (ap (c\_2Ebool\_2EIN ty\_2Erealax\_2Ereal) V3x) V2s)) \Rightarrow ((ap V0f V3x) = (ap V1g V3x)))) \wedge (p (ap (ap c\_2Ereal\_topology\_2Euniformly\_continuous\_on V0f) V2s))) \Rightarrow (p (ap (ap c\_2Ereal\_topology\_2Euniformly\_continuous\_on V1g) V2s))))))$$