

thm_2Ereal__topology_2EWITHIN__UNIV (TM- cPT5sUMoPiCFt6gwHPaVPSY4FhiLK4mAo)

October 26, 2020

Definition 1 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define $c_2Ebool_2E_2E$ to be $(ap (ap (c_2Emin_2E_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

Definition 3 We define $c_2Ebool_2E_2E1$ to be $\lambda A_27a : \iota.(\lambda V0P \in (2^{A_27a}).(ap (ap (c_2Emin_2E_3D (2^{A_27a}))$

Definition 4 We define $c_2Ebool_2E_2EF$ to be $(ap (c_2Ebool_2E_2E1 2) (\lambda V0t \in 2.V0t))$.

Definition 5 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p \Rightarrow q)$ of type ι .

Definition 6 We define $c_2Ebool_2E_2E7E$ to be $(\lambda V0t \in 2.(ap (ap c_2Emin_2E_3D_3D_3E V0t) c_2Ebool_2E_2EF$

Definition 7 We define $c_2Epred_set_2EUNIV$ to be $\lambda A_27a : \iota.(\lambda V0x \in A_27a.c_2Ebool_2E_2E7E)$.

Definition 8 We define $c_2Ebool_2E_2E5C_2E2F$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_2E1 2) (\lambda V2t \in 2.V2t))$

Let $ty_2Ehreal_2Ehreal : \iota$ be given. Assume the following.

$$nonempty\ ty_2Ehreal_2Ehreal \tag{1}$$

Let $ty_2Epair_2Eprod : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A0.nonempty\ A0 \Rightarrow \forall A1.nonempty\ A1 \Rightarrow nonempty\ (ty_2Epair_2Eprod\ A0\ A1) \tag{2}$$

Let $ty_2Erealax_2Ereal : \iota$ be given. Assume the following.

$$nonempty\ ty_2Erealax_2Ereal \tag{3}$$

Let $c_2Erealax_2Ereal_REP_CLASS : \iota$ be given. Assume the following.

$$c_2Erealax_2Ereal_REP_CLASS \in ((2^{(ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal\ ty_2Ehreal_2Ehreal)})ty_2Erealax_2Ereal) \tag{4}$$

Definition 9 We define $c_2Emin_2E_40$ to be $\lambda A.\lambda P \in 2^A.$ **if** $(\exists x \in A.p (ap P x))$ **then** $(the (\lambda x.x \in A \wedge p$
of type $\iota \Rightarrow \iota$.

Definition 10 We define $c_2Erealax_2Ereal_REP$ to be $\lambda V0a \in ty_2Erealax_2Ereal.(ap (c_2Emin_2E_40 (t$

Let $c_2Erealax_2Etrealt_lt : \iota$ be given. Assume the following.

$$c_2Erealax_2Etrealt_lt \in ((2^{(ty_2Epair_2Eprod ty_2Ehreal_2Ehreal ty_2Ehreal_2Ehreal)})(ty_2Epair_2Eprod ty_2Ehreal_2Ehreal)) \quad (5)$$

Definition 11 We define $c_2Erealax_2Ereal_lt$ to be $\lambda V0T1 \in ty_2Erealax_2Ereal.\lambda V1T2 \in ty_2Erealax_2Ereal$

Definition 12 We define $c_2Ereal_2Ereal_lte$ to be $\lambda V0x \in ty_2Erealax_2Ereal.\lambda V1y \in ty_2Erealax_2Ereal$

Definition 13 We define $c_2Ebool_2E_2F_5C$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21 2) (\lambda V2t \in 2$

Let $c_2Epair_2EABS_prod : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A_27a.nonempty A_27a \Rightarrow \forall A_27b.nonempty A_27b \Rightarrow c_2Epair_2EABS_prod A_27a A_27b \in ((ty_2Epair_2Eprod A_27a A_27b)^{(2^{A_27b})^{A_27a}}) \quad (6)$$

Definition 14 We define $c_2Epair_2E_2C$ to be $\lambda A_27a : \iota.\lambda A_27b : \iota.\lambda V0x \in A_27a.\lambda V1y \in A_27b.(ap (c_2E$

Let $c_2Ereal_topology_2EDist : \iota$ be given. Assume the following.

$$c_2Ereal_topology_2EDist \in (ty_2Erealax_2Ereal^{(ty_2Epair_2Eprod ty_2Erealax_2Ereal ty_2Erealax_2Ereal)}) \quad (7)$$

Let $c_2Enum_2EZERO_REP : \iota$ be given. Assume the following.

$$c_2Enum_2EZERO_REP \in \omega \quad (8)$$

Let $ty_2Enum_2Enum : \iota$ be given. Assume the following.

$$nonempty ty_2Enum_2Enum \quad (9)$$

Let $c_2Enum_2EABS_num : \iota$ be given. Assume the following.

$$c_2Enum_2EABS_num \in (ty_2Enum_2Enum^{\omega}) \quad (10)$$

Definition 15 We define c_2Enum_2E0 to be $(ap c_2Enum_2EABS_num c_2Enum_2EZERO_REP)$.

Let $c_2Ereal_2Ereal_of_num : \iota$ be given. Assume the following.

$$c_2Ereal_2Ereal_of_num \in (ty_2Erealax_2Ereal^{ty_2Enum_2Enum}) \quad (11)$$

Let $ty_2Ereal_topology_2Enet : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A0.nonempty A0 \Rightarrow nonempty (ty_2Ereal_topology_2Enet A0) \quad (12)$$

Let $c_2Ereal_topology_2Emk_net : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A_27a.nonempty A_27a \Rightarrow c_2Ereal_topology_2Emk_net A_27a \in ((ty_2Ereal_topology_2Enet A_27a)^{(2^{A_27a})^{A_27a}}) \quad (13)$$

Definition 16 We define $c_2Ereal_topology_2Eat$ to be $\lambda V0a \in ty_2Erealax_2Ereal.(ap (c_2Ereal_topology_2Eat$

Definition 17 We define c_2Ebool_2EIN to be $\lambda A_27a : \iota.(\lambda V0x \in A_27a.(\lambda V1f \in (2^{A_27a}).(ap V1f V0x)))$

Let $c_2Ereal_topology_2Enetord : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A_27a.nonempty A_27a \Rightarrow c_2Ereal_topology_2Enetord A_27a \in ((2^{A_27a})^{A_27a})^{(ty_2Ereal_topology_2Enet A_27a)} \quad (14)$$

Definition 18 We define $c_2Ereal_topology_2Ewithin$ to be $\lambda A_27a : \iota.\lambda V0net \in (ty_2Ereal_topology_2Enet$

Assume the following.

$$True \quad (15)$$

Assume the following.

$$\forall A_27a.nonempty A_27a \Rightarrow (\forall V0t \in 2.((\forall V1x \in A_27a.(p V0t)) \Leftrightarrow (p V0t))) \quad (16)$$

Assume the following.

$$\begin{aligned} & (\forall V0t \in 2.(((True \wedge (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \wedge True) \Leftrightarrow \\ & (p V0t)) \wedge (((False \wedge (p V0t)) \Leftrightarrow False) \wedge (((p V0t) \wedge False) \Leftrightarrow False) \wedge \\ & (((p V0t) \wedge (p V0t)) \Leftrightarrow (p V0t)))))) \quad (17) \end{aligned}$$

Assume the following.

$$\forall A_27a.nonempty A_27a \Rightarrow (\forall V0x \in A_27a.((V0x = V0x) \Leftrightarrow True)) \quad (18)$$

Assume the following.

$$\begin{aligned} & (\forall V0t \in 2.(((True \Leftrightarrow (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \Leftrightarrow True) \Leftrightarrow \\ & (p V0t)) \wedge (((False \Leftrightarrow (p V0t)) \Leftrightarrow \neg(p V0t)) \wedge (((p V0t) \Leftrightarrow False) \Leftrightarrow \neg \\ & (p V0t)))))) \quad (19) \end{aligned}$$

Assume the following.

$$\forall A_27a.nonempty A_27a \Rightarrow (\forall V0x \in A_27a.(p (ap (ap (c_2Ebool_2EIN A_27a) V0x) (c_2Epred_set_2EUNIV A_27a)))) \quad (20)$$

Assume the following.

$$\begin{aligned} & \forall A_27a.nonempty A_27a \Rightarrow \forall A_27b.nonempty A_27b \Rightarrow (\\ & (\forall V0a \in (ty_2Ereal_topology_2Enet A_27a).((ap (c_2Ereal_topology_2Emk_net \\ & A_27a) (ap (c_2Ereal_topology_2Enetord A_27a) V0a)) = V0a)) \wedge \\ & (\forall V1r \in ((2^{A_27b})^{A_27b}).((\forall V2x \in A_27b.(\forall V3y \in \\ & A_27b.((\forall V4z \in A_27b.((p (ap (ap V1r V4z) V2x)) \Rightarrow (p (ap (ap \\ & V1r V4z) V3y)))) \vee (\forall V5z \in A_27b.((p (ap (ap V1r V5z) V3y)) \Rightarrow \\ & (p (ap (ap V1r V5z) V2x)))))) \Leftrightarrow ((ap (c_2Ereal_topology_2Enetord \\ & A_27b) (ap (c_2Ereal_topology_2Emk_net A_27b) V1r)) = V1r)))) \quad (21) \end{aligned}$$

Theorem 1

$$(\forall V0x \in ty_2Erealax_2Ereal. ((ap (ap (c_2Ereal_topology_2Ewithin ty_2Erealax_2Ereal) (ap c_2Ereal_topology_2Eat V0x)) (c_2Epred_set_2EUNIV ty_2Erealax_2Ereal)) = (ap c_2Ereal_topology_2Eat V0x)))$$