

# thm\_2Ereal\_\_topology\_2EWLOG\_\_LE (TM- bgxX3nzUhddDeQwEkFcNnqUU8KaqJBgWp)

October 26, 2020

**Definition 1** We define  $c\_2Emin\_2E\_3D$  to be  $\lambda A.\lambda x \in A.\lambda y \in A.inj\_o (x = y)$  of type  $\iota \Rightarrow \iota$ .

**Definition 2** We define  $c\_2Ebool\_2E\_2T$  to be  $(ap (ap (c\_2Emin\_2E\_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

Let  $ty\_2Enum\_2Enum : \iota$  be given. Assume the following.

$$nonempty\ ty\_2Enum\_2Enum \tag{1}$$

**Definition 3** We define  $c\_2Ebool\_2E\_21$  to be  $\lambda A.\lambda a : \iota.(\lambda V0P \in (2^{A-27a}).(ap (ap (c\_2Emin\_2E\_3D (2^{A-27a}))$

**Definition 4** We define  $c\_2Ebool\_2E\_2F$  to be  $(ap (c\_2Ebool\_2E\_21 2) (\lambda V0t \in 2.V0t))$ .

**Definition 5** We define  $c\_2Emin\_2E\_3D\_3D\_3E$  to be  $\lambda P \in 2.\lambda Q \in 2.inj\_o (p P \Rightarrow p Q)$  of type  $\iota$ .

**Definition 6** We define  $c\_2Ebool\_2E\_27E$  to be  $(\lambda V0t \in 2.(ap (ap c\_2Emin\_2E\_3D\_3D\_3E V0t) c\_2Ebool\_2E\_2F$

**Definition 7** We define  $c\_2Ebool\_2E\_2F\_5C$  to be  $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c\_2Ebool\_2E\_21 2) (\lambda V2t \in 2.V2t))$

Let  $c\_2Enum\_2EREP\_num : \iota$  be given. Assume the following.

$$c\_2Enum\_2EREP\_num \in (\omega^{ty\_2Enum\_2Enum}) \tag{2}$$

Let  $c\_2Enum\_2ESUC\_REP : \iota$  be given. Assume the following.

$$c\_2Enum\_2ESUC\_REP \in (\omega^{\omega}) \tag{3}$$

Let  $c\_2Enum\_2EABS\_num : \iota$  be given. Assume the following.

$$c\_2Enum\_2EABS\_num \in (ty\_2Enum\_2Enum^{\omega}) \tag{4}$$

**Definition 8** We define  $c\_2Enum\_2ESUC$  to be  $\lambda V0m \in ty\_2Enum\_2Enum.(ap c\_2Enum\_2EABS\_num (ap c\_2Enum\_2EREP\_num (ap c\_2Enum\_2ESUC\_REP m)))$

**Definition 9** We define  $c\_2Emin\_2E\_40$  to be  $\lambda A.\lambda P \in 2^A.if (\exists x \in A.p (ap P x)) \mathbf{then} (the (\lambda x.x \in A \wedge P x))$  of type  $\iota \Rightarrow \iota$ .

**Definition 10** We define  $c\_2Ebool\_2E\_3F$  to be  $\lambda A\_27a : \iota. (\lambda V0P \in (2^{A\_27a}). (ap\ V0P\ (ap\ (c\_2Emin\_2E\_40$

**Definition 11** We define  $c\_2Eprim\_rec\_2E\_3C$  to be  $\lambda V0m \in ty\_2Enum\_2Enum. \lambda V1n \in ty\_2Enum\_2Enum$

**Definition 12** We define  $c\_2Ebool\_2E\_5C\_2F$  to be  $(\lambda V0t1 \in 2. (\lambda V1t2 \in 2. (ap\ (c\_2Ebool\_2E\_21\ 2)\ (\lambda V2t \in$

**Definition 13** We define  $c\_2Earithmetic\_2E\_3C\_3D$  to be  $\lambda V0m \in ty\_2Enum\_2Enum. \lambda V1n \in ty\_2Enum\_2Enum$

Assume the following.

$$True \tag{5}$$

Assume the following.

$$\begin{aligned} & (\forall V0t \in 2. (((True \Rightarrow (p\ V0t)) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \Rightarrow True) \Leftrightarrow \\ & True) \wedge (((False \Rightarrow (p\ V0t)) \Leftrightarrow True) \wedge (((p\ V0t) \Rightarrow (p\ V0t)) \Leftrightarrow True) \wedge (( \\ & (p\ V0t) \Rightarrow False) \Leftrightarrow (\neg(p\ V0t)))))) \end{aligned} \tag{6}$$

Assume the following.

$$\begin{aligned} & ((\forall V0t \in 2. ((\neg(\neg(p\ V0t))) \Leftrightarrow (p\ V0t))) \wedge ((\neg True) \Leftrightarrow False) \wedge \\ & ((\neg False) \Leftrightarrow True)) \end{aligned} \tag{7}$$

Assume the following.

$$\begin{aligned} & (\forall V0t \in 2. (((True \Leftrightarrow (p\ V0t)) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \Leftrightarrow True) \Leftrightarrow \\ & (p\ V0t)) \wedge (((False \Leftrightarrow (p\ V0t)) \Leftrightarrow (\neg(p\ V0t))) \wedge (((p\ V0t) \Leftrightarrow False) \Leftrightarrow (\neg( \\ & p\ V0t)))))) \end{aligned} \tag{8}$$

Assume the following.

$$\begin{aligned} & (\forall V0m \in ty\_2Enum\_2Enum. (\forall V1n \in ty\_2Enum\_2Enum. ( \\ & (p\ (ap\ (ap\ c\_2Earithmetic\_2E\_3C\_3D\ V0m)\ V1n)) \vee (p\ (ap\ (ap\ c\_2Earithmetic\_2E\_3C\_3D \\ & V1n)\ V0m)))) \end{aligned} \tag{9}$$

Assume the following.

$$(\forall V0t \in 2. ((\neg(\neg(p\ V0t))) \Leftrightarrow (p\ V0t))) \tag{10}$$

Assume the following.

$$(\forall V0A \in 2. ((p\ V0A) \Rightarrow ((\neg(p\ V0A)) \Rightarrow False))) \tag{11}$$

Assume the following.

$$\begin{aligned} & (\forall V0A \in 2. (\forall V1B \in 2. (((\neg((p\ V0A) \vee (p\ V1B))) \Rightarrow False) \Leftrightarrow \\ & (((p\ V0A) \Rightarrow False) \Rightarrow ((\neg(p\ V1B)) \Rightarrow False)))) \end{aligned} \tag{12}$$

Assume the following.

$$\begin{aligned} & (\forall V0A \in 2. (\forall V1B \in 2. (((\neg((\neg(p\ V0A)) \vee (p\ V1B))) \Rightarrow False) \Leftrightarrow \\ & ((p\ V0A) \Rightarrow ((\neg(p\ V1B)) \Rightarrow False)))) \end{aligned} \tag{13}$$

Assume the following.

$$(\forall V0A \in 2.((\neg(p V0A)) \Rightarrow False) \Rightarrow ((p V0A) \Rightarrow False) \Rightarrow False)) \quad (14)$$

Assume the following.

$$\begin{aligned} & (\forall V0p \in 2.(\forall V1q \in 2.(\forall V2r \in 2.(((p V0p) \Leftrightarrow ( \\ & (p V1q) \Leftrightarrow (p V2r))) \Leftrightarrow (((p V0p) \vee ((p V1q) \vee (p V2r))) \wedge (((p V0p) \vee (\neg( \\ & p V2r)) \vee (\neg(p V1q)))) \wedge (((p V1q) \vee (\neg(p V2r)) \vee (\neg(p V0p)))) \wedge ((p V2r) \vee \\ & ((\neg(p V1q)) \vee (\neg(p V0p)))))))))) \end{aligned} \quad (15)$$

Assume the following.

$$\begin{aligned} & (\forall V0p \in 2.(\forall V1q \in 2.(\forall V2r \in 2.(((p V0p) \Leftrightarrow ( \\ & (p V1q) \wedge (p V2r))) \Leftrightarrow (((p V0p) \vee (\neg(p V1q)) \vee (\neg(p V2r))) \wedge (((p V1q) \vee \\ & (\neg(p V0p))) \wedge ((p V2r) \vee (\neg(p V0p)))))))) \end{aligned} \quad (16)$$

Assume the following.

$$\begin{aligned} & (\forall V0p \in 2.(\forall V1q \in 2.(\forall V2r \in 2.(((p V0p) \Leftrightarrow ( \\ & (p V1q) \vee (p V2r))) \Leftrightarrow (((p V0p) \vee (\neg(p V1q))) \wedge (((p V0p) \vee (\neg(p V2r))) \wedge \\ & ((p V1q) \vee ((p V2r) \vee (\neg(p V0p)))))))))) \end{aligned} \quad (17)$$

Assume the following.

$$\begin{aligned} & (\forall V0p \in 2.(\forall V1q \in 2.(\forall V2r \in 2.(((p V0p) \Leftrightarrow ( \\ & (p V1q) \Rightarrow (p V2r))) \Leftrightarrow (((p V0p) \vee (p V1q)) \wedge (((p V0p) \vee (\neg(p V2r))) \wedge (( \\ & \neg(p V1q)) \vee ((p V2r) \vee (\neg(p V0p)))))))))) \end{aligned} \quad (18)$$

Assume the following.

$$\begin{aligned} & (\forall V0p \in 2.(\forall V1q \in 2.(((p V0p) \Leftrightarrow (\neg(p V1q))) \Leftrightarrow (((p V0p) \vee \\ & (p V1q)) \wedge ((\neg(p V1q)) \vee (\neg(p V0p)))))) \end{aligned} \quad (19)$$

Assume the following.

$$(\forall V0p \in 2.(\forall V1q \in 2.((\neg((p V0p) \Rightarrow (p V1q))) \Rightarrow (p V0p))) \quad (20)$$

Assume the following.

$$(\forall V0p \in 2.(\forall V1q \in 2.((\neg((p V0p) \Rightarrow (p V1q))) \Rightarrow (\neg(p V1q)))) \quad (21)$$

### Theorem 1

$$\begin{aligned} & (\forall V0P \in ((2^{ty\_2Enum\_2Enum})^{ty\_2Enum\_2Enum}).(((\forall V1m \in \\ & ty\_2Enum\_2Enum.(\forall V2n \in ty\_2Enum\_2Enum.((p (ap (ap V0P V1m) \\ & V2n)) \Leftrightarrow (p (ap (ap V0P V2n) V1m)))))) \wedge (\forall V3m \in ty\_2Enum\_2Enum. \\ & (\forall V4n \in ty\_2Enum\_2Enum.((p (ap (ap c\_2Earithmetic\_2E\_3C\_3D \\ & V3m) V4n)) \Rightarrow (p (ap (ap V0P V3m) V4n)))))) \Rightarrow (\forall V5m \in ty\_2Enum\_2Enum. \\ & (\forall V6n \in ty\_2Enum\_2Enum.(p (ap (ap V0P V5m) V6n)))))) \end{aligned}$$