

thm_2Erealax_2EHREAL_EQ_ADDL
(TMTJuTt6b4UoKfhcXBSSYZ2WuUGTH2Z93ye)

October 26, 2020

Let $ty_2Ehurat_2Ehurat : \iota$ be given. Assume the following.

$$nonempty\ ty_2Ehurat_2Ehurat \tag{1}$$

Let $ty_2Ehreal_2Ehreal : \iota$ be given. Assume the following.

$$nonempty\ ty_2Ehreal_2Ehreal \tag{2}$$

Let $c_2Ehreal_2Ecut : \iota$ be given. Assume the following.

$$c_2Ehreal_2Ecut \in ((2^{ty_2Ehurat_2Ehurat})^{ty_2Ehreal_2Ehreal}) \tag{3}$$

Definition 1 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p \Rightarrow Q)$ of type ι .

Definition 2 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 3 We define $c_2Ebool_2E_2T$ to be $(ap (ap (c_2Emin_2E_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

Definition 4 We define $c_2Ebool_2E_21$ to be $\lambda A_27a : \iota.(\lambda V0P \in (2^{A_27a}).(ap (ap (c_2Emin_2E_3D (2^{A_27a}))$

Definition 5 We define $c_2Ebool_2E_2F_5C$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21 2) (\lambda V2t \in 2$

Let $ty_2Enum_2Enum : \iota$ be given. Assume the following.

$$nonempty\ ty_2Enum_2Enum \tag{4}$$

Let $ty_2Epair_2Eprod : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A0.nonempty\ A0 \Rightarrow \forall A1.nonempty\ A1 \Rightarrow nonempty\ (ty_2Epair_2Eprod\ A0\ A1) \tag{5}$$

Let $c_2Ehurat_2Ehurat_REP_CLASS : \iota$ be given. Assume the following.

$$c_2Ehurat_2Ehurat_REP_CLASS \in ((2^{(ty_2Epair_2Eprod\ ty_2Enum_2Enum\ ty_2Enum_2Enum)})^{ty_2Ehurat_2Ehurat}) \tag{6}$$

Definition 6 We define $c_2Emin_2E_40$ to be $\lambda A.\lambda P \in 2^A$. **if** $(\exists x \in A.p (ap P x))$ **then** $(the (\lambda x.x \in A \wedge p$
of type $\iota \Rightarrow \iota$.

Definition 7 We define $c_2Ehrat_2Ehrat_REP$ to be $\lambda V0a \in ty_2Ehrat_2Ehrat.(ap (c_2Emin_2E_40 (ty_2E$

Let $c_2Ehrat_2Etrrat_add : \iota$ be given. Assume the following.

$$c_2Ehrat_2Etrrat_add \in (((ty_2Epair_2Eprod ty_2Enum_2Enum ty_2Enum_2Enum)^{ty_2Epair_2Eprod ty_2Enum_2Enum})$$
 (7)

Let $c_2Ehrat_2Etrrat_eq : \iota$ be given. Assume the following.

$$c_2Ehrat_2Etrrat_eq \in ((2^{(ty_2Epair_2Eprod ty_2Enum_2Enum ty_2Enum_2Enum)})^{ty_2Epair_2Eprod ty_2Enum_2Enum})$$
 (8)

Let $c_2Ehrat_2Ehrat_ABS_CLASS : \iota$ be given. Assume the following.

$$c_2Ehrat_2Ehrat_ABS_CLASS \in (ty_2Ehrat_2Ehrat^{(2^{(ty_2Epair_2Eprod ty_2Enum_2Enum ty_2Enum_2Enum)})})$$
 (9)

Definition 8 We define $c_2Ehrat_2Ehrat_ABS$ to be $\lambda V0r \in (ty_2Epair_2Eprod ty_2Enum_2Enum ty_2E$

Definition 9 We define $c_2Ehrat_2Ehrat_add$ to be $\lambda V0T1 \in ty_2Ehrat_2Ehrat.\lambda V1T2 \in ty_2Ehrat_2Eh$

Definition 10 We define $c_2Ebool_2E_3F$ to be $\lambda A_27a : \iota.(\lambda V0P \in (2^{A_27a}).(ap V0P (ap (c_2Emin_2E_40$

Let $c_2Ehreal_2Ehreal : \iota$ be given. Assume the following.

$$c_2Ehreal_2Ehreal \in (ty_2Ehreal_2Ehreal^{(2^{ty_2Ehrat_2Ehrat})})$$
 (10)

Definition 11 We define $c_2Ehreal_2Ehreal_add$ to be $\lambda V0X \in ty_2Ehreal_2Ehreal.\lambda V1Y \in ty_2Ehreal_2Eh$

Definition 12 We define $c_2Ebool_2E_7E$ to be $(ap (c_2Ebool_2E_21 2) (\lambda V0t \in 2.V0t))$.

Definition 13 We define $c_2Ebool_2E_7E$ to be $(\lambda V0t \in 2.(ap (ap c_2Emin_2E_3D_3D_3E V0t) c_2Ebool_2E_7E$

Assume the following.

$$\forall A_27a.nonempty A_27a \Rightarrow (\forall V0x \in A_27a.(\forall V1y \in A_27a.((V0x = V1y) \Leftrightarrow (V1y = V0x))))$$
 (11)

Assume the following.

$$(\forall V0x \in ty_2Ehreal_2Ehreal.(\forall V1y \in ty_2Ehreal_2Ehreal.(\neg((ap (ap c_2Ehreal_2Ehreal_add V0x) V1y) = V0x))))$$
 (12)

Theorem 1

$$(\forall V0x \in ty_2Ehreal_2Ehreal.(\forall V1y \in ty_2Ehreal_2Ehreal.(\neg(V0x = (ap (ap c_2Ehreal_2Ehreal_add V0x) V1y))))))$$