

thm_2Erelation_2ENOT__INVOL
(TMNX4wYYQScU7YwPvgkGSVpGZNhYTvp9iuA)

October 26, 2020

Definition 1 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define $c_2Ebool_2E_2ET$ to be $(ap (ap (c_2Emin_2E_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

Definition 3 We define $c_2Ebool_2E_21$ to be $\lambda A_27a : \iota.(\lambda V0P \in (2^{A_27a}).(ap (ap (c_2Emin_2E_3D (2^{A_27a}))$

Definition 4 We define $c_2Ebool_2E_2EF$ to be $(ap (c_2Ebool_2E_21 2) (\lambda V0t \in 2.V0t))$.

Definition 5 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p P \Rightarrow p Q)$ of type ι .

Definition 6 We define $c_2Ebool_2E_27E$ to be $(\lambda V0t \in 2.(ap (ap c_2Emin_2E_3D_3D_3E V0t) c_2Ebool_2E_2EF$

Definition 7 We define $c_2Ebool_2E_2F_5C$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21 2) (\lambda V2t \in 2.V2t))$

Definition 8 We define $c_2Ecombin_2EK$ to be $\lambda A_27a : \iota.\lambda A_27b : \iota.(\lambda V0x \in A_27a.(\lambda V1y \in A_27b.V0x))$

Definition 9 We define $c_2Ecombin_2ES$ to be $\lambda A_27a : \iota.\lambda A_27b : \iota.\lambda A_27c : \iota.(\lambda V0f \in ((A_27c^{A_27b})^{A_27a})$

Definition 10 We define $c_2Ecombin_2EI$ to be $\lambda A_27a : \iota.(ap (ap (c_2Ecombin_2ES A_27a (A_27a^{A_27a}) A_27a$

Definition 11 We define $c_2Ecombin_2Eo$ to be $\lambda A_27a : \iota.\lambda A_27b : \iota.\lambda A_27c : \iota.\lambda V0f \in (A_27b^{A_27c}).\lambda V1g \in (A_27c^{A_27a}).$

Definition 12 We define $c_2Erelation_2EINVOL$ to be $\lambda A_27z : \iota.\lambda V0f \in (A_27z^{A_27z}).(ap (ap (c_2Emin_2E_3D$

Assume the following.

$$True \tag{1}$$

Assume the following.

$$\forall A_27a.nonempty A_27a \Rightarrow (\forall V0t \in 2.((\forall V1x \in A_27a.(p V0t)) \Leftrightarrow (p V0t))) \tag{2}$$

Assume the following.

$$((\forall V0t \in 2.((\neg(\neg(p V0t))) \Leftrightarrow (p V0t))) \wedge (((\neg True) \Leftrightarrow False) \wedge ((\neg False) \Leftrightarrow True))) \tag{3}$$

Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow (\forall V0x \in A_27a. ((V0x = V0x) \Leftrightarrow True)) \quad (4)$$

Assume the following.

$$\forall A_27z.nonempty\ A_27z \Rightarrow (\forall V0f \in (A_27z^{A_27z}). ((p (ap (c_2Erelation_2EINVOL\ A_27z)\ V0f)) \Leftrightarrow (\forall V1x \in A_27z. (ap\ V0f\ (ap\ V0f\ V1x)) = V1x)))) \quad (5)$$

Theorem 1 $(p (ap (c_2Erelation_2EINVOL\ 2)\ c_2Ebool_2E_7E)).$