

thm_2Erelation_2EO_Id

(TMY67jcFk4kWRBWKDos4JnFpBLck72ZDrMK)

October 26, 2020

Definition 1 We define `c_2Emin_2E_3D_3D_3E` to be $\lambda P \in 2. \lambda Q \in 2. \text{inj_o } (p \Rightarrow P \Rightarrow Q)$ of type ι .

Definition 2 We define `c_2Emin_2E_3D` to be $\lambda A. \lambda x \in A. \lambda y \in A. \text{inj_o } (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 3 We define `c_2Ebool_2E_2T` to be $(\text{ap } (\text{ap } (\text{c_2Emin_2E_3D } (2^2)) (\lambda V 0x \in 2. V 0x)) (\lambda V 1x \in 2. V 1x))$

Definition 4 We define `c_2Ebool_2E_21` to be $\lambda A. 27a : \iota. (\lambda V 0P \in (2^{A-27a}). (\text{ap } (\text{ap } (\text{c_2Emin_2E_3D } (2^{A-27a}))))$

Definition 5 We define `c_2Ebool_2E_2F_5C` to be $(\lambda V 0t1 \in 2. (\lambda V 1t2 \in 2. (\text{ap } (\text{c_2Ebool_2E_21 } 2)) (\lambda V 2t \in 2. V 2t)))$

Definition 6 We define `c_2Emin_2E_40` to be $\lambda A. \lambda P \in 2^A. \text{if } (\exists x \in A. p (\text{ap } P x)) \text{ then } (\text{the } (\lambda x. x \in A \wedge p (\text{ap } P x)))$ of type $\iota \Rightarrow \iota$.

Definition 7 We define `c_2Ebool_2E_3F` to be $\lambda A. 27a : \iota. (\lambda V 0P \in (2^{A-27a}). (\text{ap } V 0P (\text{ap } (\text{c_2Emin_2E_40 } A))))$

Definition 8 We define `c_2Erelation_2EO` to be $\lambda A. 27g : \iota. \lambda A. 27h : \iota. \lambda A. 27k : \iota. \lambda V 0R1 \in ((2^{A-27k})^{A-27h}).$

Assume the following.

$$\text{True} \tag{1}$$

Assume the following.

$$(\forall V 0t1 \in 2. (\forall V 1t2 \in 2. (((p \Rightarrow V 0t1) \Rightarrow (p \Rightarrow V 1t2)) \Rightarrow (((p \Rightarrow V 1t2) \Rightarrow (p \Rightarrow V 0t1)) \Rightarrow ((p \Rightarrow V 0t1) \Leftrightarrow (p \Rightarrow V 1t2)))))) \tag{2}$$

Assume the following.

$$\forall A. 27a. \text{nonempty } A. 27a \Rightarrow (\forall V 0t \in 2. ((\forall V 1x \in A. 27a. (p \Rightarrow V 0t)) \Leftrightarrow (p \Rightarrow V 0t))) \tag{3}$$

Assume the following.

$$\forall A. 27a. \text{nonempty } A. 27a \Rightarrow (\forall V 0x \in A. 27a. ((V 0x = V 0x) \Leftrightarrow \text{True})) \tag{4}$$

Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow (\forall V0x \in A_27a. (\forall V1y \in A_27a. ((V0x = V1y) \Leftrightarrow (V1y = V0x)))) \quad (5)$$

Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow \forall A_27b.nonempty\ A_27b \Rightarrow (\forall V0f \in (A_27b^{A_27a}). (\forall V1g \in (A_27b^{A_27a}). ((V0f = V1g) \Leftrightarrow (\forall V2x \in A_27a. ((ap\ V0f\ V2x) = (ap\ V1g\ V2x)))))) \quad (6)$$

Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow (\forall V0P \in (2^{A_27a}). (\forall V1a \in A_27a. ((\exists V2x \in A_27a. ((V2x = V1a) \wedge (p\ (ap\ V0P\ V2x)))) \Leftrightarrow (p\ (ap\ V0P\ V1a)))))) \quad (7)$$

Theorem 1

$$\forall A_27a.nonempty\ A_27a \Rightarrow \forall A_27b.nonempty\ A_27b \Rightarrow (\forall V0R \in ((2^{A_27b})^{A_27a}). ((ap\ (ap\ (c_2Erelation_2EO\ A_27a\ A_27a\ A_27b)\ V0R)\ (c_2Emin_2E_3D\ A_27a)) = V0R))$$