

thm\_2Eres\_\_quan\_2ERES\_\_EXISTS\_\_DISJ\_\_DIST  
 (TMThgLCnfEbvsCDGss-  
 doWNkkNVV3AHzLTbW)

October 26, 2020

**Definition 1** We define  $c\_2Emin\_2E\_3D\_3D\_3E$  to be  $\lambda P \in 2.\lambda Q \in 2.inj\_o (p P \Rightarrow p Q)$  of type  $\iota$ .

**Definition 2** We define  $c\_2Emin\_2E\_3D$  to be  $\lambda A.\lambda x \in A.\lambda y \in A.inj\_o (x = y)$  of type  $\iota \Rightarrow \iota$ .

**Definition 3** We define  $c\_2Ebool\_2E\_2T$  to be  $(ap (ap (c\_2Emin\_2E\_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

**Definition 4** We define  $c\_2Ebool\_2E\_21$  to be  $\lambda A\_27a : \iota.(\lambda V0P \in (2^{A\_27a}).(ap (ap (c\_2Emin\_2E\_3D (2^{A\_27a}))$

**Definition 5** We define  $c\_2Ebool\_2E\_5C\_2F$  to be  $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c\_2Ebool\_2E\_21 2) (\lambda V2t \in 2.V2t)))$

**Definition 6** We define  $c\_2Ebool\_2E\_IN$  to be  $\lambda A\_27a : \iota.(\lambda V0x \in A\_27a.(\lambda V1f \in (2^{A\_27a}).(ap V1f V0x)))$

**Definition 7** We define  $c\_2Ebool\_2E\_2F\_5C$  to be  $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c\_2Ebool\_2E\_21 2) (\lambda V2t \in 2.V2t)))$

**Definition 8** We define  $c\_2Emin\_2E\_40$  to be  $\lambda A.\lambda P \in 2^A$ .if  $(\exists x \in A.p (ap P x))$  then  $(the (\lambda x.x \in A \wedge p (ap P x)))$  of type  $\iota \Rightarrow \iota$ .

**Definition 9** We define  $c\_2Ebool\_2E\_3F$  to be  $\lambda A\_27a : \iota.(\lambda V0P \in (2^{A\_27a}).(ap V0P (ap (c\_2Emin\_2E\_40 A\_27a P))))$

**Definition 10** We define  $c\_2Ebool\_2ERES\_EXISTS$  to be  $\lambda A\_27a : \iota.(\lambda V0p \in (2^{A\_27a}).(\lambda V1m \in (2^{A\_27a}).(ap V1m (ap (c\_2Emin\_2E\_40 A\_27a p))))$

Assume the following.

$$(\forall V0t1 \in 2.(\forall V1t2 \in 2.(((p V0t1) \wedge (p V1t2)) \Leftrightarrow ((p V1t2) \wedge (p V0t1)))))) \quad (1)$$

Assume the following.

$$\begin{aligned} & \forall A\_27a.nonempty A\_27a \Rightarrow (\forall V0P \in (2^{A\_27a}).(\forall V1Q \in \\ & (2^{A\_27a}).((\exists V2x \in A\_27a.((p (ap V0P V2x)) \vee (p (ap V1Q V2x)))))) \Leftrightarrow \\ & ((\exists V3x \in A\_27a.(p (ap V0P V3x))) \vee (\exists V4x \in A\_27a.(p (ap V1Q V4x)))))) \end{aligned} \quad (2)$$

Assume the following.

$$(\forall V0A \in 2. (\forall V1B \in 2. (\forall V2C \in 2. (((p V1B) \vee (p V2C)) \wedge (p V0A)) \Leftrightarrow (((p V1B) \wedge (p V0A)) \vee ((p V2C) \wedge (p V0A)))))) \quad (3)$$

Assume the following.

$$\forall A\_27a. \text{nonempty } A\_27a \Rightarrow (\forall V0P \in (2^{A\_27a}). (\forall V1x \in A\_27a. ((p (ap (ap (c\_2Ebool\_2EIN A\_27a) V1x) V0P)) \Leftrightarrow (p (ap V0P V1x)))))) \quad (4)$$

Assume the following.

$$\forall A\_27a. \text{nonempty } A\_27a \Rightarrow (\forall V0P \in (2^{A\_27a}). (\forall V1f \in (2^{A\_27a}). ((p (ap (ap (c\_2Ebool\_2ERES\_EXISTS A\_27a) V0P) V1f)) \Leftrightarrow (\exists V2x \in A\_27a. ((p (ap (ap (c\_2Ebool\_2EIN A\_27a) V2x) V0P)) \wedge (p (ap V1f V2x)))))))) \quad (5)$$

**Theorem 1**

$$\forall A\_27a. \text{nonempty } A\_27a \Rightarrow (\forall V0P \in (2^{A\_27a}). (\forall V1Q \in (2^{A\_27a}). (\forall V2R \in (2^{A\_27a}). ((p (ap (ap (c\_2Ebool\_2ERES\_EXISTS A\_27a) V0P) (\lambda V3i \in A\_27a. (ap (ap c\_2Ebool\_2E\_5C\_2F (ap V1Q V3i)) (ap V2R V3i)))))) \Leftrightarrow ((p (ap (ap (c\_2Ebool\_2ERES\_EXISTS A\_27a) V0P) (\lambda V4i \in A\_27a. (ap V1Q V4i)))) \vee (p (ap (ap (c\_2Ebool\_2ERES\_EXISTS A\_27a) V0P) (\lambda V5i \in A\_27a. (ap V2R V5i))))))))))$$