

thm_2Eres__quan_2ERES__EXISTS__UNIQUE__NOT__EMPTY (TMT3sz3AdQu41a7L98kb8ykTKQ36Zih51WL)

October 26, 2020

Definition 1 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$
of type $\iota \Rightarrow \iota$.

Definition 2 We define $c_2Ebool_2E_2T$ to be $(ap (ap (c_2Emin_2E_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

Definition 3 We define $c_2Ebool_2E_21$ to be $\lambda A_27a : \iota.(\lambda V0P \in (2^{A_27a}).(ap (ap (c_2Emin_2E_3D (2^{A_27a}))$

Definition 4 We define $c_2Ebool_2E_2F$ to be $(ap (c_2Ebool_2E_21 2) (\lambda V0t \in 2.V0t))$.

Definition 5 We define $c_2Epred_set_2EEMPTY$ to be $\lambda A_27a : \iota.(\lambda V0x \in A_27a.c_2Ebool_2E_2F)$.

Definition 6 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p P \Rightarrow p Q)$
of type ι .

Definition 7 We define $c_2Ebool_2E_7E$ to be $(\lambda V0t \in 2.(ap (ap c_2Emin_2E_3D_3D_3E V0t) c_2Ebool_2E_2F))$

Definition 8 We define $c_2Ebool_2E_2IN$ to be $\lambda A_27a : \iota.(\lambda V0x \in A_27a.(\lambda V1f \in (2^{A_27a}).(ap V1f V0x)))$

Definition 9 We define $c_2Ebool_2E_2F_5C$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21 2) (\lambda V2t \in 2.V2t)))$

Definition 10 We define $c_2Emin_2E_40$ to be $\lambda A.\lambda P \in 2^A.if (\exists x \in A.p (ap P x)) \mathbf{then} (the (\lambda x.x \in A))$
of type $\iota \Rightarrow \iota$.

Definition 11 We define $c_2Ebool_2E_3F$ to be $\lambda A_27a : \iota.(\lambda V0P \in (2^{A_27a}).(ap V0P (ap (c_2Emin_2E_40$

Definition 12 We define $c_2Ebool_2ERES_EXISTS$ to be $\lambda A_27a : \iota.(\lambda V0p \in (2^{A_27a}).(\lambda V1m \in (2^{A_27a}).(ap$

Definition 13 We define $c_2Ebool_2ERES_FORALL$ to be $\lambda A_27a : \iota.(\lambda V0p \in (2^{A_27a}).(\lambda V1m \in (2^{A_27a}).(ap$

Definition 14 We define $c_2Ebool_2ERES_EXISTS_UNIQUE$ to be $\lambda A_27a : \iota.(\lambda V0p \in (2^{A_27a}).(\lambda V1m \in$

Assume the following.

$$\forall A.27a.nonempty\ A.27a \Rightarrow (\forall V0P \in (2^{A.27a}). (\forall V1s \in (2^{A.27a}). ((p\ (ap\ (ap\ (c.2Ebool.2ERES_EXISTS\ A.27a)\ V1s)\ V0P)) \Rightarrow (\neg(V1s = (c.2Epred_set.2EEMPTY\ A.27a)))))) \quad (1)$$

Assume the following.

$$\forall A.27a.nonempty\ A.27a \Rightarrow (\forall V0P \in (2^{A.27a}). (\forall V1s \in (2^{A.27a}). ((p\ (ap\ (ap\ (c.2Ebool.2ERES_EXISTS_UNIQUE\ A.27a)\ V0P)\ V1s)) \Rightarrow (p\ (ap\ (ap\ (c.2Ebool.2ERES_EXISTS\ A.27a)\ V0P)\ V1s)))))) \quad (2)$$

Theorem 1

$$\forall A.27a.nonempty\ A.27a \Rightarrow (\forall V0P \in (2^{A.27a}). (\forall V1s \in (2^{A.27a}). ((p\ (ap\ (ap\ (c.2Ebool.2ERES_EXISTS_UNIQUE\ A.27a)\ V1s)\ V0P)) \Rightarrow (\neg(V1s = (c.2Epred_set.2EEMPTY\ A.27a))))))$$