

thm_2Eres__quan_2ERES__EXISTS__UNIQUE__SING (TMYZpfW3kULPKbniimS9heVBdoSgZZkez2F)

October 26, 2020

Definition 1 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$
of type $\iota \Rightarrow \iota$.

Definition 2 We define c_2Ebool_2ET to be $(ap (ap (c_2Emin_2E_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

Definition 3 We define $c_2Ebool_2E_21$ to be $\lambda A_27a : \iota.(\lambda V0P \in (2^{A_27a}).(ap (ap (c_2Emin_2E_3D (2^{A_27a})))$

Definition 4 We define c_2Ebool_2EF to be $(ap (c_2Ebool_2E_21 2) (\lambda V0t \in 2.V0t))$.

Definition 5 We define $c_2Epred_set_2EEMPTY$ to be $\lambda A_27a : \iota.(\lambda V0x \in A_27a.c_2Ebool_2EF)$.

Definition 6 We define c_2Ebool_2EIN to be $\lambda A_27a : \iota.(\lambda V0x \in A_27a.(\lambda V1f \in (2^{A_27a}).(ap V1f V0x)))$

Definition 7 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p P \Rightarrow p Q)$
of type ι .

Definition 8 We define $c_2Ebool_2E_5C_2F$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21 2) (\lambda V2t \in 2.V2t)))$

Definition 9 We define $c_2Ebool_2E_2F_5C$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21 2) (\lambda V2t \in 2.V2t)))$

Let $ty_2Epair_2Eprod : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A0.nonempty A0 \Rightarrow \forall A1.nonempty A1 \Rightarrow nonempty (ty_2Epair_2Eprod A0 A1) \quad (1)$$

Let $c_2Epair_2EABS_prod : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A_27a.nonempty A_27a \Rightarrow \forall A_27b.nonempty A_27b \Rightarrow c_2Epair_2EABS_prod A_27a A_27b \in ((ty_2Epair_2Eprod A_27a A_27b)^{(2^{A_27b})^{A_27a}}) \quad (2)$$

Definition 10 We define $c_2Epair_2E_2C$ to be $\lambda A_27a : \iota.\lambda A_27b : \iota.\lambda V0x \in A_27a.\lambda V1y \in A_27b.(ap (c_2Emin_2E_3D (2^{A_27a})))$

Let $c_2Epred_set_2EGSPEC : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned} \forall A_27a.nonempty\ A_27a \Rightarrow \forall A_27b.nonempty\ A_27b \Rightarrow c_2Epred_set_2EGSPEC \\ A_27a\ A_27b \in ((2^{A_27a})^{((ty_2Epair_2Eprod\ A_27a\ 2)^{A_27b})}) \end{aligned} \quad (3)$$

Definition 11 We define $c_2Epred_set_2EINSERT$ to be $\lambda A_27a : \iota. \lambda V0x \in A_27a. \lambda V1s \in (2^{A_27a}). (ap\ (c_2Emin_2E40$

Definition 12 We define c_2Emin_2E40 to be $\lambda A. \lambda P \in 2^A. \text{if } (\exists x \in A. p\ (ap\ P\ x)) \text{ then } (the\ (\lambda x. x \in A \wedge p\ x))$ of type $\iota \Rightarrow \iota$.

Definition 13 We define c_2Ebool_2E3F to be $\lambda A_27a : \iota. (\lambda V0P \in (2^{A_27a}). (ap\ V0P\ (ap\ (c_2Emin_2E40$

Definition 14 We define $c_2Ebool_2ERES_FORALL$ to be $\lambda A_27a : \iota. (\lambda V0p \in (2^{A_27a}). (\lambda V1m \in (2^{A_27a}).$

Definition 15 We define $c_2Ebool_2ERES_EXISTS$ to be $\lambda A_27a : \iota. (\lambda V0p \in (2^{A_27a}). (\lambda V1m \in (2^{A_27a}).$

Definition 16 We define $c_2Ebool_2ERES_EXISTS_UNIQUE$ to be $\lambda A_27a : \iota. (\lambda V0p \in (2^{A_27a}). (\lambda V1m \in$

Assume the following.

$$True \quad (4)$$

Assume the following.

$$\begin{aligned} \forall A_27a.nonempty\ A_27a \Rightarrow (\forall V0t \in 2. ((\forall V1x \in \\ A_27a. (p\ V0t)) \Leftrightarrow (p\ V0t))) \end{aligned} \quad (5)$$

Assume the following.

$$\begin{aligned} (\forall V0t \in 2. (((True \wedge (p\ V0t)) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \wedge True) \Leftrightarrow \\ (p\ V0t)) \wedge (((False \wedge (p\ V0t)) \Leftrightarrow False) \wedge (((p\ V0t) \wedge False) \Leftrightarrow False) \wedge \\ (((p\ V0t) \wedge (p\ V0t)) \Leftrightarrow (p\ V0t)))))) \end{aligned} \quad (6)$$

Assume the following.

$$\begin{aligned} \forall A_27a.nonempty\ A_27a \Rightarrow (\forall V0x \in A_27a. ((V0x = V0x) \Leftrightarrow \\ True)) \end{aligned} \quad (7)$$

Assume the following.

$$\begin{aligned} \forall A_27a.nonempty\ A_27a \Rightarrow (\forall V0p \in (2^{A_27a}). (\forall V1m \in \\ 2. ((p\ (ap\ (ap\ (c_2Ebool_2ERES_EXISTS_UNIQUE\ A_27a)\ V0p)\ (\lambda V2x \in \\ A_27a. V1m))) \Leftrightarrow ((\exists V3x \in A_27a. (V0p = (ap\ (ap\ (c_2Epred_set_2EINSERT \\ A_27a)\ V3x)\ (c_2Epred_set_2EEMPTY\ A_27a)))) \wedge (p\ V1m)))))) \end{aligned} \quad (8)$$

Theorem 1

$$\begin{aligned} \forall A_27a.nonempty\ A_27a \Rightarrow \forall A_27b.nonempty\ A_27b \Rightarrow \forall A_27c. \\ nonempty\ A_27c \Rightarrow (\forall V0P \in A_27a. (\forall V1s \in (2^{A_27b}). \\ \forall V2x \in A_27c. ((p\ (ap\ (ap\ (c_2Ebool_2ERES_EXISTS_UNIQUE \\ A_27b)\ V1s)\ (\lambda V3x \in A_27b. c_2Ebool_2ET)))) \Leftrightarrow (\exists V4y \in A_27b. \\ (V1s = (ap\ (ap\ (c_2Epred_set_2EINSERT\ A_27b)\ V4y)\ (c_2Epred_set_2EEMPTY \\ A_27b))))))))) \end{aligned}$$