

thm_2Eres__quan_2ERES__EXISTS__UNIV
(TMavJjHcGKr2DaCS8sk8L3Y2KVvnbkuuDuD)

October 26, 2020

Definition 1 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define $c_2Ebool_2E_21$ to be $(ap (ap (c_2Emin_2E_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

Definition 3 We define $c_2Ebool_2E_21$ to be $\lambda A_27a : \iota.(\lambda V0P \in (2^{A-27a}).(ap (ap (c_2Emin_2E_3D (2^{A-27a})) (\lambda V1f \in (2^{A-27a}).(ap V1f V0x))) (\lambda V2t \in 2.V2t)))$

Definition 4 We define $c_2Ebool_2E_21$ to be $(ap (c_2Ebool_2E_21 2) (\lambda V0t \in 2.V0t))$.

Definition 5 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p P \Rightarrow p Q)$ of type ι .

Definition 6 We define $c_2Ebool_2E_7E$ to be $(\lambda V0t \in 2.(ap (ap c_2Emin_2E_3D_3D_3E V0t) c_2Ebool_2E_21))$

Definition 7 We define $c_2Epred_set_2EUNIV$ to be $\lambda A_27a : \iota.(\lambda V0x \in A_27a.c_2Ebool_2E_21)$.

Definition 8 We define $c_2Ebool_2E_IN$ to be $\lambda A_27a : \iota.(\lambda V0x \in A_27a.(\lambda V1f \in (2^{A-27a}).(ap V1f V0x)))$

Definition 9 We define $c_2Ebool_2E_2F_5C$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21 2) (\lambda V2t \in 2.V2t)))$

Definition 10 We define $c_2Emin_2E_40$ to be $\lambda A.\lambda P \in 2^A.if (\exists x \in A.p (ap P x)) \text{ then } (the (\lambda x.x \in A \wedge p (ap P x)))$ of type $\iota \Rightarrow \iota$.

Definition 11 We define $c_2Ebool_2E_3F$ to be $\lambda A_27a : \iota.(\lambda V0P \in (2^{A-27a}).(ap V0P (ap (c_2Emin_2E_40 (2^{A-27a})) (\lambda V1m \in (2^{A-27a}).(ap V1m V0P))))$

Definition 12 We define $c_2Ebool_2ERES_EXISTS$ to be $\lambda A_27a : \iota.(\lambda V0p \in (2^{A-27a}).(\lambda V1m \in (2^{A-27a}).(ap V1m V0p)))$

Assume the following.

$$True \tag{1}$$

Assume the following.

$$\begin{aligned} & (\forall V0t \in 2.(((True \wedge (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \wedge True) \Leftrightarrow \\ & (p V0t)) \wedge (((False \wedge (p V0t)) \Leftrightarrow False) \wedge (((p V0t) \wedge False) \Leftrightarrow False) \wedge \\ & (((p V0t) \wedge (p V0t)) \Leftrightarrow (p V0t)))))) \end{aligned} \tag{2}$$

Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow (\forall V0x \in A_27a. ((V0x = V0x) \Leftrightarrow True)) \quad (3)$$

Assume the following.

$$\begin{aligned} & (\forall V0t \in 2. (((True \Leftrightarrow (p\ V0t)) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \Leftrightarrow True) \Leftrightarrow \\ & (p\ V0t)) \wedge (((False \Leftrightarrow (p\ V0t)) \Leftrightarrow \neg(p\ V0t)) \wedge (((p\ V0t) \Leftrightarrow False) \Leftrightarrow \neg(\\ & p\ V0t)))))) \end{aligned} \quad (4)$$

Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow (\forall V0x \in A_27a. (p\ (ap\ (ap\ (c_2Ebool_2EIN\ A_27a)\ V0x)\ (c_2Epred_set_2EUNIV\ A_27a)))) \quad (5)$$

Assume the following.

$$\begin{aligned} & \forall A_27a.nonempty\ A_27a \Rightarrow (\forall V0P \in (2^{A_27a}). (\forall V1f \in \\ & (2^{A_27a}). ((p\ (ap\ (ap\ (c_2Ebool_2ERES_EXISTS\ A_27a)\ V0P)\ V1f)) \Leftrightarrow \\ & (\exists V2x \in A_27a. ((p\ (ap\ (ap\ (c_2Ebool_2EIN\ A_27a)\ V2x)\ V0P)) \wedge \\ & (p\ (ap\ V1f\ V2x)))))) \end{aligned} \quad (6)$$

Theorem 1

$$\begin{aligned} & \forall A_27a.nonempty\ A_27a \Rightarrow (\forall V0p \in (2^{A_27a}). ((p\ (ap \\ & (ap\ (c_2Ebool_2ERES_EXISTS\ A_27a)\ (c_2Epred_set_2EUNIV\ A_27a)) \\ & V0p)) \Leftrightarrow (p\ (ap\ (c_2Ebool_2E_3F\ A_27a)\ V0p)))) \end{aligned}$$