

# thm\_2Erich\_list\_2EEVERY\_FOLDR (TMGVLrD4htUdWaLofo3ESZT0i97vHkeMYNj)

October 26, 2020

**Definition 1** We define  $c\_2Emin\_2E\_3D$  to be  $\lambda A.\lambda x \in A.\lambda y \in A.inj\_o (x = y)$  of type  $\iota \Rightarrow \iota$ .

**Definition 2** We define  $c\_2Ebool\_2ET$  to be  $(ap (ap (c\_2Emin\_2E\_3D (2^2))) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x)$

**Definition 3** We define  $c\_2Ebool\_2E\_21$  to be  $\lambda A\_27a : \iota.(\lambda V0P \in (2^{A\_27a}).(ap (ap (c\_2Emin\_2E\_3D (2^{A\_27a}))))$

**Definition 4** We define  $c\_2Ebool\_2EF$  to be  $(ap (c\_2Ebool\_2E\_21 2) (\lambda V0t \in 2.V0t))$ .

**Definition 5** We define  $c\_2Emin\_2E\_3D\_3D\_3E$  to be  $\lambda P \in 2.\lambda Q \in 2.inj\_o (p P \Rightarrow p Q)$  of type  $\iota$ .

**Definition 6** We define  $c\_2Ebool\_2E\_7E$  to be  $(\lambda V0t \in 2.(ap (ap c\_2Emin\_2E\_3D\_3D\_3E V0t) c\_2Ebool\_2EF))$

Let  $ty\_2Elist\_2Elist : \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A0.nonempty A0 \Rightarrow nonempty (ty\_2Elist\_2Elist A0) \quad (1)$$

Let  $c\_2Elist\_2EFOLDR : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\begin{aligned} \forall A\_27a.nonempty A\_27a \Rightarrow \forall A\_27b.nonempty A\_27b \Rightarrow c\_2Elist\_2EFOLDR \\ A\_27a A\_27b \in (((A\_27b^{(ty\_2Elist\_2Elist A\_27a)})^{A\_27b})^{((A\_27b^{A\_27b})^{A\_27a})}) \end{aligned} \quad (2)$$

Let  $c\_2Elist\_2EEVERY : \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A\_27a.nonempty A\_27a \Rightarrow c\_2Elist\_2EEVERY A\_27a \in ((2^{(ty\_2Elist\_2Elist A\_27a)})^{(2^{A\_27a})}) \quad (3)$$

Let  $c\_2Elist\_2ECONS : \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A\_27a.nonempty A\_27a \Rightarrow c\_2Elist\_2ECONS A\_27a \in (((ty\_2Elist\_2Elist A\_27a)^{(ty\_2Elist\_2Elist A\_27a)})^{A\_27a}) \quad (4)$$

Let  $c\_2Elist\_2ENIL : \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A\_27a.nonempty A\_27a \Rightarrow c\_2Elist\_2ENIL A\_27a \in (ty\_2Elist\_2Elist A\_27a) \quad (5)$$

**Definition 7** We define  $c\_2Ebool\_2E\_2F\_5C$  to be  $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c\_2Ebool\_2E\_21 2) (\lambda V2t \in 2.$

Assume the following.

$$True \tag{6}$$

Assume the following.

$$\forall A\_27a.nonempty A\_27a \Rightarrow (\forall V0t \in 2.((\forall V1x \in A\_27a.(p V0t)) \Leftrightarrow (p V0t))) \tag{7}$$

Assume the following.

$$\forall A\_27a.nonempty A\_27a \Rightarrow (\forall V0x \in A\_27a.((V0x = V0x) \Leftrightarrow True)) \tag{8}$$

Assume the following.

$$(\forall V0t \in 2.(((True \Leftrightarrow (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \Leftrightarrow True) \Leftrightarrow (p V0t)) \wedge (((False \Leftrightarrow (p V0t)) \Leftrightarrow \neg(p V0t)) \wedge (((p V0t) \Leftrightarrow False) \Leftrightarrow \neg(p V0t)))))) \tag{9}$$

Assume the following.

$$\begin{aligned} & \forall A\_27a.nonempty A\_27a \Rightarrow \forall A\_27b.nonempty A\_27b \Rightarrow ( \\ & (\forall V0f \in ((A\_27b^{A\_27b})^{A\_27a}).(\forall V1e \in A\_27b.(ap ( \\ & ap (ap (c\_2Elist\_2EFOLDR A\_27a A\_27b) V0f) V1e) (c\_2Elist\_2ENIL \\ & A\_27a)) = V1e))) \wedge (\forall V2f \in ((A\_27b^{A\_27b})^{A\_27a}).(\forall V3e \in \\ & A\_27b.(\forall V4x \in A\_27a.(\forall V5l \in (ty\_2Elist\_2Elist A\_27a). \\ & ((ap (ap (ap (c\_2Elist\_2EFOLDR A\_27a A\_27b) V2f) V3e) (ap (ap (c\_2Elist\_2ECONS \\ & A\_27a) V4x) V5l)) = (ap (ap V2f V4x) (ap (ap (ap (c\_2Elist\_2EFOLDR \\ & A\_27a A\_27b) V2f) V3e) V5l))))))))) \end{aligned} \tag{10}$$

Assume the following.

$$\begin{aligned} & \forall A\_27a.nonempty A\_27a \Rightarrow ((\forall V0P \in (2^{A\_27a}).((p (ap \\ & (ap (c\_2Elist\_2EVERY A\_27a) V0P) (c\_2Elist\_2ENIL A\_27a))) \Leftrightarrow True)) \wedge \\ & (\forall V1P \in (2^{A\_27a}).(\forall V2h \in A\_27a.(\forall V3t \in (ty\_2Elist\_2Elist \\ & A\_27a).((p (ap (ap (c\_2Elist\_2EVERY A\_27a) V1P) (ap (ap (c\_2Elist\_2ECONS \\ & A\_27a) V2h) V3t))) \Leftrightarrow ((p (ap V1P V2h)) \wedge (p (ap (ap (c\_2Elist\_2EVERY \\ & A\_27a) V1P) V3t)))))))))) \end{aligned} \tag{11}$$

Assume the following.

$$\begin{aligned} & \forall A\_27a.nonempty A\_27a \Rightarrow (\forall V0P \in (2^{(ty\_2Elist\_2Elist A\_27a)}). \\ & (((p (ap V0P (c\_2Elist\_2ENIL A\_27a))) \wedge (\forall V1t \in (ty\_2Elist\_2Elist \\ & A\_27a).((p (ap V0P V1t)) \Rightarrow (\forall V2h \in A\_27a.(p (ap V0P (ap (ap ( \\ & c\_2Elist\_2ECONS A\_27a) V2h) V1t)))))) \Rightarrow (\forall V3l \in (ty\_2Elist\_2Elist \\ & A\_27a).(p (ap V0P V3l)))))) \end{aligned} \tag{12}$$

**Theorem 1**

$$\begin{aligned} & \forall A_{.27a}. \text{nonempty } A_{.27a} \Rightarrow (\forall V0P \in (2^{A_{.27a}}). (\forall V1l \in \\ & (ty\_2Elist\_2Elist \ A_{.27a}). ((p \ (ap \ (ap \ (c\_2Elist\_2EVERY \ A_{.27a}) \\ V0P) \ V1l)) \Leftrightarrow (p \ (ap \ (ap \ (ap \ (c\_2Elist\_2EFOLDER \ A_{.27a} \ 2) \ (\lambda V2x \in \\ & A_{.27a}. (\lambda V3l_{.27} \in 2. (ap \ (ap \ c\_2Ebool\_2E\_2F\_5C \ (ap \ V0P \ V2x)) \\ & \ V3l_{.27})))) \ c\_2Ebool\_2ET) \ V1l)))))) \end{aligned}$$