

# thm\_2Erich\_\_list\_2EOR\_\_EL\_\_FOLDL (TM- MuKt6oxXC4s5iBsLZL3cwNKfCzaNhGgwZ)

October 26, 2020

**Definition 1** We define `c_2Emin_2E_3D` to be  $\lambda A.\lambda x \in A.\lambda y \in A.inj\_o (x = y)$  of type  $\iota \Rightarrow \iota$ .

**Definition 2** We define `c_2Ecombin_2EK` to be  $\lambda A_27a : \iota.\lambda A_27b : \iota.(\lambda V0x \in A_27a.(\lambda V1y \in A_27b.V0x))$

**Definition 3** We define `c_2Ecombin_2ES` to be  $\lambda A_27a : \iota.\lambda A_27b : \iota.\lambda A_27c : \iota.(\lambda V0f \in ((A_27c^{A_27b})^{A_27a}))$

**Definition 4** We define `c_2Ecombin_2EI` to be  $\lambda A_27a : \iota.(ap (ap (c_2Ecombin_2ES A_27a (A_27a^{A_27a})) A_27a))$

Let `ty_2Elist_2Elist` :  $\iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A0.nonempty A0 \Rightarrow nonempty (ty_2Elist_2Elist A0) \quad (1)$$

Let `c_2Elist_2EEXISTS` :  $\iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A_27a.nonempty A_27a \Rightarrow c_2Elist_2EEXISTS A_27a \in ((2^{(ty_2Elist_2Elist A_27a)})^{(2^{A_27a})}) \quad (2)$$

**Definition 5** We define `c_2Erich__list_2EOR__EL` to be  $(ap (c_2Elist_2EEXISTS 2) (c_2Ecombin_2EI 2))$ .

**Definition 6** We define `c_2Ebool_2ET` to be  $(ap (ap (c_2Emin_2E_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

**Definition 7** We define `c_2Ebool_2E_21` to be  $\lambda A_27a : \iota.(\lambda V0P \in (2^{A_27a}).(ap (ap (c_2Emin_2E_3D (2^{A_27a}))$

**Definition 8** We define `c_2Ebool_2EF` to be  $(ap (c_2Ebool_2E_21 2) (\lambda V0t \in 2.V0t))$ .

**Definition 9** We define `c_2Emin_2E_3D_3D_3E` to be  $\lambda P \in 2.\lambda Q \in 2.inj\_o (p P \Rightarrow p Q)$  of type  $\iota$ .

**Definition 10** We define `c_2Ebool_2E_5C_2F` to be  $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21 2) (\lambda V2t \in 2.V2t)))$

Let `c_2Elist_2EFOLDL` :  $\iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A_27a.nonempty A_27a \Rightarrow \forall A_27b.nonempty A_27b \Rightarrow c_2Elist_2EFOLDL A_27a A_27b \in (((A_27b^{(ty_2Elist_2Elist A_27a)})^{A_27b})^{((A_27b^{A_27a})^{A_27b})}) \quad (3)$$

Assume the following.

$$\forall A_{.27a}.nonempty\ A_{.27a} \Rightarrow (\forall V0x \in A_{.27a}.((ap\ (c_{.2Ecombin_{.2EI}}\ A_{.27a})\ V0x) = V0x)) \quad (4)$$

Assume the following.

$$\begin{aligned} \forall A_{.27a}.nonempty\ A_{.27a} \Rightarrow (\forall V0P \in (2^{A_{.27a}}).(\forall V1l \in \\ (ty_{.2Elist_{.2Elist}}\ A_{.27a}).((p\ (ap\ (ap\ (c_{.2Elist_{.2EEXISTS}}\ A_{.27a}) \\ V0P)\ V1l)) \Leftrightarrow (p\ (ap\ (ap\ (ap\ (c_{.2Elist_{.2EFOLDL}}\ A_{.27a}\ 2)\ (\lambda V2l_{.27} \in \\ 2.(\lambda V3x \in A_{.27a}.(ap\ (ap\ c_{.2Ebool_{.2E_{.5C_{.2F}}}\ V2l_{.27})\ (ap\ V0P\ V3x)))) \\ c_{.2Ebool_{.2EF}}\ V1l)))))) \end{aligned} \quad (5)$$

**Theorem 1**

$$\begin{aligned} (\forall V0l \in (ty_{.2Elist_{.2Elist}}\ 2).((p\ (ap\ c_{.2Erich_{.list_{.2EOR_{.EL}}}} \\ V0l)) \Leftrightarrow (p\ (ap\ (ap\ (ap\ (c_{.2Elist_{.2EFOLDL}}\ 2\ 2)\ c_{.2Ebool_{.2E_{.5C_{.2F}}}} \\ c_{.2Ebool_{.2EF}}\ V0l)))) \end{aligned}$$